

Facilitated transport of sodium or potassium chloride across vesicle membranes using a ditopic salt-binding macrobicyclic

Atanas V. Koulov, Joseph M. Mahoney and Bradley D. Smith*


Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556-5670, USA. E-mail: smith.115@nd.edu; Fax: +1 (574) 631 6652; Tel: +1 (574) 631 8632

Received 11th September 2002, Accepted 30th September 2002
First published as an Advance Article on the web 8th November 2002

A synthetic receptor, with an ability to bind sodium or potassium chloride as a contact ion-pair, is shown to effectively transport either salt across vesicle membranes. Significant transport is observed even when the transporter : phospholipid ratio is as low as 1 : 2500. Chloride efflux from unilamellar vesicles is monitored using a chloride selective electrode. Mechanistic studies indicate that the facilitated efflux is due to the uncomplexed transporter diffusing into the vesicle and the transporter-salt complex diffusing out. Vesicle influx experiments are also reported, where the facilitated influx of chloride and sodium ions into vesicles is observed directly by ^{35}Cl and ^{23}Na NMR, respectively.

While nature has produced mobile carrier molecules, such as valinomycin and monensin, to transport metal cations across biological membranes, it is intriguing that there are no analogous biotic carriers that selectively transport anions, or salts.¹ There are likely chemical and biological reasons for this situation. Chemically, anion transport across a bilayer membrane is less favorable than cation transport because anions are often more hydrophilic. In addition, most biological membranes have a high phospholipid content and the phosphate residues present in the phospholipid head groups compete strongly for anion binding sites. In fact our research group has recently demonstrated that various anionophores can bind and translocate certain types of phospholipids across bilayer membranes, a dynamic process also known as flip-flop.² Nonetheless, facilitated transmembrane anion transport is a ubiquitous cellular process, and thus an interesting challenge for supramolecular chemists. Biologically, the most important target anion is Cl^- since defective Cl^- transport is related to a number of disease states, the most common being cystic fibrosis.³ It is thought by some researchers that synthetic Cl^- transporters have potential as therapeutic agents.⁴ While the last few years have witnessed the first examples of synthetic Cl^- channels,⁵ we know of no publication describing rationally-designed carrier-mediated transport of Cl^- or chloride salts across phospholipid bilayer membranes.⁶ Recently, we reported the synthesis of macrocyclic receptor **1** (mp 121–123 °C)⁷ and described its ability to bind KCl or NaCl as contact ion-pairs in organic solution (Scheme 1). We now disclose that **1** is a very efficient transporter of these salts across vesicle membranes.

Cl^- efflux from unilamellar 1-palmitoyl-2-oleoylphosphatidylcholine (POPC) vesicles was monitored by a Cl^- selective electrode.⁸ The Cl^- efflux profiles in Fig. 1 show that **1** is a very effective transporter. For example, even a phospholipid:1 ratio of 2500 : 1 leads to release of half of the vesicle Cl^- content in about 300 s. Control experiments show that transporter **1** induces no leakage of entrapped fluorescent dyes or glucose-6-phosphate, reflecting the selectivity of the transport process.⁹ The importance of the ditopic salt-binding ability of macrobicyclic **1** is highlighted by the complete lack of Cl^- efflux induced by



Scheme 1 Structures of $1 \cdot \text{M}^+ \text{Cl}^-$ complex and partial ion receptors **2** and **3**.




Fig. 1 Cl^- efflux upon addition of **1** (0.4 μM , green \circ ; 4.0 μM , red ∇ ; 40.0 μM , blue \square) or a 1 : 1 molar mixture of **2** and **3** (40 μM each, blue \blacksquare) to unilamellar POPC vesicles (200 nm mean diameter, 1 mM phospholipid) containing 500 mM NaCl and dispersed in 500 mM NaHCO_3 .

high concentrations of a binary mixture of crown **2** and isophthalamide **3**, the two ion-binding components of **1**.

Mechanistic insight was gained by monitoring transporter-

promoted Cl^- efflux from POPC vesicles containing NaCl, KCl or CsCl. Furthermore, the Cl^- efflux was monitored with three different extravesicle solutions, Na_2SO_4 , NaHCO_3 and NaNO_3 (Fig. 2). Previously, we have shown that receptor **1** has a much




Fig. 2 Cl^- efflux upon addition of **1** ($4.0 \mu\text{M}$) to unilamellar POPC vesicles (200 nm mean diameter, 1 mM phospholipid) containing 500 mM of NaCl (blue \square), KCl (red ∇) or CsCl (green \circ) and dispersed in: (A) 375 mM Na_2SO_4 , (B) 500 mM NaHCO_3 , or (C) 500 mM NaNO_3 .

weaker affinity for CsCl than NaCl or KCl,¹⁰ and as shown in Fig. 2 the rates of Cl^- efflux from vesicles containing CsCl are considerably lower than the efflux from vesicles containing NaCl or KCl. This is strong evidence that the Cl^- is transported from the vesicles as a 1-salt complex.¹¹ A related mechanistic question is whether transporter **1** enters the membrane as an uncomplexed receptor, or as a salt complex. The data in Fig. 3 shows that the rate of Cl^- efflux from vesicles containing NaCl is unaltered if the extravesicle solution is changed from Na_2SO_4 to Cs_2SO_4 . The fact that Cl^- efflux rates are independent of external metal cation identity (and external anion identity, see Fig. 2) indicates that transporter **1** enters the vesicle as an uncomplexed receptor. The proposed major transport pathway for facilitated Cl^- efflux from vesicles is shown in Scheme 2.





Fig. 3 Cl^- efflux upon addition of **1** ($4.0 \mu\text{M}$) to unilamellar POPC vesicles (200 nm mean diameter, 1 mM phospholipid) containing 500 mM of NaCl and dispersed in 375 mM Na_2SO_4 (blue \square) or 375 mM Cs_2SO_4 (red ∇).



Scheme 2 Proposed mechanism for Cl^- efflux mediated by transporter **1**.

In addition to Cl^- efflux, established ^{23}Na and ^{35}Cl NMR transport assays were employed to directly observe facilitated influx of Na^+ and Cl^- ions into vesicles.¹² The influx experiments started with vesicles containing Cs_2SO_4 which were dispersed in NaCl. Both NMR assays used the same principle, that is, a membrane impermeable shift reagent was added to the vesicle solutions which allowed internalized Na^+ (or Cl^-) to be distinguished from externalized ion. The shift reagent for the ^{23}Na NMR was a DyCl_3 -sodium tripolyphosphate mixture which moves the ^{23}Na resonance upfield.¹³ As shown in Fig. 4A and B, an unshifted peak, corresponding to internalized Na^+ ,




Fig. 4 Na^+ and Cl^- influx into vesicles (egg-PC : cholesterol, 7 : 3). (A) ^{23}Na NMR spectrum of vesicles containing 150 mM Cs_2SO_4 and dispersed in 20 mM $\text{Na}_5\text{P}_3\text{O}_6$ -100 mM NaCl-5.5 mM DyCl_3 . (B) ^{23}Na NMR spectrum one hour after addition of **1** (lipid : **1**, 250 : 1). (C) ^{35}Cl NMR spectrum of vesicles containing 225 mM Cs_2SO_4 and dispersed in 300 mM NaCl-15 mM CoCl_2 . (D) ^{35}Cl NMR spectrum one hour after addition of **1** (lipid : **1**, 250 : 1).

appeared after addition of **1** to the vesicles. The ^{35}Cl NMR shift reagent was CoCl_2 which moves the broadened ^{35}Cl resonance downfield.^{14,15} As shown in Fig. 4C and D, an unshifted peak, corresponding to internalized Cl^- , appeared after addition of **1** to the vesicles.

In summary, a salt-binding macrobicyclic is shown for the first time to transport NaCl or KCl across vesicle membranes. The ditopic receptor **1** is an extremely effective transporter, whereas a binary mixture of crown **2** and isophthalamide **3**, the two ion-binding components of **1**, has essentially no transport activity. Our results suggest that salt transporters, such as **1**, are likely to induce interesting biological effects.

Acknowledgements

This work was supported by the National Science Foundation (USA).

References

- 1 A very rare group of chloride transporters are the prodigiosins. See for example: K. Tanigaki, T. Sato, Y. Tanaka, T. Ochi, A. Nishikawa, K. Nagai, H. Kawashima and S. Ohkuma, *FEBS Lett.*, 2002, **524**, 37–42.
- 2 J. M. Boon and B. D. Smith, *J. Am. Chem. Soc.*, 2001, **123**, 6221–6226.
- 3 B. J. Rosenstein and P. L. Zeitlin, *Lancet*, 1998, **351**, 277–282.
- 4 (a) L. Gao, J. R. Broughman, T. Iwamoto, J. M. Tomich, C. J. Venglarik and H. J. Forman, *Am. J. Physiol. Lung Cell Mol. Physiol.*, 2001, **281**, L24–L30; (b) C. Jiang, E. R. Lee, M. B. Lane, X.-F. Xiao, D. J. Harris and S. H. Cheng, *Am. J. Physiol. Lung Cell Mol. Physiol.*, 2001, **281**, L1164–L1172; (c) H. C. Rodgers and A. J. Knox, *Eur. Respir. J.*, 2001, **17**, 1314–1321.
- 5 (a) L. A. Weiss, N. Sakai, B. Ghebremariam, C. Ni and S. Matile, *J. Am. Chem. Soc.*, 1997, **119**, 12142–12149; (b) M. Merritt, M. Lanier, G. Deng and S. L. Regen, *J. Am. Chem. Soc.*, 1998, **120**, 8494–8501; (c) V. Sidorov, F. W. Kotch, G. Abdrakhmanova, R. Mizani, J. C. Fettingner and J. T. Davis, *J. Am. Chem. Soc.*, 2002, **124**, 2267–2278; (d) P. H. Schlesinger, R. Ferdani, J. Liu, J. Pajewska, R. Pajewski, M. Saito, H. Shabany and G. W. Gokel, *J. Am. Chem. Soc.*, 2002, **124**, 1848–1849.
- 6 There are a number of papers in the literature describing facilitated anion or salt transport across liquid organic membranes. See, for example: (a) J. L. Sessler, P. I. Sansom, A. Andrievsky and V. Kral, in *Supramolecular Chemistry of Anions*, eds. A. Bianchi, K. Bowman-James and E. Garcia-Espana, Wiley, New York, 1997; (b) L. A. J. Chrisstoffels, D. N. Reinhoudt, S. Sivelli, L. Gazzola, A. Casnati and R. Ungaro, *J. Am. Chem. Soc.*, 1999, **121**, 10142–10151; (c) P. K. Beer and J. D. McKinney, *Chem. Commun.*, 1999, 1253–1254; (d) D. M. Rudkevich, W. Verboom, R. Ungaro, F. de Jong and D. N. Reinhoudt, *J. Am. Chem. Soc.*, 1995, **117**, 6124–6125; (e) H. Tsukube, *Tetrahedron Lett.*, 1981, **22**, 3981–3984.
- 7 J. M. Mahoney, A. M. Beatty and B. D. Smith, *J. Am. Chem. Soc.*, 2001, **123**, 5847–5848.
- 8 Briefly, unilamellar vesicles (mean diameter of 200 nm) containing 500 mM of salt were prepared by freeze–thawing, followed by extrusion through polycarbonate membranes. The non-encapsulated salt was removed by dialysis. The chloride efflux experiments used a chloride selective electrode (Accumet) and 5 mL vesicle solutions (1 mM phospholipid). The initial chloride reading was considered as 0% release and the reading after detergent addition at 300 s considered as 100% release. Typical chloride concentrations were 6.0×10^{-5} M in the beginning and 6.0×10^{-4} M after 100% release.
- 9 P. R. Westmark, S. J. Gardiner and B. D. Smith, *J. Am. Chem. Soc.*, 1996, **118**, 11093–11100.
- 10 J. M. Mahoney and B. D. Smith, unpublished results.
- 11 Additional experiments show that Cl^- efflux is very slow from vesicles containing 500 mM CsCl and dispersed in 500 mM CsNO_3 (e.g., 4.0 μM of **1** induces 10% Cl^- release after 300 s) which is strong evidence against a simple anion antiport mechanism.
- 12 (a) F. G. Riddell in *Encyclopedia of Spectroscopy and Spectrometry*, eds. J. C. Lindon, G. E. Tranter and J. L. Holmes, Academic Press, London, 1999, 1584–1593; (b) P. W. Kuchel, B. E. Chapman, and A. S. L. Xu, in *Progress in Cell Research*, vol. 2, eds. E. Bamberg and H. Passow, Elsevier, Amsterdam, 1992.
- 13 Q. Xie, Y. Li, G. Gokel, J. Hernandez and L. Echevoyen, *J. Am. Chem. Soc.*, 1994, **116**, 690–696.
- 14 Y. Shachar-Hill and R. Shulman, *Biochemistry*, 1992, **31**, 6272–6278.
- 15 In our hands, CoCl_2 disrupts vesicles composed of only POPC. Therefore, the NMR-based transport assays used vesicles composed of egg-PC : cholesterol (7 : 3) which were mechanically stable in the presence of CoCl_2 .