

ON TAMING AND COMPATIBLE SYMPLECTIC FORMS

RICHARD HIND, COSTANTINO MEDORI, ADRIANO TOMASSINI

ABSTRACT. Let (X, J) be an almost complex manifold. The almost complex structure J acts on the space of 2-forms on X as an involution. A 2-form α is J -anti-invariant if $J\alpha = -\alpha$. We investigate the anti-invariant forms and their relation to taming and compatible symplectic forms. For every closed almost complex manifold, in contrast to invariant forms, we show that the space of closed anti-invariant forms has finite dimension.

If X is a closed almost-complex manifold with a taming symplectic form then we show that there are no non trivial exact anti-invariant forms. On the other hand we construct many examples of almost-complex manifolds with exact anti-invariant forms, which are therefore not tamed by any symplectic form. In particular we use our analysis to give an explicit example of an almost-complex structure which is locally almost-Kähler but not globally tamed.

The non-existence of exact anti-invariant forms however does not in itself imply that there exists a taming symplectic form. We show how to construct examples in all dimensions.

INTRODUCTION

Almost-complex structures on a manifold X can be categorized according to whether or not there exist taming or compatible symplectic forms. We recall that a symplectic form ω tames an almost-complex structure J if $\omega(v, Jv) > 0$ for all nonzero tangent vectors v , and ω is compatible with J if the formula $g(v, w) = \omega(v, Jw)$ defines a Riemannian metric g on X . If ω is compatible with J then the triple (X, J, ω) is sometimes called an almost-Kähler manifold. A Kähler manifold is an almost-Kähler manifold with J integrable.

Let $\mathcal{J} = \mathcal{J}(X)$ be the set of almost-complex structures on X , then we can define subsets $\mathcal{J}_{tame} = \mathcal{J}_{tame}(X)$ and $\mathcal{J}_{comp} = \mathcal{J}_{comp}(X)$ of \mathcal{J} to be the almost-complex structures for which there exists a taming or compatible symplectic form respectively. We also define a subset $\mathcal{J}_{loc.tame}$ of \mathcal{J}_{tame} which consists of locally tame almost-complex structures, that is, $J \in \mathcal{J}_{loc.tame}$ if there exists an open cover $\{U_i\}$ of X such that $J|_{U_i} \in \mathcal{J}_{tame}(U_i)$ for all i . Similarly we can define locally compatible almost-complex structures $\mathcal{J}_{loc.comp} \subset \mathcal{J}_{comp}$. It is immediate that $\mathcal{J}_{loc.tame} = \mathcal{J}$ and that the set of (integrable) complex structures $\mathcal{I} \subset \mathcal{J}_{loc.comp}$.

In summary we have the following diagram of inclusions.

Date: April 16, 2014.

2010 *Mathematics Subject Classification.* 32Q60, 53C15, 58A12.

Key words and phrases. pure and full almost complex structure; J -invariant form; J -anti-invariant form. Partially supported by *Fondazione Bruno Kessler-CIRM (Trento)* and by GNSAGA of INdAM.

$$(1) \quad \mathcal{I} \subset \begin{array}{c} \mathcal{J}_{loc.comp} \\ \bigcup_{k_1} \\ \mathcal{J}_{comp} \end{array} \subset_j \begin{array}{c} \mathcal{J}_{loc.tame} \\ \bigcup_{k_2} \\ \mathcal{J}_{tame} \end{array} = \mathcal{J}$$

When our manifold X has dimension 4 the map j is actually a surjection, in other words $\mathcal{J} = \mathcal{J}_{loc.comp}$. For a complete proof of this see Lejmi [13, Theorem 1]. It is a question of Donaldson [4, question 2] as to whether the map i is also a surjection.

In this paper a key observation is the following.

Proposition 0.1. *If X is closed (compact without boundary) and $J \in \mathcal{J}_{tame}$ then there are no non-zero exact J -anti-invariant 2 forms.*

Recall that a 2-form α is anti-invariant if $J\alpha = -\alpha$, where $J\alpha(v, w) = \alpha(Jv, Jw)$. In dimension 4 there are no non-zero exact anti-invariant forms with respect to any J , see Corollary 1.2, but in higher dimensions the existence of an exact anti-invariant form is an obstruction to the existence of a taming symplectic form.

It is in fact quite easy to find examples of almost-complex structures admitting exact anti-invariant forms. The following is a consequence of Theorem 1.4.

Theorem 0.2. *Suppose that W^{4n} is a $4n$ dimensional manifold with trivial tangent bundle. Then $X = W \times S^1 \times S^1$ has an almost-complex structure J for which there exist non-zero exact anti-invariant 2-forms.*

The methods used to establish Theorem 0.2 are very topological, they rely on Gromov's h-principle. Therefore we have little control on the almost-complex structure, in particular it is difficult in this way to find examples which lie in $\mathcal{J}_{loc.comp}$. This issue is addressed in section 3. For example, in section 3.3 we explicitly construct a nonintegrable almost-complex structure on a 6-dimensional manifold which is locally compatible yet admits a non-zero exact anti-invariant form, and so lies in $\mathcal{J}_{loc.comp} \setminus (\mathcal{J}_{tame} \cup \mathcal{I})$.

We remark however that the non-existence of exact anti-invariant forms is not a sufficient condition for an almost-complex structure to be tamed by a symplectic form. In dimension 4, since we never have any exact anti-invariant forms, examples are given by any almost-complex manifolds which are not symplectic. In higher dimensions, we can use a theorem of Peternell [17, Theorem 1.4] to imply the following.

Theorem 0.3. *A non-Kähler Moisëzon manifold has no non-zero exact anti-invariant forms but no taming symplectic form.*

In dimension 6, a simpler concrete example is the following.

Theorem 0.4. *The product of the Hopf surface and $\mathbb{C}P^1$ does not have non-zero exact anti-invariant forms or any symplectic forms at all.*

Cohomology properties can also be used to categorize almost-complex structures. Following [5] we can define subspaces $H_J^+(X), H_J^-(X) \subset H^2(X, \mathbb{R})$, the second de Rham cohomology of X , as follows. A class $\mathbf{a} \in H_J^+(X)$ if there exists a 2 form α with $[\alpha] = \mathbf{a}$ and $J\alpha = \alpha$. Similarly a class $\mathbf{a} \in H_J^-(X)$ if it has a representative α which is anti-invariant with respect to J . The almost-complex manifold (X, J) is called \mathcal{C}^∞ -pure if

$H_J^+(X) \cap H_J^-(X) = \{0\}$ and \mathcal{C}^∞ -full if $H_J^+(X) + H_J^-(X) = H^2(X, \mathbb{R})$. In [5, Theorem 2.3], Drahjici, Li and Zhang show that an almost complex structure on a compact 4-dimensional manifold is \mathcal{C}^∞ -pure-and-full. Furthermore, in [14, Theorem 1.3], Li and Zhang proved that if J is \mathcal{C}^∞ -full and if the compatible cone

$$\mathcal{K}_J^c = \{[\omega] \in H^2(X; \mathbb{R}) \mid \omega \text{ is compatible with } J\}$$

is non-empty, then

$$\mathcal{K}_J^t = \mathcal{K}_J^c + H_J^-(X).$$

Here we focus on $H_J^-(X)$ and study $\mathcal{Z}_J^-(X)$, the real vector space of closed anti-invariant 2-forms. We have already seen that if X is closed and $J \in \mathcal{J}_{tame}$ then there are no nonzero exact anti-invariant forms and so the map

$$\mathcal{Z}_J^-(X) \rightarrow H_J^-(X) \subset H^2(X, \mathbb{R})$$

is an injection. This can be contrasted with the case of invariant forms $\mathcal{Z}_J^+(X)$. At least if J is integrable then $\mathcal{Z}_J^+(X)$ is always infinite dimensional.

In the case when $J \in \mathcal{J}_{comp}$ we can be more precise. Let g be the Riemannian metric associated to a compatible symplectic form. Then we show in Proposition 2.2 that $\mathcal{Z}_J^-(X) \subset \mathcal{H}_g(X)$, the set of harmonic 2-forms with respect to g . In other words, we have the following.

Proposition 0.5. *J -anti-invariant forms are harmonic with respect to any Riemannian metric associated to a compatible symplectic form.*

It turns out that even if $J \notin \mathcal{J}_{comp}$ the closed anti-invariant forms \mathcal{Z}_J^- lie in the kernel of a second order elliptic operator, see Proposition 2.4. Hence we have an alternative proof of a theorem from [10] saying that anti-invariant forms satisfy a unique continuation principle, see Proposition 2.6.

The paper is organized as follows. After fixing some notation, we establish some basic facts about anti-invariant forms in section 1 and prove Proposition 0.1 and Theorem 0.2. To complement these results we also derive the examples of Theorems 0.3 and 0.4. In section 2 we discuss the relation between anti-invariant forms and harmonic forms and in particular prove the unique continuation theorem for anti-invariant forms. Finally in section 3 we construct our explicit examples.

Acknowledgements. We would like to thank *Fondazione Bruno Kessler-CIRM (Trento)* for their support and very pleasant working environment.

1. ANTI-INVARIANT FORMS ON ALMOST-COMPLEX MANIFOLDS.

We start by fixing some notation. Let (X, J) be a $2n$ -dimensional almost complex manifold and denote by $\Lambda^2(X)$ the space of 2-forms on X . The almost complex structure acts on $\Lambda^2(X)$ as an involution by setting $J\alpha(u, v) = \alpha(Ju, Jv)$. Following [5], $\alpha \in \Lambda^2(X)$ is said to be J -invariant or invariant, if $J\alpha = \alpha$ and J -anti-invariant or anti-invariant if $J\alpha = -\alpha$. We denote by $\Lambda_J^+(X)$ and by $\Lambda_J^-(X)$ the space of J -invariant, J -anti-invariant forms respectively. Let $\mathcal{Z}(X)$ be the space of closed 2-forms on X . We set $\mathcal{Z}_J^\pm(X) = \Lambda_J^\pm(X) \cap \mathcal{Z}(X)$ and

$$H_J^\pm(X) = \{\mathbf{a} \in H^2(X; \mathbb{R}) \mid \mathbf{a} = [\alpha], \alpha \in \mathcal{Z}_J^\pm(X)\}.$$

According to [5], an almost complex structure J is said to be C^∞ -pure if $H_J^+(X) \cap H_J^-(X) = \{0\}$, C^∞ -full if $H^2(X; \mathbb{R}) = H_J^+(X) + H_J^-(X)$.

We begin our study of anti-invariant forms in dimension 4.

Lemma 1.1. *Let (X, J, g) be a 4-dimensional almost Hermitian manifold. Let $\alpha \in \Lambda^2_-(X)$. Then $\alpha^2 = f \text{Vol}_g$, where $f : X \rightarrow \mathbb{R}$ is a smooth non-negative function on X and Vol_g denotes the Riemannian volume form.*

Proof. Let $p \in X$ and let $\{v_1, Jv_1, v_2, Jv_2\}$ be a g -orthonormal positive basis of $T_p X$. Then, if $\{v_1^*, Jv_1^*, v_2^*, Jv_2^*\}$ denotes the dual basis of $\{v_1, Jv_1, v_2, Jv_2\}$, any J -anti-invariant 2-form at p can be written as

$$\alpha(p) = \lambda(v_1^* \wedge v_2^* - Jv_1^* \wedge Jv_2^*) + \mu(v_1^* \wedge Jv_2^* + Jv_1^* \wedge v_2^*).$$

Hence,

$$\alpha^2(p) = 2(\lambda^2 + \mu^2)(v_1^* \wedge Jv_1^* \wedge v_2^* \wedge Jv_2^*).$$

□

Corollary 1.2. *Let (X, J, g) be a compact 4-dimensional almost Hermitian manifold. Then there are no non-trivial exact anti-invariant forms on X .*

Proof. Let $\alpha \in \mathcal{Z}_J^-(X)$. By assumption $\alpha \neq 0$; assume by contradiction that $\alpha = d\beta$. Then,

$$0 = \int_X d(\beta \wedge d\beta) = \int_X \alpha^2 = \int_X f \text{Vol}_g > 0,$$

and this is absurd. □

We will see in Theorem 1.4 and in section 3 that non-zero exact anti-invariant forms can exist on higher dimensional almost-complex manifolds, but the following proposition rules this out if the almost-complex structure $J \in \mathcal{J}_{tame}$.

Proposition 1.3. *Let (X, J) be a compact $2n$ -dimensional almost complex manifold. If ω is a symplectic form taming J , then there are no non-zero exact J -anti-invariant forms.*

Proof. By contradiction. Let ω be a symplectic form taming J . Let $\alpha \in \mathcal{Z}_J^-(X)$ be exact, $\alpha = d\beta$. Let $2k$ be the maximal rank of $\alpha(p)$, for $p \in X$. The following claim from linear algebra is useful.

Claim. A skew-symmetric anti-invariant 2-form η on a complex vector space V has rank $2r$ divisible by 4. Moreover η^r generates the complex orientation on $V/\ker \eta$ (with its induced complex structure) if $r/2$ is even and the opposite of the complex orientation if $r/2$ is odd.

Proof of claim. First note that as η is anti-invariant $\ker(\eta)$ is indeed a complex subspace of V and so the quotient $W = V/\ker(\eta)$ inherits a complex structure. Then η^r is a volume form on W which implies that $J\eta^r = \lambda\eta^r$ for some $\lambda > 0$. As η is anti-invariant this in turn implies that r must be even and we can take $\lambda = 1$.

Now we choose a basis of W of the form $e_1, f_1, e_2, f_2, \dots, e_r, f_r$ such that if i is odd we have $\eta(e_i, f_i) = 1$ and $Je_i = e_{i+1}$ and $Jf_i = f_{i+1}$. Then necessarily if i is even we have $\eta(e_i, f_i) = -1$. We may also assume that $\eta(e_i, e_j) = \eta(f_i, f_j) = 0$ for all i, j and $\eta(e_i, f_j) = 0$ for all $i \neq j$.

Given this we compute

$$\eta^r(e_1, e_2, f_1, f_2, \dots, f_{r-1}, f_r) = (-1)^r \eta(e_1, f_1) \eta(e_2, f_2) \dots \eta(e_r, f_r) = (-1)^{r/2}$$

and the claim follows. \square

Returning to the proof, we have that k is even and $\alpha^k \wedge \omega^{n-k} = (-1)^{k/2} f \omega^n$, where f is a non-negative function on X . This is because ω gives the complex orientation on any complex subspace. The function f is positive exactly when α has maximum rank. Therefore

$$0 = (-1)^{k/2} \int_X d(\beta \wedge (d\beta)^{k-1} \wedge \omega^{n-k}) = (-1)^{k/2} \int_X \alpha^k \wedge \omega^{n-k} = \int_X f \omega^n > 0$$

and this is absurd. \square

To complement the above proposition, the following theorem shows that almost-complex structures admitting exact anti-invariant forms can be constructed under fairly general hypotheses.

Theorem 1.4. *Suppose that an orientable manifold M^{4n+1} admits a 2 form $\tilde{\alpha}$ of everywhere maximal rank $4n$ such that the quotient bundle $TM/\ker \tilde{\alpha} \rightarrow M$ has an almost-complex structure for which $\tilde{\alpha}$ is anti-invariant. Then there exists an almost-complex structure J on $M \times S^1$ which admits an exact nonzero anti-invariant 2 form.*

We emphasize that the hypotheses of the theorem are purely topological, in particular we do not need to assume that $\tilde{\alpha}$ is closed. The proof does not use the hypothesis that M has dimension $4n + 1$, only that the dimension is odd. However we have seen above that the rank of an anti-invariant form is necessarily a multiple of 4.

Proof. The result is a consequence of a theorem of McDuff, see [15] and [7, Thm 10.4.1], which states (in a simple form) that a 2-form of maximal rank on an odd dimensional manifold can be deformed through forms of maximal rank to an exact form. Hence we can find maximal rank 2-forms α_t on M such that $\alpha_0 = \tilde{\alpha}$ and α_1 is exact. Fixing a Riemannian metric on M the $4n$ dimensional subbundles $(\ker \tilde{\alpha})^\perp$ and $(\ker \alpha_1)^\perp$ are isomorphic as symplectic vector bundles with forms $\tilde{\alpha}$ and α_1 respectively. Hence $(\ker \alpha_1)^\perp$ also admits an almost-complex structure J anti-invariant with respect to α_1 . The corresponding orientation on $(\ker \alpha_1)^\perp$ together with one on M determines a trivialization of $\ker \alpha$. Hence we can extend J to an almost-complex structure on $M \times S^1$ such that J maps $\ker \alpha_1$ onto TS^1 . Let us pull back α_1 to a 2-form α on $M \times S^1$ using the natural projection. Then α is also nonzero and exact. Finally since α vanishes on the complex planes spanned by $\ker \alpha_1$ and TS^1 it is anti-invariant as required. \square

To close this section we discuss our examples of complex manifolds which have no exact anti-invariant 2-forms but still have no taming symplectic forms.

First let X be a Moisézon manifold, that is, a compact complex manifold which admits a proper modification from a projective manifold. Then the following result holds, see Peternell [17, Thm.1.4]

Theorem 1.5. *Let X be a Moisézon manifold. Assume there exists a real $(1, 1)$ -form ω and a real 2-form φ on X such that*

- i) ω is positive definite,

- ii) $d(\omega - \varphi) = 0$,
- iii) $\int_C \varphi = 0$ for all curves $C \subset X$.

Then X is projective.

This directly implies Theorem 0.3 as follows.

Proposition 1.6. *Any non-Kähler Moisëzon manifold X has no non-trivial d -exact anti-invariant 2-forms and no taming symplectic forms.*

Proof. First, if α is a d -exact anti-invariant 2-form α on X then its pull back to a projective manifold is also exact and anti-invariant. By Proposition 1.3 this implies that α must be identically zero.

The fact that X has no taming symplectic form has already been pointed out by Draghici and Zhang, [6], but we give the argument here for completeness. Arguing by contradiction, suppose that η is a taming symplectic form. We can write $\eta = \omega - \psi_1 - \bar{\psi}_2$ where ω is a real $(1, 1)$ -form, ψ_1 is a real $(2, 0)$ -form and $\bar{\psi}_2$ is a real $(0, 2)$ -form. Then ψ_1 and $\bar{\psi}_2$ vanish on complex lines and so since η is taming the form ω is positive definite. Setting $\varphi = \psi_1 + \bar{\psi}_2$ the remaining two conditions of Theorem 1.5 are clearly satisfied and so X must be projective, a contradiction. \square

Finally we give a proof of Theorem 0.4. Let Y be the Hopf surface, that is, $Y = (\mathbb{C}^2 \setminus 0) / z \sim 2z$ with its induced complex structure.

Proposition 1.7. *The product $X = \mathbb{C}P^1 \times Y$ does not have exact anti-invariant forms or any symplectic forms at all.*

Proof. The 6-manifold X is diffeomorphic to $S^2 \times S^3 \times S^1$ and so has no cohomology classes \mathfrak{a} with $\mathfrak{a}^3 \neq 0$. Therefore it admits no symplectic forms at all.

There are two projections $p_1, p_2 : X \rightarrow \mathbb{C}P^1$. The first is just projection onto the first factor, the second is induced by projection onto Y and then quotienting by \mathbb{C}^* to get $Y/\mathbb{C}^* = (\mathbb{C}^2 \setminus 0)/\mathbb{C}^* = \mathbb{C}P^1$. Therefore we can pull-back the Fubini-Study form using p_1 and p_2 to get invariant 2-forms ω_1 and ω_2 on X .

Suppose that there exists a non-zero exact anti-invariant 2-form α on X . As we are working in dimension 6 we have that $\alpha(x)$ has rank 0 or 4 at all points $x \in X$. Observe that applying Stokes' Theorem as in Proposition 1.3 gives a contradiction if there exists a closed 2-form Ω on X which satisfies $\alpha^2 \wedge \Omega \geq 0$ and $\alpha^2 \wedge \Omega(x) > 0$ at least for some $x \in X$. When $\alpha \neq 0$ its kernel is a complex line. Therefore as ω_1 and ω_2 are invariant we have $\alpha^2 \wedge (\omega_1 + \omega_2) \geq 0$ (for the complex orientation on X) and hence by the argument above we must have $\alpha^2 \wedge (\omega_1 + \omega_2) \equiv 0$.

This implies that when $\alpha(x) \neq 0$ its kernel is generated by r and ir , where r is the radial, or S^1 , direction in Y (coming from a suitably scaled radial vector in \mathbb{C}^2) and ir is parallel to the Hopf fibration. Indeed, if the kernel were transverse to this plane the form $\omega_1 + \omega_2$ would evaluate nontrivially. Hence r and ir lie in $\ker(\alpha(x))$ for all $x \in X$ and α is invariant under the vectorfields r and ir . These vectorfields generate a torus action on X whose projection onto the orbit space is just the holomorphic projection $(p_1, p_2) : X \rightarrow \mathbb{C}P^1 \times \mathbb{C}P^1$. Hence α is a pull-back of a form α' on $\mathbb{C}P^1 \times \mathbb{C}P^1$. As α is a closed anti-invariant form so is α' . Furthermore, as α' is anti-invariant it must vanish when restricted to both $\mathbb{C}P^1$ factors. Therefore its cohomology class is trivial and so

α' is exact. But by Corollary 1.2 the only exact anti-invariant forms on a 4-dimensional manifold are identically 0, and this completes our proof. \square

2. HODGE STAR OPERATOR FOR ANTI-INVARIANT FORMS

Let (X, J, g) be an almost Hermitian manifold of dimension $2n$ and denote by ω the fundamental form of g . Then we have the following

Proposition 2.1. *Let α be J -anti-invariant 2-form on (X, J, g) . Then*

$$(2) \quad *\alpha = \frac{1}{(n-2)!} \alpha \wedge \omega^{n-2}.$$

For the sake of completeness we give the proof of (2).

Proof. Let α be any J -anti-invariant form on (X, J) . Then $*\alpha$ is a $(2n-2)$ -form. The Lefschetz decomposition applied to $\Lambda^{2n-2}(X)$ yields to

$$\Lambda^{2n-2}(X) = \bigoplus_{i \geq 0} L^i(P^{2(n-i)-2}(X)),$$

where $L : \Lambda^k(X) \rightarrow \Lambda^{k+2}$, $L(\gamma) = \gamma \wedge \omega$ is the Lefschetz operator and $P^k(X)$ is the space of primitive forms, which can be identified with $\ker L^{n-k+1}|_{\Lambda^k(X)}$ (see e.g., [11, Prop.1.2.30]). Therefore,

$$*\alpha = f\omega^{n-1} + L^{n-2}(\gamma),$$

where f is a smooth function and $\gamma \in P^2(X)$. Then, taking $*$ in the last formula, by [11, Prop.1.2.30], we get

$$\alpha = f\omega - (n-2)!J\gamma.$$

Since α is J -anti-invariant, by the last formula, $f = 0$ and $\gamma = \frac{1}{(n-2)!}\alpha$. Then (2) is proved. \square

As a consequence, we obtain the following

Proposition 2.2. *Let (X, J, g, ω) be a $2n$ -dimensional almost Kähler manifold. Then $\mathcal{Z}_J^-(X) \subset \mathcal{H}^2(X)$, where $\mathcal{H}^2(X)$ denotes the space of 2-harmonic forms on X with respect to the Hermitian metric g .*

Proof. Let $\alpha \in \mathcal{Z}_J^-(X)$. Then by formula (2), since α and ω are closed, we get:

$$d^*\alpha = - * d * (\alpha) = - \frac{1}{(n-2)!} * d(\alpha \wedge \omega^{n-2}) = 0,$$

that is α co-closed. Since α is closed by assumption, then α is harmonic. \square

We record the following corollary, which of course also follows from Proposition 1.3.

Corollary 2.3. *If (X, J, g, ω) is a compact $2n$ -dimensional almost Kähler manifold, then the natural map*

$$\mathcal{Z}_J^-(X) \hookrightarrow H_{dR}^2(X; \mathbb{R})$$

is an injection. In particular $\dim_{\mathbb{R}}(\mathcal{Z}_J^-(X)) \leq b_2(X)$. Furthermore, the map is an isomorphism if and only if $H_J^-(X) = H_{dR}^2(X; \mathbb{R})$.

In general, on an almost Hermitian manifold (X, J, g) of dimension $2n$, define a generalized co-differential on the space of 2-forms $\Gamma(\Lambda^2(X))$, $d_-^* : \Gamma(\Lambda^2(X)) \rightarrow \Gamma(\Lambda^1(X))$, by setting

$$d_-^*(\alpha) = d^*(\alpha) + \frac{1}{(n-2)!} *(\alpha \wedge d(\omega^{n-2})),$$

where d^* denotes the usual co-differential on (X, g) . By formula (2), it follows that d_-^* vanishes on $\mathcal{Z}_-^2(X)$. Let E be the differential operator on $\Gamma(\Lambda^2(X))$ defined as

$$\mathbb{E} = \Delta(\alpha) + \frac{1}{(n-2)!} d(*(\alpha \wedge d(\omega^{n-2})))$$

Proposition 2.4. *The differential operator \mathbb{E} is a second order elliptic operator, the principal part is the Hodge-de Rham laplacian Δ and $\mathcal{Z}_J^-(X) \subset \ker(\mathbb{E})$.*

Proof. By the definition of \mathbb{E} , for any $\alpha \in \mathcal{Z}_J^-(X)$, we have:

$$\begin{aligned} \mathbb{E}(\alpha) &= \Delta(\alpha) + \frac{1}{(n-2)!} d(*(\alpha \wedge d(\omega^{n-2}))) = dd^*(\alpha) + d^*d(\alpha) + \\ &\quad \frac{1}{(n-2)!} d(*(\alpha \wedge d(\omega^{n-2}))) \\ &= dd^*(\alpha) + \frac{1}{(n-2)!} d(*(\alpha \wedge d(\omega^{n-2}))) = dd_-^*(\alpha) = 0. \end{aligned}$$

□

Corollary 2.5. *If (X, J) is a compact $2n$ -dimensional almost complex manifold, then $\dim \mathcal{Z}_J^-(X) < +\infty$.*

In contrast, $\mathcal{Z}_J^+(X)$ has infinite dimension if J is integrable, because for any smooth function $f : X \rightarrow \mathbb{R}$ we have $dd^c f \in \mathcal{Z}_J^+(X)$.

We can now give another proof of the analytic continuation property for closed anti-invariant 2-forms (see [10, Thm.4.1])

Proposition 2.6. *Let X be a $2n$ -dimensional connected almost complex manifold. Let $\alpha \in \mathcal{Z}_J^-(X)$ be vanishing at infinite order at some point $p \in X$. Then α is identically zero.*

Proof. By Proposition 2.4, α is a solution of an elliptic PDE, whose leading term is the Laplacian. Hence by [1] (see also [12]), the form α has strong unique continuation. □

In contrast, this is false for $\mathcal{Z}_J^+(X)$ for the same reason as before.

3. COMPUTATIONS OF $\mathcal{Z}_-^2(X)$

In this section we will do some explicit computations on the space of anti-invariant forms on complex manifolds, to contrast with the indirect Theorem 1.4. In section 3.1 we give an example of a complex manifold with $\dim_{\mathbb{R}} \mathcal{Z}_J^-(X) > \dim_{\mathbb{R}} H_J^-(X)$. By Corollary 2.3 this implies that the manifold is not almost Kähler. Indeed by Proposition 1.3 there is not even a taming symplectic form. Another such example is given in section 3.2. Finally in section 3.3 we construct an almost-complex manifold for which we can write down explicitly a compatible symplectic form on small open sets. However it also admits a non-zero exact anti-invariant form and so by Proposition 1.3 has no globally defined taming symplectic form.

3.1. Iwasawa manifold. On \mathbb{C}^3 , consider the product $*$ defined as

$$(z_1, z_2, z_3) * (w_1, w_2, w_3) = (z_1 + w_1, z_2 + w_2, z_3 + z_1 w_2 + w_3) .$$

It is immediate to check that $(\mathbb{C}^3, *)$ is a nilpotent Lie group isomorphic to

$$\mathbb{H}(3) = \left\{ \begin{pmatrix} 1 & z_1 & z_3 \\ 0 & 1 & z_2 \\ 0 & 0 & 1 \end{pmatrix} \in GL(3; \mathbb{C}) \mid z_1, z_2, z_3 \in \mathbb{C} \right\} .$$

We have that $(\mathbb{Z}[i])^3 \subset \mathbb{C}^3$ is a cocompact discrete subgroup of $(\mathbb{C}^3, *)$. The *Iwasawa manifold* X is defined as the manifold

$$X = (\mathbb{Z}[i])^3 \backslash (\mathbb{C}^3, *) .$$

It is a compact complex 3-dimensional nilmanifold; by [8], it follows that X is not formal; hence, it has no Kähler metrics, see [3, Main Theorem]; nevertheless, there exists a balanced metric on X . Let $(z^i)_{i \in \{1,2,3\}}$ be the standard complex coordinate system on \mathbb{C}^3 ; the following $(1, 0)$ -forms on \mathbb{C}^3 are invariant for the action (on the left) of $(\mathbb{Z}[i])^3$, so they give rise to a global coframe for $T^{*1,0}X$:

$$\begin{cases} \varphi^1 = dz^1, \\ \varphi^2 = dz^2, \\ \varphi^3 = dz^3 - z^1 dz^2. \end{cases}$$

The structure equations are therefore

$$\begin{cases} d\varphi^1 = 0, \\ d\varphi^2 = 0, \\ d\varphi^3 = -\varphi^1 \wedge \varphi^2. \end{cases}$$

By Hattori-Nomizu theorem, we compute the real cohomology group $H_{dR}^2(X; \mathbb{R})$ of X (for simplicity, we list the harmonic representative instead of its class and write φ^{AB} for $\varphi^A \wedge \varphi^B$):

$$\begin{aligned} H_{dR}^2(X; \mathbb{R}) = \text{span}_{\mathbb{R}} \{ & \varphi^{13} + \varphi^{\bar{1}\bar{3}}, i(\varphi^{13} - \varphi^{\bar{1}\bar{3}}), \varphi^{23} + \varphi^{\bar{2}\bar{3}}, \\ & i(\varphi^{23} - \varphi^{\bar{2}\bar{3}}), \varphi^{1\bar{2}} - \varphi^{2\bar{1}}, i(\varphi^{1\bar{2}} + \varphi^{2\bar{1}}), i\varphi^{1\bar{1}}, i\varphi^{2\bar{2}} \} , \end{aligned}$$

Note that each harmonic representative is of pure degree and hence the complex structure is \mathcal{C}^∞ -pure and full. The Betti numbers of X are

$$b^0 = 1, \quad b^1 = 4, \quad b^2 = 8, \quad b^3 = 10 .$$

Then

$$\begin{aligned} & \frac{1}{2}(\varphi^2 \wedge \varphi^3 + \bar{\varphi}^2 \wedge \bar{\varphi}^3), \frac{1}{2i}(\varphi^2 \wedge \varphi^3 - \bar{\varphi}^2 \wedge \bar{\varphi}^3), \frac{1}{2}(\varphi^1 \wedge \varphi^2 + \bar{\varphi}^1 \wedge \bar{\varphi}^2), \\ & \frac{1}{2i}(\varphi^1 \wedge \varphi^2 - \bar{\varphi}^1 \wedge \bar{\varphi}^2), \frac{1}{2}(\varphi^1 \wedge \varphi^3 + \bar{\varphi}^1 \wedge \bar{\varphi}^3), \frac{1}{2i}(\varphi^1 \wedge \varphi^3 - \bar{\varphi}^1 \wedge \bar{\varphi}^3), \end{aligned}$$

are J -anti-invariant closed 2-forms on X and consequently $\dim_{\mathbb{R}} \mathcal{Z}_J^-(X) > \dim_{\mathbb{R}} H_J^-(X)$.

3.2. Nakamura manifold. The *Nakamura manifold* is the compact quotient $X = \Gamma \backslash G$ of G by a uniform discrete subgroup Γ .

By [2, Corollary 4.2] we have

$$H_{dR}^2(X; \mathbb{R}) = \text{span}_{\mathbb{R}} \left\{ [e^{14}], [e^{26} - e^{35}], [e^{23} - e^{56}], [\cos(2x_4)(e^{23} + e^{56}) - \sin(2x_4)(e^{26} + e^{35})], \right. \\ \left. [\sin(2x_4)(e^{23} + e^{56}) - \cos(2x_4)(e^{26} + e^{35})] \right\},$$

i.e. in this case the de Rham cohomology of M is not isomorphic to $H^*(\mathfrak{g})$. The previous representatives are all harmonic forms. The complex structure on the solvmanifold X can be defined in term of $(1, 0)$ -forms as follows:

$$\varphi^1 = e^1 + ie^4, \quad \varphi^2 = e^2 + ie^5, \quad \varphi^3 = e^3 + ie^6$$

We have that the real forms

$$\frac{1}{2}(\varphi^2 \wedge \varphi^3 + \bar{\varphi}^2 \wedge \bar{\varphi}^3), \frac{1}{2i}(\varphi^2 \wedge \varphi^3 - \bar{\varphi}^2 \wedge \bar{\varphi}^3), \frac{1}{2}(\varphi^1 \wedge \varphi^2 + \bar{\varphi}^1 \wedge \bar{\varphi}^2), \\ \frac{1}{2i}(\varphi^1 \wedge \varphi^2 - \bar{\varphi}^1 \wedge \bar{\varphi}^2), \frac{1}{2}(\varphi^1 \wedge \varphi^3 + \bar{\varphi}^1 \wedge \bar{\varphi}^3), \frac{1}{2i}(\varphi^1 \wedge \varphi^3 - \bar{\varphi}^1 \wedge \bar{\varphi}^3),$$

are anti-invariant and closed. Therefore, $\dim \mathcal{Z}_{\bar{J}}(X) > b_2(X)$ and by Corollary 2.3, the complex structure J does not admit any compatible Kähler metric.

This can be also derived by complex Hodge theory, since φ^2 is a non closed holomorphic 1-form.

It has also to be remarked that as a consequence of a result by Hasegawa (see [9, main theorem]) X does not admit any Kähler structure.

3.3. Locally almost-Kähler non globally almost Kähler manifold. In this section we will provide a family of 6-dimensional almost complex (non complex) manifolds (N, J) which are locally almost Kähler but not globally. We first recall the construction of N (see [16] and [2]). Let $A \in \text{SL}(2, \mathbb{Z})$ with two distinct real eigenvalues e^λ and $e^{-\lambda}$, where $\lambda > 0$. Let $Q \in \text{GL}(2, \mathbb{R})$ such that

$$QAQ^{-1} = \Lambda = \begin{pmatrix} e^{-\lambda} & 0 \\ 0 & e^\lambda \end{pmatrix}.$$

On \mathbb{C}^2 , with coordinates (z, w) , let \sim be defined by

$$\begin{pmatrix} z' \\ w' \end{pmatrix} \sim \begin{pmatrix} z \\ w \end{pmatrix} \iff \begin{pmatrix} z' \\ w' \end{pmatrix} = \begin{pmatrix} z \\ w \end{pmatrix} + Q \begin{pmatrix} m_1 + 2\pi i n_1 \\ m_2 + 2\pi i n_2 \end{pmatrix},$$

where $m_1, m_2, n_1, n_2 \in \mathbb{Z}$. Then \mathbb{C}^2 / \sim is a complex torus $\mathbb{T}_{\mathbb{C}}^2$ and

$$\Lambda \left[\begin{pmatrix} z \\ w \end{pmatrix} \right] = \left[\Lambda \begin{pmatrix} z \\ w \end{pmatrix} \right]$$

is a well defined automorphism of $\mathbb{T}_{\mathbb{C}}^2$. Indeed, if $\begin{pmatrix} z' \\ w' \end{pmatrix} \sim \begin{pmatrix} z \\ w \end{pmatrix}$, then

$$\begin{aligned} \Lambda \begin{pmatrix} z' \\ w' \end{pmatrix} &= \Lambda \begin{pmatrix} z \\ w \end{pmatrix} + \Lambda Q \begin{pmatrix} m_1 + 2\pi i n_1 \\ m_2 + 2\pi n_2 \end{pmatrix} = \\ &= \Lambda \begin{pmatrix} z \\ w \end{pmatrix} + QA \begin{pmatrix} m_1 + 2\pi i n_1 \\ m_2 + 2\pi n_2 \end{pmatrix} = \Lambda \begin{pmatrix} z \\ w \end{pmatrix} + Q \begin{pmatrix} m_1 + 2\pi i n_1 \\ m_2 + 2\pi n_2 \end{pmatrix} \end{aligned}$$

so that $\Lambda \begin{pmatrix} z' \\ w' \end{pmatrix} \sim \Lambda \begin{pmatrix} z \\ w \end{pmatrix}$.

For example, take

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}.$$

Then $\lambda = \log \frac{3+\sqrt{5}}{2}$ and we can choose

$$(3) \quad P = \begin{pmatrix} \frac{1-\sqrt{5}}{2} & 1 \\ 1 & \frac{\sqrt{5}-1}{2} \end{pmatrix}.$$

Set

$$\lambda = \log \frac{3 + \sqrt{5}}{2}, \quad \mu = \frac{\sqrt{5} - 1}{2}.$$

Let x_1, x_3, x_4, x_5, x_6 denote coordinates on \mathbb{R}^6 and, according to the previous notation, set $z = x_3 + ix_5, w = x_4 + ix_6$. Consider the following transformation of \mathbb{R}^5 :

$$T_1(x_1, x_3, x_4, x_5, x_6) = \left(x_1 + \lambda, e^\lambda x_3, e^{-\lambda} x_4, e^\lambda x_5, e^{-\lambda} x_6 \right).$$

We set

$$N = \frac{\mathbb{R}_{x_2}}{2\pi\mathbb{Z}} \times \frac{\mathbb{R}_{x_1} \times \mathbb{R}_{x_3, x_4, x_5, x_6}^4 / \Gamma}{\langle T_1(x) \rangle}$$

where

$$\Gamma = \text{Span}_{\mathbb{Z}} \langle (1, \mu, 0, 0)^t, (-\mu, 1, 0, 0)^t, (0, 0, 2\pi, 2\pi\mu)^t, (0, 0, -2\pi\mu, 2\pi)^t \rangle$$

and $\langle T_1(x) \rangle$ denotes the subgroup of transformations generated by $T_1(x)$, so that $\mathbb{T}_{\mathbb{C}}^2 \simeq \mathbb{R}_{x_3, x_4, x_5, x_6}^4 / \Gamma$. Then N is a compact 6-dimensional manifold. The following six 1-forms on \mathbb{R}^6

$$\begin{cases} e^1 = dx_1, \\ e^2 = dx_2, \\ e^3 = \exp(-x_1) dx_3, \\ e^4 = \exp(x_1) dx_4, \\ e^5 = \exp(-x_1) dx_5, \\ e^6 = \exp(x_1) dx_6, \end{cases}$$

induce 1-forms on the manifold N . Therefore, we immediately get

$$(4) \quad \begin{cases} de^1 = 0, \\ de^2 = 0, \\ de^3 = -e^1 \wedge e^3, \\ de^4 = e^1 \wedge e^4, \\ de^5 = -e^1 \wedge e^5, \\ de^6 = e^1 \wedge e^6. \end{cases}$$

The dual global frame $\{e_1, \dots, e_6\}$ on N is given by

$$\begin{aligned} e_1 &= \frac{\partial}{\partial x_1}, & e_2 &= \frac{\partial}{\partial x_2}, & e_3 &= \exp(x_1) \frac{\partial}{\partial x_3} \\ e_4 &= \exp(-x_1) \frac{\partial}{\partial x_4}, & e_5 &= \exp(x_1) \frac{\partial}{\partial x_5}, & e_6 &= \exp(-x_1) \frac{\partial}{\partial x_6} \end{aligned}$$

Let $f = f(x_2)$ be a never vanishing \mathbb{Z} -periodic function; let us define the almost complex structure J on N as

$$Je_1 = e_2, \quad Je_2 = -e_1, \quad Je_3 = f(x_2)e_5, \quad Je_4 = e_6, \quad Je_5 = -\frac{1}{f(x_2)}e_3, \quad Je_6 = -e_4.$$

Then it can be checked that J is integrable if and only if f is constant. We show that J is locally almost Kähler. Indeed, let ω be the local non degenerate and closed 2-form defined as

$$\omega = dx_1 \wedge dx_2 + dx_3 \wedge dx_5 + dx_4 \wedge dx_6;$$

then, since

$$\begin{aligned} J \frac{\partial}{\partial x_1} &= \frac{\partial}{\partial x_2}, & J \frac{\partial}{\partial x_2} &= -\frac{\partial}{\partial x_1}, & J \frac{\partial}{\partial x_3} &= f(x_2) \frac{\partial}{\partial x_5}, \\ J \frac{\partial}{\partial x_4} &= \frac{\partial}{\partial x_6}, & J \frac{\partial}{\partial x_5} &= -\frac{1}{f(x_2)} \frac{\partial}{\partial x_3}, & J \frac{\partial}{\partial x_6} &= -\frac{\partial}{\partial x_4}, \end{aligned}$$

we immediately get that $J\omega = \omega$ and $\omega(J\cdot, \cdot) > 0$ for any non-zero tangent vector, i.e., J is locally almost Kähler. Now we prove that J cannot be globally Kähler, and more generally that there is no global taming symplectic form. In view of Proposition 1.3, it is sufficient to find a nonzero J -anti-invariant exact form. To this purpose, let

$$\alpha = \cos(x_2)e^2 \wedge e^4 + \sin(x_2)e^1 \wedge e^4 - \sin(x_2)e^2 \wedge e^6 + \cos(x_2)e^1 \wedge e^6;$$

then, according to (4) and to definition of J , we have that $\alpha = d(\sin(x_2)e^4 + \cos(x_2)e^6)$ and that $J\alpha = -\alpha$, i.e., α is a J -anti-invariant exact 2-form.

REFERENCES

- [1] N. Aronszajn, A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order, *J. Math. Pure Appl.* **36** (1957), 235–249.
- [2] P. de Bartolomeis, A. Tomassini, On solvable generalized Calabi-Yau manifolds, *Ann. Inst. Fourier* **56** (2006), 1281–1296.
- [3] P. Deligne, P. Griffiths, J. Morgan, D. Sullivan, Real homotopy theory of Kähler manifolds, *Invent. Math.* **29** (1975), no. 3, 245–274.
- [4] S. K. Donaldson, *Two-forms on four-manifolds and elliptic equations*, Inspired by S. S. Chern, Nankai Tracts Math., vol. 11, World Sci. Publ., Hackensack, NJ, 2006, pp. 153–172.

- [5] T. Drăghici, T.-J. Li, W. Zhang, Symplectic forms and cohomology decomposition of almost complex four-manifolds, *Int. Math. Res. Not. IMRN* **2010** (2010), no. 1, 1–17.
- [6] T. Draghici, W. Zhang, A note on exact forms on almost complex manifolds, *Math. Res. Lett.* **19** (2012), 691–697.
- [7] Y. Eliashberg, N. Mishashev, *Introduction to the h-principle*, Graduate Studies in Mathematics, 48. American Mathematical Society, Providence, RI, 2002. xviii+206 pp..
- [8] M. Fernández, A. Gray, *The Iwasawa manifold*, Differential geometry, Peñíscola 1985, Lecture Notes in Math., vol. **1209**, Springer, Berlin, 1986, 157–159.
- [9] K. Hasegawa, A note on compact solvmanifolds with Kähler structures, *Osaka J. Math.* **43** (2006), 131–135.
- [10] R. Hind, C. Medori, A. Tomassini, On non pure forms on almost complex manifolds, to appear in *Proc. Amer. Math. Soc.*
- [11] D. Huybrechts, *Complex Geometry An Introduction*, Springer, Berlin, 2005.
- [12] J.L. Kazdan, Unique continuation in geometry, *Comm. Pure Appl. Math.* **41** (1988), 667–681.
- [13] M. Lejmi, Strictly nearly Kähler 6-manifolds are not compatible with symplectic forms, *C. R. Math. Acad. Sci. Paris* **343** (2006), 759–762.
- [14] T.-J. Li, W. Zhang, Comparing tamed and compatible symplectic cones and cohomological properties of almost complex manifolds, *Comm. Anal. Geom.* **17** (2009), 651–684.
- [15] D. McDuff, Application of convex integration to symplectic and contact geometry, *Ann. Inst. Fourier, (Grenoble)*, **37** (1987), 107–133.
- [16] I. Nakamura, Complex parallelisable manifolds and their small deformations, *J. Differential Geometry* **10** (1975), 85–112.
- [17] T. Peternell, Algebraicity Criteria for Compact Complex Manifolds, *Math. Ann.* **275** (1986), 653–672.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556
E-mail address: hind.1@nd.edu

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI PARMA, PARCO AREA DELLE SCIENZE
 53/A, 43124, PARMA, ITALY
E-mail address: costantino.medori@unipr.it
E-mail address: adriano.tomassini@unipr.it