


**General 1<sup>st</sup> derivative approximation (obtained by Lagrange interpolation)**

The interpolation nodes are given as:

- $(x_0, f(x_0))$
- $(x_1, f(x_1))$
- $(x_2, f(x_2))$
- ...
- $(x_N, f(x_N))$



By Lagrange Interpolation Theorem (Thm 3.3):

$$f(x) = \sum_{k=0}^n f(x_k)L_{N,k}(x) + \frac{(x-x_0)\cdots(x-x_N)}{(N+1)!} f^{(N+1)}(\xi(x)) \quad (1)$$

Take 1<sup>st</sup> derivative for Eq. (1):

$$f'(x) = \sum_{k=0}^n f(x_k)L'_{N,k}(x) + \frac{(x-x_0)\cdots(x-x_N)}{(N+1)!} \left( \frac{d(f^{(N+1)}(\xi(x)))}{dx} \right) + \frac{1}{(N+1)!} \left( \frac{d((x-x_0)\cdots(x-x_N))}{dx} \right) f^{(N+1)}(\xi(x))$$

Set  $x = x_j$ , with  $x_j$  being x-coordinate of one of interpolation nodes.  $j = 0, \dots, N$ .

$$f'(x_j) = \sum_{k=0}^n f(x_k)L'_{N,k}(x_j) + \frac{f^{(N+1)}(\xi(x))}{(N+1)!} \prod_{\substack{k=0 \\ k \neq j}}^N (x_j - x_k) \text{ ----- (N+1)-point formula to approximate } f'(x_j).$$

**The error of (N+1)-point formula is**  $\frac{f^{(N+1)}(\xi(x))}{(N+1)!} \prod_{\substack{k=0 \\ k \neq j}}^N (x_j - x_k).$

**Example.** The three-point formula with error to approximate  $f'(x_j)$ .


Let interpolation nodes be  $(x_0, f(x_0))$ ,  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ .

$$f'(x_j) = f(x_0) \left[ \frac{2x_j - x_1 - x_2}{(x_0 - x_1)(x_0 - x_2)} \right] + f(x_1) \left[ \frac{2x_j - x_0 - x_2}{(x_1 - x_0)(x_1 - x_2)} \right] + f(x_2) \left[ \frac{2x_j - x_0 - x_1}{(x_2 - x_0)(x_2 - x_1)} \right] + \frac{f^{(3)}(\xi(x))}{6} \prod_{\substack{k=0; \\ k \neq j}}^2 (x_j - x_k)$$

**Mostly used three-point formula (see Figure 1)**

Let  $x_0, x_1$ , and  $x_2$  be **equally spaced** and the grid spacing be  $h$ .

Thus  $x_1 = x_0 + h$ ; and  $x_2 = x_0 + 2h$ .



1.  $f'(x_0) = \frac{1}{2h} [-3f(x_0) + 4f(x_1) - f(x_2)] + \frac{h^2}{3} f^{(3)}(\xi(x_0))$  (three-point endpoint formula)


2.  $f'(x_1) = \frac{1}{2h} [-f(x_0) + f(x_2)] + \frac{h^2}{6} f^{(3)}(\xi(x_1))$  (three-point midpoint formula)

3.  $f'(x_2) = \frac{1}{2h} [f(x_0) - 4f(x_1) + 3f(x_2)] + \frac{h^2}{3} f^{(3)}(\xi(x_2))$  (three-point endpoint formula)

**Figure 1. Schematic for three-point formula**


**Mostly used five-point formula**

1. Five-point midpoint formula




$$f'(x_0) = \frac{1}{12h} [f(x_0 - 2h) - 8f(x_0 - h) + 8f(x_0 + h) - f(x_0 + 2h)] + \frac{h^4}{30} f^{(5)}(\xi)$$

## 2. Five-point endpoint formula



$$f'(x_0) = \frac{1}{12h} [-25f(x_0) + 48f(x_0 + h) - 36f(x_0 + 2h) + 16f(x_0 + 3h) - 3f(x_0 + 4h)] + \frac{h^4}{5} f^{(5)}(\xi)$$

**2<sup>nd</sup> derivative approximation (obtained by Taylor polynomial)**



Approximate  $f(x_0 + h)$  by expansion about  $x_0$ :

$$f(x_0 + h) = f(x_0) + f'(x_0)h + \frac{1}{2}f''(x_0)h^2 + \frac{1}{6}f'''(x_0)h^3 + \frac{1}{24}f^{(4)}(\xi_1)h^4 \quad (3)$$

Approximate  $f(x_0 - h)$  by expansion about  $x_0$ :

$$f(x_0 - h) = f(x_0) - f'(x_0)h + \frac{1}{2}f''(x_0)h^2 - \frac{1}{6}f'''(x_0)h^3 + \frac{1}{24}f^{(4)}(\xi_2)h^4 \quad (4)$$

Add Eqns. (3) and (4):

$$f(x_0 - h) + f(x_0 + h) = 2f(x_0) + f''(x_0)h^2 + \left[ \frac{1}{24}f^{(4)}(\xi_1)h^4 + \frac{1}{24}f^{(4)}(\xi_2)h^4 \right]$$

Thus

**Second derivative midpoint formula**

$$f''(x_0) = \frac{1}{h^2} [f(x_0 - h) - 2f(x_0) + f(x_0 + h)] - \frac{h^2}{12} f^{(4)}(\xi)$$