

Part I: Multiple choice questions (7 points each)

1. Find the closest point to $\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ in the subspace of \mathbb{R}^3 spanned by $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix}$.

- (a) $\begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$ (b) $\begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix}$ (c) $\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ (d) $\begin{bmatrix} 8/5 \\ 1 \\ 6/5 \end{bmatrix}$ (e) $\begin{bmatrix} -3/5 \\ 1 \\ 6/5 \end{bmatrix}$

$$W = \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix} \right\} = \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 2 \end{bmatrix} \right\} \quad \vec{x} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

↑
orthogonal basis

$$\text{proj}_W \vec{x} = \frac{\vec{x} \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 + \frac{\vec{x} \cdot \vec{v}_2}{\vec{v}_2 \cdot \vec{v}_2} \vec{v}_2 = \frac{1}{1} \vec{v}_1 + \frac{3}{5} \vec{v}_2 = \begin{bmatrix} -3/5 \\ 1 \\ 6/5 \end{bmatrix}$$

2. Which of the following is a least square solution \hat{x} to the equation

$$A \vec{x} = \vec{b} \quad \begin{matrix} \nearrow & \left[\begin{array}{cc} 1 & -2 \\ 2 & 1 \\ 1 & -2 \\ 2 & 1 \end{array} \right] & \left[\begin{array}{c} 1 \\ 3 \\ 1 \\ 3 \end{array} \right] \leftarrow \vec{b} \\ & & ? \end{matrix}$$

- (a) $\begin{bmatrix} 11/9 \\ 1/9 \end{bmatrix}$ (b) $\begin{bmatrix} 3/2 \\ 1/2 \end{bmatrix}$ (c) $\begin{bmatrix} 7/5 \\ 1/5 \end{bmatrix}$ (d) $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$ (e) $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$

$$A^T A = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix} \quad A^T \vec{b} = \begin{bmatrix} 14 \\ 2 \end{bmatrix}$$

Normal equation: $A^T A \vec{x} = A^T \vec{b}$

$$\left[\begin{array}{cc|c} 10 & 0 & 14 \\ 0 & 10 & 2 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & 0 & 14/10 \\ 0 & 1 & 2/10 \end{array} \right]$$

$$\text{so } \hat{x} = \begin{bmatrix} 7/5 \\ 1/5 \end{bmatrix}$$

3. Which of the following functions is a solution to the initial value problem

$$\frac{dy}{dt} = (y-t)^2 + 1; \quad y(0) = -1?$$

(a) $y = \frac{1}{t+1} - 2$ (b) $y = t$ (c) $y = \frac{-1}{t+1} + t$
 (d) $y = t - 1$ (e) $y = \frac{-2}{t+1} + 1$

Trial and error: eliminate (b) since $y(0) = 0 \neq -1$
 try (a), (c), (d), (e).

(c): $\frac{dy}{dt} = \frac{1}{(t+1)^2} + 1$

$$\Rightarrow \frac{dy}{dt} = (y-t)^2 + 1.$$

$$(y-t)^2 = \left(\frac{-1}{t+1}\right)^2 = \frac{1}{(t+1)^2}$$

4. Let A be an $m \times n$ matrix. Which of the following may be false?

- (a) The equation $A^T Ax = A^T b$ is always consistent for any b in \mathbb{R}^m .
- (b) $A^T A$ is invertible.
- (c) A solution to $A^T Ax = A^T b$ is a least squares solution of $Ax = b$.
- (d) The columns of A^T lie in the column space of $A^T A$.
- (e) If $A^T Ax = A^T b$ then $Ax - b$ is orthogonal to $\text{Col}(A)$.

(b) may be false

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \text{not invertible.}$$

5. Which of the following is a general solution to the differential equation

$$1 + \left(\frac{x}{y} - \sin y\right) \frac{dy}{dx} = 0?$$

- (a) $xy + y \sin y - \sin y = c$ (b) $xy + y \cos y - \sin y = cy$
 (c) $xy + y \sin y - \cos y = c$ (d) $xy + y \cos y - \sin y = c$
 (e) $xy + y \cos y - \cos y = c$

NOT on the exam!

6. Consider the initial value problem

$$\sin(2x) + \cos(3y) \frac{dy}{dx} = 0 \quad y(\pi/2) = \pi/3$$

Which of the following implicitly defines the solution?

- (a) $\frac{-\cos(2x)}{2} + \frac{\sin(3y)}{3} = \frac{-1}{2}$ (b) $-\cos(2x) + \sin(3y) = \frac{1}{2}$
 (c) $\sin(2x) + \cos(3y) = 1$ (d) $-\cos(2x) + \sin(3y) = \frac{-1}{2}$
 (e) $\frac{-\cos(2x)}{2} + \frac{\sin(3y)}{3} = \frac{1}{2}$

$$\int \cos(3y) dy = \int \sin(2x) dx$$

$$\frac{\sin(3y)}{3} = \frac{\cos(2x)}{2} + C$$

$$\frac{\sin(3y)}{3} = \frac{\cos(2x)}{2} + \frac{1}{2}$$

$$x = \frac{\pi}{2} \\ \implies \\ y\left(\frac{\pi}{2}\right) = \frac{\pi}{3}$$

$$\frac{\sin \pi}{3} = \frac{\cos \pi}{2} + C$$

$$0 = \frac{-1}{2} + C$$

$$C = \frac{1}{2}$$

7. Let $y(t)$ be the unique solution of the initial value problem

$$(t^2 - t) \frac{dy}{dt} + \cos(\pi t)y = \frac{t^2 - t}{t - 2} \quad y(3/2) = 0$$


What is the largest interval where y is defined?

- (a) $t > 0$ (b) $0 < t < 2$ (c) $1 < t < 2$ (d) $t < 1/2$ (e) $t < 2$

$$\frac{dy}{dt} + \frac{\cos(\pi t)}{t^2 - t} y = \frac{1}{t - 2}$$

$t=0, 1$
bad

$t=2$ bad



8. A tank initially contains 100l of pure water. Then, at $t = 0$, a sugar solution with concentration of 4g/l starts being pumped into the tank at a rate of 5l/min. The tank is kept well mixed, and the solution is being pumped out at the rate of 4l/min. Which of the following is the initial value problem for $y(t) =$ quantity of sugar, in grams, in the tank at time t ?

- (a) $\frac{dy}{dt} = 5y - 4(100 + t) \quad y(0) = 0$
- (b) $\frac{dy}{dt} = 20 - 4y \quad y(0) = 0$
- (c) $\frac{dy}{dt} = 4 \quad y(0) = 100$
- (d) $\frac{dy}{dt} = 20 - \frac{4y}{100 + t} \quad y(0) = 0$
- (e) $\frac{dy}{dt} = 20 - \frac{y}{(100 + t)^2} \quad y(0) = 100$

tank at time t
has
 $100 + 5t - 4t = 100 + t$
liters of mixture

$$\left\{ \begin{aligned} \frac{dy}{dt} &= 4 \cdot 5 - \frac{y}{100+t} \cdot 4 \\ &\quad \uparrow \text{going in} \quad \downarrow \text{going out} \\ y(0) &= 0 \end{aligned} \right.$$

no sugar at time 0

Part II: Partial credit questions (11 points each). Show your work.

9. Using the Gram-Schmidt Process, find an orthonormal basis of the subspace of \mathbb{R}^4

spanned by the vectors $\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 1 \\ 2 \\ 1 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 3 \\ 1 \\ 3 \end{bmatrix}$.

$\xrightarrow{\parallel} \quad \xrightarrow{\parallel} \quad \xrightarrow{\parallel}$
 $\vec{x}_1 \quad \vec{x}_2 \quad \vec{x}_3$

$$\vec{v}_1 = \vec{x}_1$$

$$\vec{v}_2 = \vec{x}_2 - \frac{\vec{x}_2 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1$$

$$= \begin{bmatrix} 1 \\ 2 \\ 1 \\ 2 \end{bmatrix} - \frac{1}{1} \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ 1 \\ 2 \end{bmatrix}$$

$$\vec{v}_3 = \vec{x}_3 - \frac{\vec{x}_3 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 - \frac{\vec{x}_3 \cdot \vec{v}_2}{\vec{v}_2 \cdot \vec{v}_2} \vec{v}_2$$

$$= \begin{bmatrix} 1 \\ 3 \\ 1 \\ 3 \end{bmatrix} - \frac{1}{1} \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} - \frac{13}{9} \begin{bmatrix} 0 \\ 2 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1/9 \\ -4/9 \\ 1/9 \end{bmatrix}$$

Orthonormal basis: $\vec{u}_i = \frac{1}{\|\vec{v}_i\|} \cdot \vec{v}_i$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2/3 \\ 1/3 \\ 2/3 \end{bmatrix}, \begin{bmatrix} 0 \\ 1/3\sqrt{2} \\ -2/3 \\ 1/3\sqrt{2} \end{bmatrix} \right\}$$

Orthogonal

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \quad \|\vec{v}_1\| = 1$$

$$\vec{v}_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \\ 2 \end{bmatrix} \quad \|\vec{v}_2\| = \sqrt{4+1+4} = 3$$

$$\vec{v}_3 = \begin{bmatrix} 0 \\ 1/9 \\ -4/9 \\ 1/9 \end{bmatrix}$$


$$\|\vec{v}_3\| = \frac{1}{9} \cdot \sqrt{1+16+1} = \frac{\sqrt{18}}{9} = \frac{\sqrt{2}}{3}$$

10. By drawing a direction field, sketch two solutions to the ODE

$$\frac{dy}{dt} = y^2(y-2)$$

with initial conditions $y(0) = 1$ and $y(0) = 3$.

Indicate clearly the limiting behavior $\lim_{t \rightarrow \infty} y(t)$ and $\lim_{t \rightarrow -\infty} y(t)$.



11. Find the function $y(t)$, for $t > 0$, which solves the initial value problem

$$t \frac{dy}{dt} + 4y = \frac{e^{-t}}{t^2}, \quad y(1) = 0$$

$$\frac{dy}{dt} + \underbrace{\left(\frac{4}{t}\right)}_{p(t)} y = \underbrace{\left(\frac{e^{-t}}{t^2}\right)}_{g(t)}$$

$$\mu(t) = e^{\int p(t) dt} = e^{4 \ln t} = t^4 \quad \text{integrating factor}$$

$$y(t) = \frac{\int \mu(t) \cdot g(t) dt}{\mu(t)}$$

$$\begin{aligned} \int t^4 \cdot \frac{e^{-t}}{t^2} dt &= \int t e^{-t} dt = t \cdot (-e^{-t}) - \int (-e^{-t}) dt \\ &= -t e^{-t} - e^{-t} + C \end{aligned}$$

$$\text{So } y(t) = \frac{-e^{-t}(t+1) + C}{t^4}$$

$$y(1) = 0 \quad \Rightarrow \quad \boxed{C = 2e^{-1}}$$

$$\boxed{y(t) = \frac{2e^{-1} - e^{-t}(t+1)}{t^4}}$$

12. Consider the differential equation

$$2y \frac{dy}{dx} = -e^x$$

- (a) Find the general solution.
(b) Find the solution with $y(0) = 1$.
(c) What is the largest interval in which the solution in part (b) is defined?

(a) $\int 2y dy = \int -e^x dx$
 $y^2 = -e^x + C$
 $y = \pm \sqrt{-e^x + C}$

(b) $y(0) = 1 \Rightarrow$ ~~positive~~ sign

$$1 = \sqrt{-e^0 + C} \Rightarrow \underline{\underline{C=2}}$$

$$y = \sqrt{2 - e^x}$$

(c) Need $2 - e^x \geq 0$ so

$$e^x \leq 2$$

$$\Leftrightarrow \boxed{x \leq \ln 2}$$

$$\boxed{I = (-\infty, \ln 2]}$$