

INTERMEDIATE GEOMETRY AND TOPOLOGY EXERCISES 2,
9/3/2021

1. Consider the Hopf fibration $S^3 \rightarrow \mathbb{C}\mathbb{P}^1$ as a principal S^1 -bundle. Show that the associated complex line bundle (using the standard action of $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ on \mathbb{C} by multiplication) is isomorphic to the tautological line bundle τ over $\mathbb{C}\mathbb{P}^1$.
2. Consider an Ehresmann connection in a trivial vector bundle $\pi: U \times \mathbb{R}^k \rightarrow U$ given by a horizontal distribution assigning to a point $(x, v) \in U \times \mathbb{R}^k$ the subspace $H_{(x,v)} = \text{graph}(-A \circ v) \subset T_{(x,v)}(U \times \mathbb{R}^k)$. Here $A \in \Omega^1(U, \mathfrak{gl}(k))$ – a matrix-valued 1-form on the base.

(a) Show that the corresponding covariant derivative operator¹ $\nabla: \Omega^p(U, \underline{\mathbb{R}}^k) \rightarrow \Omega^{p+1}(U, \underline{\mathbb{R}}^k)$ acts as $\nabla = d + A$.²

(b) Show that the curvature 2-form on the total space $\mathcal{F} \in \Omega^2(U \times \mathbb{R}^k, V)$, defined by $\mathcal{F}(X, Y) = [X_H, Y_H]_V$ for X, Y vector fields on the total space, has the form $\mathcal{F} = \pi^*F$ where

$$F = dA + \frac{1}{2}[A, A] \in \Omega^2(U, \mathfrak{gl}(k))$$

– a matrix-valued 2-form on the base.

3. Consider the trivial principal G -bundle $\mathcal{P} = U \times G$ over U with G a matrix Lie group. Assume the bundle \mathcal{P} is equipped with a connection defined by a 1-form $\mathcal{A} \in \Omega^1(\mathcal{P}, \mathfrak{g})$ on the total space. Let $\sigma: x \mapsto (x, 1)$ be the unit section of \mathcal{P} and let $A = \sigma^*\mathcal{A} \in \Omega^1(U, \mathfrak{g})$ be the connection 1-form on the base. Show that \mathcal{A} can be expressed in terms of A as

$$\mathcal{A}|_{(x,g)} = g^{-1}dg + g^{-1}A|_xg$$

where $(x, g) \in U \times G$ is a point in the total space.

4. (a) Assume that a connection in a vector bundle $E \rightarrow M$ of rank k is described in a local trivialization $\{U_\alpha, \phi_\alpha\}$ by matrix-valued 1-forms $A_\alpha \in \Omega^1(U_\alpha, \mathfrak{gl}(k))$. (I.e. in a trivialization the covariant derivative is $\nabla = d + A_\alpha$.) Show that on an overlap $U_\alpha \cap U_\beta$, one has

$$(1) \quad A_\beta = t_{\beta\alpha}A_\alpha t_{\alpha\beta} + t_{\beta\alpha}dt_{\alpha\beta}$$

with $t_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow GL(k)$ the transition functions.

¹Recall that we define ∇ on sections by $(\nabla\sigma)(v) = \frac{d}{dt}\Big|_{t=0} (\text{Hol}_{\gamma_0^t})^{-1}\sigma(\gamma(t))$ where $v \in T_xU$ is a tangent vector on a base at a point $x \in U$, γ is any curve $\gamma: [0, 1] \rightarrow U$ satisfying $\gamma(0) = x$, $\dot{\gamma}(0) = v$. $\text{Hol}_{\gamma_0^t}: \underbrace{E_x}_{\mathbb{R}^k} \rightarrow \underbrace{E_{\gamma(t)}}_{\mathbb{R}^k}$ is the parallel transport along the stretch of the curve γ from time

0 to time t .

²First consider the case $p = 0$, i.e., show that ∇ maps a section σ (understood as a vector-valued function on U) to $d\sigma + A \circ \sigma$, then extend the result to $p > 0$ by Leibniz identity.

- (b) Assume we have a connection in a principal G -bundle $\mathcal{P} \rightarrow M$ determined by a 1-form $\mathcal{A} \in \Omega^1(\mathcal{P}, \mathfrak{g})$. Assume that \mathcal{P} is trivialized over a cover $\{U_\alpha\}$ of M with trivializing local sections $\{\sigma_\alpha: U_\alpha \rightarrow \mathcal{P}|_{U_\alpha}\}$. Define local 1-forms of the connection as pullbacks $A_\alpha = \sigma_\alpha^* \mathcal{A} \in \Omega^1(U_\alpha, \mathfrak{g})$. Show that on an overlap $U_\alpha \cap U_\beta$ one has again the relation (1) between local connection 1-forms, where now transition functions take values in G .³
5. (a) Prove that for any connection in the tautological line bundle τ over $\mathbb{C}\mathbb{P}^1$, the integral of the curvature 2-form $F \in \Omega^2(\mathbb{C}\mathbb{P}^1)$ is

$$\int_{\mathbb{C}\mathbb{P}^1} F = 2\pi i$$

(In particular, there is no flat connection in τ .)⁴

- (b) Prove that for any connection in $\tau^{\otimes n}$, $n \in \mathbb{Z}$,⁵ one has

$$\int_{\mathbb{C}\mathbb{P}^1} F = 2\pi i n$$

³Here we assume for simplicity that G is a matrix Lie group. More generally, instead of (1), we should write $A_\beta = \text{Ad}_{t_{\beta\alpha}} A_\alpha + t_{\alpha\beta}^* \mu$ where $\mu \in \Omega^1(G, \mathfrak{g})$ is the Maurer-Cartan left-invariant 1-form on the group (which for a matrix group has the form $\mu = g^{-1} dg$).

⁴Hint: cut $\mathbb{C}\mathbb{P}^1$ into two disks B_\pm contained in open sets D_\pm of the trivializing cover for τ from Exercise sheet 1. A connection is represented by local 1-forms A_\pm on D_\pm , related on the overlap. Use this to evaluate $\int_{\mathbb{C}\mathbb{P}^1} F$ as $\int_{B_+} F + \int_{B_-} F$ where the two integrals can be evaluated in terms of local connection 1-forms A_\pm .

⁵By convention for any line bundle L , the inverse L^{-1} is understood as the dual bundle L^* . Thus, e.g., $\tau^{\otimes(-5)} = (\tau^*)^{\otimes 5}$. (Generally, isomorphism classes of line bundles over a fixed base M form a group under tensor product, with unit being the trivial line bundle and the inverse given by dualization.)