


2.1 The Bisection Method



Facts to remember:

1. The sequence of intervals $\{(a_i, b_i)\}_{i=1}^{\infty}$ contains the desired root.
2. Intervals containing the root: $(a_1, b_1) \supset (a_2, b_2) \supset (a_3, b_3) \supset (a_4, b_4) \dots$
3. After n steps, the interval (a_n, b_n) has the length:
$$b_n - a_n = \left(\frac{1}{2}\right)^{n-1} (b - a)$$
4. Let $p_n = \frac{b_n + a_n}{2}$ be the mid-point of (a_n, b_n) . The limit of sequence $\{p_n\}_{n=1}^{\infty}$ is the root.

Convergence

- **Theorem 2.1**

Suppose function $f(x)$ is continuous on $[a, b]$, and $f(a) \cdot f(b) < 0$. The Bisection method generates a sequence $\{p_n\}_{n=1}^{\infty}$ approximating a zero p of $f(x)$ with

$$|p_n - p| \leq \left(\frac{1}{2}\right)^n (b - a), \quad \text{when } n \geq 1$$

- **Convergence rate**

The sequence $\{p_n\}_{n=1}^{\infty}$ converges to p with the rate of convergence $O\left(\left(\frac{1}{2}\right)^n\right)$:

$$p_n = p + O\left(\left(\frac{1}{2}\right)^n\right)$$

Example 2.1.1. Show that $f(x) = x^3 + 4x^2 - 10 = 0$ has a root in $[1, 2]$, and use the Bisection method to determine an approximation to the root that is accurate to at least within 10^{-4} .

Remark: $|p_n - p| \leq (1/2)^n (b - a)$

or $|p_n - p| \leq (1/2)^n (b_n - a_n)$

- **Example 2.1.2.** Determine the number of iteration to solve $f(x) = x^3 + 4x^2 - 10 = 0$ with accuracy 10^{-3} . Use $a_1 = 1, b_1 = 2$.

Solution: Since $|p_n - p| \leq (1/2)^n (b_1 - a_1) \leq 10^{-3}, \rightarrow 2^{-n}(2 - 1) \leq 10^{-3}$.

Solve for $n \rightarrow n \approx 9.96$.

So $n = 10$ is needed.

- **Exercise 2.1.13.** Find an approximation to $\sqrt[3]{25}$ Correct within 10^{-4} using bisection method.

Solution: Consider to solve $f(x) = x^3 - 25 = 0$ by the Bisection method.

By trial and error, we can choose $a_1 = 2, b_1 = 3$.

Because $f(a_1) \cdot f(b_1) < 0$.

The Algorithm

INPUT **a,b**; tolerance **TOL**; maximum number of iterations **N0**.

OUTPUT solution p or message of failure.

STEP1 Set $i = 1$;
FA = $f(\mathbf{a})$;

STEP2 While $i \leq \mathbf{N0}$ do STEPs 3-6.

STEP3 Set $\mathbf{p} = \mathbf{a} + (\mathbf{b}-\mathbf{a})/2$; // a good way of computing middle point
FP = $f(\mathbf{p})$.

STEP4 IF FP = 0 or $(\mathbf{b}-\mathbf{a}) < \mathbf{TOL}$ then
OUTPUT (\mathbf{p});
STOP.

STEP5 Set $i = i + 1$.

STEP6 If $\text{FP} \cdot \text{FA} > 0$ then
Set $\mathbf{a} = \mathbf{p}$;
FA = FP.
else
set $\mathbf{b} = \mathbf{p}$;

STEP7 OUTPUT("Method failed after N0 iterations");
STOP.

Matlab Code

```
function p=bisection(f,a,b,tol)
```

```
while 1
```

```
    p=(a+b)/2;
```

```
    if p-a<tol, break; end
```

```
    if f(a)*f(p)>0
```

```
        a=p;
```

```
    else
```

```
        b=p;
```

```
    end
```

```
end %while 1
```