Math 43900 Problem Solving Fall 2016 Lecture 11 Functions and functional equations

Andrei Jorza

These problems are taken from the textbook, from Engels' *Problem solving* strategies, from Ravi Vakil's Putnam seminar notes and from Po-Shen Loh's Putnam seminar notes.

1 Functions and functional equations

You've seen in physics and calculus differential equations where you were supposed to determine a particular function f(x) satisfying a particular equation involing differentials. These are special examples of "functional equations", i.e., problems where you were supposed to determine a particular function f(x) given only an equation satisfied by f(x). They are a popular topic in math contests and solving them requires ingenuity and playfulness.

Example 1 (Cauchy's functional equation). The most classical example of a simple (nondifferential) functional equation is to determine functions $f : \mathbb{R} \to \mathbb{R}$ such that

$$f(x+y) = f(x) + f(y)$$

for all $x, y \in \mathbb{R}$.

As it stands the example has countless solutions (qnd I mean it in a technical way, there are uncountably many solutions). However, assuming mild properties of f(x) one can show that f(x) = ax for a fixed $a \in \mathbb{R}$ are the only solutions. This is the case when f(x) is assumed to be continuous, or even integrable.

Remark 1. A large number of functional equations can be reduced to Cauchy's functional equation via algebraic manipulations.

I identified 3 main topics:

- 1. Functional equations with integers, where you use the fact that the integers are discrete.
- 2. Functional equations over \mathbb{R} where you use algebraic manipulations.
- 3. Functional equations over \mathbb{R} where you use analytic properties of f(x), such that continuity or differentiability or integrability.

2 Problems

2.1 Functional equations and the integers

Easier

- 1. Suppose $f : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$ satisfies f(f(n)) = n + 3 for all $n \geq 0$ integer.
 - (a) Show that f(n+3) = f(n) + 3.
 - (b) Deduce that f(3k) = 3k + f(0), f(3k+1) = 3k + f(1) and f(3k+2) = 3k + f(2) for all nonnegative integers k.
- 2. Suppose $f : \mathbb{Q}_{>0} \to \mathbb{Q}_{>0}$ satisfies $f(xf(y)) = \frac{f(x)}{y}$ for all $x, y \in \mathbb{Q}_{>0}$.
 - (a) Show that f(f(y)) = f(1)/y, that f(f(1)) = 1 and deduce that f(1) = 1.
 - (b) Deduce that f(f(y)) = 1/y and show that f(1/y) = 1/f(y). [Hint: Apply f to the first equation.]
- 3. Suppose $f : \mathbb{Z}_{\geq 1} \to \mathbb{Z}_{\geq 1}$ satisfies f(n+1) > f(f(n)) for all $n \geq 1$.
 - (a) Show that f(1) is the minimum value of f.
 - (b) Show that $f(1) < f(2) < f(3) < \dots$

Harder

- 4. (Continuation of Exercise 1)
 - (c) Show that $f(f(n)) \equiv n \pmod{3}$ and conclude that either $f(x) \equiv x \pmod{3}$ for at least one of $x \in \{0, 1, 2\}$.

- (d) Deduce that no such function f(n) exists. [Hint: Use the previous part.]
- 5. (Continuation of Exercise 2)
 - (c) Show that f(x/y) = f(x)/f(y). [Hint: You know that f(f(y)) = 1/y.]
 - (d) Deduce that f(xy) = f(x)f(y) for all x, y.
 - (e) Can you find ONE example of such f.
- 6. (Continuation of Exercise 3)
 - (c) Show that f(n) > n can never happen.
 - (d) Deduce that f(n) = n for all n.

2.2 Functional equations and algebraic manipulations

Easier

7. Suppose $f : \mathbb{R} \to \mathbb{R}$ satisfies f(0) = 1/2 and there is some real α for which

$$f(x+y) = f(x)f(\alpha - y) + f(y)f(\alpha - x)$$

for all $x, y \in \mathbb{R}$.

- (a) Show that $f(\alpha) = 1/2$.
- (b) Show that $f(\alpha x) = f(x)$ for all x.
- 8. Suppose $f : \mathbb{R} \to \mathbb{R}$ satisfies xf(y) + yf(x) = (x+y)f(x)f(y). Show that for every $x \in \mathbb{R}$ we have $f(x) \in \{0, 1\}$. Can you show that f is an even function? [Hint: Play around with special values of x and y.]
- 9. Suppose $f : \mathbb{R} \to \mathbb{R}$ satisfies f(x)f(y) = f(x y) for all x, y and also suppose that f is not the 0 function. Show that f(0) = 1 and that for every $x \in \mathbb{R}$, $f(x) \in \{-1, 1\}$. [Hint: Play around with special values of x and y.]

Harder

- 10. (Continuation of Exercise 7)
 - (c) Show that $f(x) = \pm 1/2$ for all x.
 - (d) Show that in fact f(x) = 1/2 for all x.
- 11. Determine all functions $f : [0, \infty) \to [0, \infty)$ satisfying the following properties: (a) f(2) = 0, (b) if $x \in [0, 2)$ then $f(x) \neq 0$ and (c) if $x, y \in [0, \infty)$ then f(x+y) = f(xf(y))f(y).
- 12. Find the polynomials P(X) such that P(X+1) = P(X) + 2X + 1.

2.3 Functional equations and calculus

Easier

- 13. For each of the following functional equations find f(x) continuous that satisfy the equation:
 - (a) f(x+y) = f(x)f(y) with $f : \mathbb{R} \to (0, \infty)$. [Hint: Use log.]
 - (b) f(x+y) = f(x) + f(y) + f(x)f(y). [Hint: Reduce case (a).]
 - (c) f(xy) = f(x) + f(y) for $f: (0, \infty) \to \mathbb{R}$.
 - (d) f(xy) = xf(y) + yf(x) for $f: (0, \infty) \to \mathbb{R}$. [Hint: Divide by xy.]

Harder

- 14. Determine the continuous functions $f : \mathbb{R} \to \mathbb{R}$ such that f(x+y) = f(x)f(y). [Hint: Can you reduce to Exercise 13 (a)?]
- 15. Find the continuous functions $f : \mathbb{R} \to \mathbb{R}$ satisfying the functional equation

$$f\left(\frac{x+y}{2}\right) = \frac{f(x)+f(y)}{2}$$

[Hint: Compute f(x/2) in terms of f(x) and find a different functional equation satisfied by f.]

16. Determine the continuous functions $f : \mathbb{R} \to \mathbb{R}_{\neq 0}$ such that for all x, y

$$f(x+y) = \frac{f(x)f(y)}{f(x) + f(y)}$$