Math 43900 Problem Solving Fall 2016 Lecture 12 Inequalities

Andrei Jorza

These problems are taken from the textbook, from Engels' *Problem solving strategies*, from Ravi Vakil's Putnam seminar notes and from Po-Shen Loh's Putnam seminar notes.

1 Basics

Inequalities are a frequent and difficult topic on math competitions, and they are at the core of a huge number of results in analysis. Problem solving inequalities tend to be on the tricky side with ingenious algebra necessary to reduce them to some known inequalities. Nevertheless a handful of basic examples can be helpful in proving a large number of inequalities.

The basic inequalities:

- 1. By far the most useful inequality is that $x^2 \ge 0$ for all x real.
- 2. **AM-GM:** If $x_1, \ldots, x_n \geq 0$ then

$$\frac{x_1 + x_2 + \dots + x_n}{n} \ge \sqrt[n]{x_1 x_2 \cdots x_n}$$

with equality when $x_1 = x_2 = \ldots = x_n$.

3. Cauchy-Schwarz: If $x_1, \ldots, x_n, y_1, \ldots, y_n$ are real numbers then

$$(x_1^2 + x_2^2 + \dots + x_n^2)(y_1^2 + y_2^2 + \dots + y_n^2) \ge (x_1y_1 + x_2y_2 + \dots + x_ny_n)^2$$

with equality when $x_1 = \lambda y_1, x_2 = \lambda y_2, \ldots, x_n = \lambda y_n$ for a scalar λ .

4. Chebyshev's inequality: If $x_1 \le x_2 \le ... \le x_n$ and $y_1 \le y_2 \le ... \le y_n$ then

$$x_1y_1 + x_2y_2 + \dots + x_ny_n \ge x_1y_{\sigma(1)} + x_2y_{\sigma(2)} + \dots + x_ny_{\sigma(n)} \ge x_1y_n + x_2y_{n-1} + \dots + x_ny_1$$

for any permutation σ .

Needless to say you may use any method from calculus to show inequalities, from minimization/maximization to Lagrange multipliers. Typically, however, reducing inequalities to the basic ones via algebraic manipulations is the most effective strategy. Brute force methods sometimes work, but they are very laborious.

Inequalities come is lots of guises but the following are major themes in problem solving:

- 1. Inequalities based on AM-GM
- 2. Inequalities based on Cauchy-Schwarz
- 3. Inequalities in geometry, where a useful fact is the triangle inequality.
- 4. Inequalities in calculus

2 Problems

2.1 AM-GM, Completing the square

Easier

1. Show that for all real numbers x

$$2^x + 3^x - 4^x + 6^x - 9^x \le 1$$

[Hint: Complete the square.]

2. Show that $x^4 + 4x + 3 \ge 0$ for all real x. Find all positive integers n such that the equation

$$nx^4 + 4x + 3 = 0$$

has a real root.

Harder

3. Suppose $x_1, \ldots, x_n \in (1/4, 1)$. Show that

$$\log_{x_1}(x_2 - 1/4) + \log_{x_2}(x_3 - 1/4) + \dots + \log_{x_n}(x_1 - 1/4) \ge 2n$$

[Hint: Show that $x^2 \ge x - 1/4$ and then use AM-GM.]

- 4. Suppose a_1, \ldots, a_n are real numbers such that $a_1 + \cdots + a_n \ge n^2$ and $a_1^2 + \cdots + a_n^2 \le n^3 + 1$. Show that $a_1, \ldots, a_n \in [n-1, n+1]$. [Hint: Enough to show that $a_k n \in [-1, 1]$, or equivalently that $(a_k n)^2 \le 1$.]
- 5. Consider the real numbers $x_0 > x_1 > x_2 > \cdots > x_n$. Show that

$$x_0 + \frac{1}{x_0 - x_1} + \frac{1}{x_1 - x_2} + \dots + \frac{1}{x_{n-1} - x_n} \ge x_n + 2n$$

[Hint: Write $a_k = x_k - x_{k-1}$ and rewrite the inequality in terms of the a_k .]

2.2 Cauchy-Schwarz, Chebyshev

Easier

- 6. Find the maximum of the function f(x, y, z) = 5x 6y + 7z on the ellipsoid $2x^2 + 3y^2 + 4z^2 \le 1$. [Hint: Use calc 3 if you're up for it, but it's much easier with Cauchy-Schwarz. For the latter, maximize $f(x, y, z)^2$.]
- 7. If $a_1 + a_2 + \cdots + a_n = n$ show that $a_1^4 + a_2^4 + \cdots + a_n^4 \ge n$. [Hint: Apply Cauchy-Schwarz twice.]
- 8. Show that the positive real numbers a_0, a_1, \ldots, a_n form a geometric progression if and only if

$$(a_0a_1 + a_1a_2 + \dots + a_{n-1}a_n)^2 = (a_0^2 + a_1^2 + \dots + a_{n-1}^2)(a_1^2 + a_2^2 + \dots + a_n^2)$$

Harder

9. Show that if $0 < a, b < \pi/2$ then

$$\frac{\sin^3 a}{\sin b} + \frac{\cos^3 a}{\cos b} \ge \sec(a - b)$$

10. Find the positive integers n, k_1, \ldots, k_n are positive integers such that $k_1 + \cdots + k_n = 5n - 4$ and

$$\frac{1}{k_1} + \dots + \frac{1}{k_n} = 1$$

[Hint: Apply Cauchy-Schwarz to find n. Then play around.]

2.3 Inequalities in calculus and geometry

Easier

11. Suppose $f,g:[0,1]\to\mathbb{R}$ are continuous functions. Show that

$$\int_{0}^{1} f^{2}(x)dx \int_{0}^{1} g^{2}(x)dx \ge \left(\int_{0}^{1} f(x)g(x)dx\right)^{2}$$

[Hint: Use Riemann sums and Cauchy-Schwarz.]

Harder

12. Show that in a triangle with sides a, b, c and area A one has

$$a^2 + b^2 + c^2 \ge 4\sqrt{3}A$$

[Hint: $A = \frac{1}{2}bc\sin A$ and $a^2 = b^2 + c^2 - 2bc\cos A$.]