

# Math 30820 Honors Algebra 4

## Homework 3

Andrei Jorza

Due Wednesday, 2/8/2017

### Do 6 of the following questions.

Throughout this problem set  $R$  is an **integral domain**, unless otherwise specified.

- Let  $R$  be a ring,  $I$  an ideal of  $R$  and  $M$  an  $R$ -module.
  - Show that  $IM = \{\sum_{\text{finite } a_i, m_i | a_i \in I, m_i \in M}\}$  is an  $R$ -submodule of  $M$ .
  - Show that  $M/IM$  is an  $R/I$ -module.
- Consider  $\mathbb{C}$  as a  $\mathbb{Z}[i]$ -module under usual multiplication of complex numbers. Determine the torsion submodule of the  $\mathbb{Z}[i]$ -module  $\mathbb{C}/\mathbb{Z}[i]$ .
- Consider the ring  $A = \mathbb{F}_2[X]/(X^2 - X)$  with 4 elements. Show that torsion elements of the free  $A$ -module  $A$  do not form a submodule. (Note that  $A$  is not a domain so this does not contradict the statement from class.)
- Let  $M$  be a finitely generated  $R$ -module and  $\{m_1, \dots, m_n\}$  a linearly independent subset of  $M$ .
  - Show that  $N = \langle m_1, \dots, m_n \rangle$  is free  $\cong R^n$ .
  - If  $\{m_1, \dots, m_n\}$  is a maximal linearly independent subset show that  $M/N$  is torsion, i.e., every element of  $M/N$  is annihilated by a nonzero element of  $R$ .
- Let  $M$  be an  $R$ -module and  $\text{Tor}(M)$  its torsion submodule. Show that  $M/\text{Tor}(M)$  is torsion-free, i.e.,  $\text{Tor}(M/\text{Tor}(M)) = 0$ .
- Consider the  $R$ -module  $M = R^n \oplus N$  where  $N$  is a torsion module, i.e.,  $N = \text{Tor}(N)$ . Let  $e_1, \dots, e_n$  be the standard basis of  $R^n$  and  $t_1, \dots, t_n \in N$  arbitrary elements. Show that  $v_1 = e_1 + t_1, \dots, v_n = e_n + t_n$  are linearly independent and the map  $f : M \rightarrow M$  defined as the identity on  $N$  and sending  $e_i \mapsto v_i$  is an isomorphism. (The point of this exercise is that while  $N = \text{Tor}(M)$  is well-defined solely in terms of  $M$ , the free part  $R^n$  is not as every basis can be changed by torsion elements to get another basis.)
- Let  $f : M \rightarrow N$  be a homomorphism of  $R$ -modules. If  $\text{Im } f$  and  $\ker f$  are finitely generated, show that  $M$  is finitely generated.
- Let  $\phi : R \rightarrow S$  be a ring homomorphism and  $M$  an  $S$ -module. For  $r \in R$  and  $m \in M$  define  $r \cdot m := \phi(r)m$ , the later being scalar multiplication in  $M$  by  $\phi(r) \in S$ .
  - Show that this operation yields an  $R$ -module structure on the abelian group  $M$ . Call  $\phi^*M$  this  $R$ -module.
  - If  $f : M \rightarrow N$  is a homomorphism of  $S$ -modules define  $\phi^*f : \phi^*M \rightarrow \phi^*N$  by  $\phi^*f(m) = f(m)$ . Show that  $\phi^*f \in \text{Hom}_R(\phi^*M, \phi^*N)$ .

9. (An extra problem whose solution I won't write up) Let  $X^2 - aX + b \in \mathbb{R}[X]$  have complex roots  $u \pm vi$  with  $v \neq 0$ . Find a basis of the  $\mathbb{R}$ -vector space  $\mathbb{R}[X]/((X^2 - aX + b)^n)$  with respect to which the linear map "multiplication by  $X$ " has matrix

$$\begin{pmatrix} C & I_2 & & \\ & C & I_2 & \\ & & \ddots & I_2 \\ & & & C \end{pmatrix}$$

where  $C = \begin{pmatrix} u & v \\ -v & u \end{pmatrix}$ . This procedure yields the Jordan canonical form for real matrices.