# Math 43900 Problem Solving Fall 2018

# Lecture 12 Functions and functional equations

## Andrei Jorza

These problems are taken from the textbook, from Engels' *Problem solving strategies*, from Ravi Vakil's Putnam seminar notes and from Po-Shen Loh's Putnam seminar notes.

# 1 Functions and functional equations

You've seen in physics and calculus differential equations where you were supposed to determine a particular function f(x) satisfying a particular equation involing differentials. These are special examples of "functional equations", i.e., problems where you were supposed to determine a particular function f(x) given only an equation satisfied by f(x). They are a popular topic in math contests and solving them requires ingenuity and playfulness.

**Example 1** (Cauchy's functional equation). The most classical example of a simple (nondifferential) functional equation is to determine functions  $f: \mathbb{R} \to \mathbb{R}$  such that for all  $x, y \in \mathbb{R}$ :

$$f(x+y) = f(x) + f(y)$$

As it stands the example has countless solutions (qnd I mean it in a technical way, there are uncountably many solutions). However, assuming mild properties of f(x) one can show that f(x) = ax for a fixed  $a \in \mathbb{R}$  are the only solutions. This is the case when f(x) is assumed to be continuous, or even integrable.

Remark 1. A large number of functional equations can be reduced to Cauchy's functional equation via alegbraic manipulations.

I identified 3 main topics:

- 1. Functional equations with integers, where you use the fact that the integers are discrete.
- 2. Functional equations over  $\mathbb{R}$  where you use algebraic manipulations.
- 3. Functional equations over  $\mathbb{R}$  where you use analytic properties of f(x), such that continuity or differentiability or integrability.

## 2 Problems

## 2.1 Functional equations and the integers

#### Easier

- 1. (Putnam 1992) Show that f(n) = 1 n is the only integer-valued function defined on the integers that satisfies the following conditions:
  - (a) f(f(n)) = n for all integers n
  - (b) f(f(n+2)+2) = n for all integers n
  - (c) f(0) = 1.

#### Harder

- 2. Suppose  $f: \mathbb{Z}_{\geq 1} \to \mathbb{Z}_{\geq 1}$  satisfies f(n+1) > f(f(n)) for all  $n \geq 1$ .
  - (a) Show that f(1) is the minimum value of f.
  - (b) Show that  $f(1) < f(2) < f(3) < \dots$
  - (c) Show that f(n) > n can never happen.
  - (d) Deduce that f(n) = n for all n.

# 2.2 Functional equations and algebraic manipulations

#### Easier

3. (Putnam 1971) Let f(x) be a function defined on real numbers except 0 and 1. Find f(x) knowing that it satisfies f(x) + f(1 - 1/x) = 1 + x.

### Harder

- 4. (Putnam 1988) Show that there exists a unique function  $f(x):(0,\infty)\to(0,\infty)$  such that f(f(x))=6x-f(x) for all x>0.
- 5. (Putnam 1996) Let  $c \ge 0$  be a constant. Give a complete description of the set of continuous functions  $f: \mathbb{R} \to \mathbb{R}$  such that  $f(x) = f(x^2 + c)$  for all  $x \in \mathbb{R}$ .

# 2.3 Functional equations and calculus

#### Easier

- 6. (Putnam 1971) Find all polynomials P(x) such that  $P(x^2 + 1) = P(x)^2 + 1$  and P(0) = 0.
- 7. (Putnam 1991) Suppose f and g are nonconstant differentiable real-valued functions on  $\mathbb{R}$ . Also suppose that for all x, y real

$$f(x+y) = f(x)f(y) - g(x)g(y)$$
$$g(x+y) = f(x)g(y) + g(x)f(y)$$

If f'(0) = 0 show that  $f(x)^2 + g(x)^2 = 1$  for all x.

#### Harder

- 8. (Putnam 2000) Let  $f: [-1,1] \to \mathbb{R}$  be a continuous function such that  $f(2x^2 1) = 2xf(x)$  for all x. Show that f(x) = 0 for all x.
- 9. (Putnam 1971) Let u, f, g be real functions such that u(x+1) u(x-1) = 2f(x) and u(x+4) u(x-4) = 2g(x). Determine u in terms of f and g.

## 2.4 Extra problems

# Easier

- 10. Suppose  $f: \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$  satisfies f(f(n)) = n+3 for all  $n \geq 0$  integer.
  - (a) Show that f(n+3) = f(n) + 3.
  - (b) Deduce that f(3k) = 3k + f(0), f(3k+1) = 3k + f(1) and f(3k+2) = 3k + f(2) for all nonnegative integers k.

- 11. Suppose  $f: \mathbb{Q}_{>0} \to \mathbb{Q}_{>0}$  satisfies  $f(xf(y)) = \frac{f(x)}{y}$  for all  $x, y \in \mathbb{Q}_{>0}$ .
  - (a) Show that f(f(y)) = f(1)/y, that f(f(1)) = 1 and deduce that f(1) = 1.
  - (b) Deduce that f(f(y)) = 1/y and show that f(1/y) = 1/f(y).
- 12. Suppose  $f: \mathbb{R} \to \mathbb{R}$  satisfies f(0) = 1/2 and there is some real  $\alpha$  for which

$$f(x+y) = f(x)f(\alpha - y) + f(y)f(\alpha - x)$$

for all  $x, y \in \mathbb{R}$ .

- (a) Show that  $f(\alpha) = 1/2$ .
- (b) Show that  $f(\alpha x) = f(x)$  for all x.
- 13. Suppose  $f: \mathbb{R} \to \mathbb{R}$  satisfies xf(y) + yf(x) = (x+y)f(x)f(y). Show that for every  $x \in \mathbb{R}$  we have  $f(x) \in \{0,1\}$ . Can you show that f is an even function?
- 14. Suppose  $f: \mathbb{R} \to \mathbb{R}$  satisfies f(x)f(y) = f(x-y) for all x, y and also suppose that f is not the 0 function. Show that f(0) = 1 and that for every  $x \in \mathbb{R}$ ,  $f(x) \in \{-1, 1\}$ .
- 15. For each of the following functional equations find f(x) continuous that satisfy the equation:
  - (a) f(x+y) = f(x)f(y) with  $f: \mathbb{R} \to (0, \infty)$ .
  - (b) f(x+y) = f(x) + f(y) + f(x)f(y).
  - (c) f(xy) = f(x) + f(y) for  $f: (0, \infty) \to \mathbb{R}$ .
  - (d) f(xy) = xf(y) + yf(x) for  $f:(0,\infty) \to \mathbb{R}$ .

#### Harder

- 16. (Continuation of Exercise 10)
  - (c) Show that  $f(f(n)) \equiv n \pmod{3}$  and conclude that either  $f(x) \equiv x \pmod{3}$  for at least one of  $x \in \{0, 1, 2\}$ .
  - (d) Deduce that no such function f(n) exists.
- 17. (Continuation of Exercise 11)
  - (c) Show that f(x/y) = f(x)/f(y).
  - (d) Deduce that f(xy) = f(x)f(y) for all x, y.
  - (e) Can you find ONE example of such f.
- 18. (Continuation of Exercise 12)
  - (c) Show that  $f(x) = \pm 1/2$  for all x.
  - (d) Show that in fact f(x) = 1/2 for all x.
- 19. Determine all functions  $f:[0,\infty)\to[0,\infty)$  satisfying the following properties: (a) f(2)=0, (b) if  $x\in[0,2)$  then  $f(x)\neq 0$  and (c) if  $x,y\in[0,\infty)$  then f(x+y)=f(xf(y))f(y).
- 20. Find the polynomials P(X) such that P(X+1) = P(X) + 2X + 1.
- 21. (Putnam 2016) Find all functions  $f:(1,\infty)\to(1,\infty)$  with the following property: if  $x,y\in(1,\infty)$  and  $x^2\leq y\leq x^3$  then  $(f(x))^2\leq f(y)\leq (f(x))^3$ .

- 22. Determine the continuous functions  $f: \mathbb{R} \to \mathbb{R}$  such that f(x+y) = f(x)f(y). [Hint: Can you reduce to Exercise 15 (a)?]
- 23. Find the continuous functions  $f: \mathbb{R} \to \mathbb{R}$  satisfying the functional equation

$$f\left(\frac{x+y}{2}\right) = \frac{f(x) + f(y)}{2}$$

24. Determine the continuous functions  $f: \mathbb{R} \to \mathbb{R}_{\neq 0}$  such that for all x, y

$$f(x+y) = \frac{f(x)f(y)}{f(x) + f(y)}$$