

Multiple Choice

1.(6 pts) Use cylindrical coordinates to evaluate $\iiint_E (x^2 + y^2) dV$, where

$$E = \{(x, y, z) \mid \sqrt{x^2 + y^2} \leq z \leq 2\}.$$

- (a) $\frac{3\pi}{4}$ (b) $\frac{\pi}{4}$ (c) $\frac{4\pi}{9}$ (d) $\frac{16\pi}{5}$ (e) $\frac{4\pi}{3}$

2.(6 pts) Evaluate $\int_C xy ds$, where C is given by $\vec{r}(t) = \langle 4 \cos t, 4 \sin t, 3t \rangle$ for $0 \leq t \leq \frac{\pi}{2}$.

- (a) 10 (b) 40 (c) 5 (d) 0 (e) -40

3.(6 pts) Find the total mass of the laminated (i.e., thin) region D having density $\rho(x, y) = \sqrt{x^2 + y^2}$, where

$$D = \{(x, y) \mid x^2 + y^2 \leq 4, y \geq 0\}.$$

- (a) $\frac{4\pi}{3}$ (b) $\frac{8\pi}{3}$ (c) $\frac{3\pi}{2}$ (d) $\frac{4}{3}$ (e) $\frac{2\pi}{3}$

4.(6 pts) Use spherical coordinates to evaluate $\iiint_E e^{(x^2+y^2+z^2)^{3/2}} dV$, where

$$E = \{(x, y, z) \mid y \geq 0, z \geq 0, x^2 + y^2 + z^2 \leq 1\}.$$

- (a) $4\pi e$ (b) 0 (c) $\frac{\pi}{3}e$ (d) $\frac{\pi}{3}(e - 1)$ (e) $\frac{4\pi}{3}(e - 1)$

5.(6 pts) Let $\vec{F} = \langle xz, xyz, -y^2 \rangle$. Compute $\text{curl } \vec{F}$.

- (a) $\langle -y(2+x), x, yz \rangle$ (b) 0 (c) $z + xy$
 (d) $\langle x, -y(2+x), yz \rangle$ (e) $\langle -y(2+x), -x, yz \rangle$

6.(6 pts) Use a double integral to find the area enclosed by one loop of the four-leaved rose $r = \sin(2\theta)$. The region inside the loop is described in polar coordinates by $0 \leq \theta \leq \frac{\pi}{2}$ and $0 \leq r \leq \sin(2\theta)$.

- (a) $\frac{\pi}{2}$ (b) 0 (c) $-\frac{\pi}{2}$ (d) $\frac{\pi}{8}$ (e) $\frac{\pi}{4}$

7.(6 pts) Find the work $\int_C \vec{F} \cdot d\vec{r}$ done by the force field $\vec{F} = \langle xy, yz, zx \rangle$ in moving a particle along the curve C given by $\vec{r}(t) = \langle t, t^2, t^3 \rangle$ for $-1 \leq t \leq 1$.

- (a) $\frac{1}{4}$ (b) $\frac{5}{7}$ (c) $\frac{1}{2}$ (d) $\frac{10}{7}$ (e) $\frac{27}{28}$

8.(6 pts) Use Green's theorem to evaluate $\int_C \left((3y - e^{x^2})dx + (7x + \sqrt{y^{99} + y + 100})dy \right)$ where C is the circle $x^2 + y^2 = 9$ with the counter-clockwise orientation.

- (a) 3π (b) 36π (c) 0 (d) -36π (e) 7π

9.(6 pts) Let $x = 2u$ and $y = -3v$. Then $\int_{-3}^3 \int_{-2}^2 f(x, y) dx dy$ can be written as:

- (a) $\frac{1}{6} \int_{-1}^1 \int_{-1}^1 f(2u, -3v) dudv$ (b) $6 \int_{-3}^3 \int_{-2}^2 f(2u, -3v) dudv$
 (c) $6 \int_{-1}^1 \int_{-1}^1 f(2u, -3v) dudv$ (d) $-4 \int_{-1}^1 \int_{-1}^1 f(2u, -3v) dudv$
 (e) $-6 \int_{-1}^1 \int_{-1}^1 f(2u, -3v) dudv$

10.(6 pts) Which of the following vector fields cannot be written as $\text{curl } \vec{F}$?

- (a) $\langle -x - y + 1, xy - 1, -xz + y + z \rangle$ (b) $\langle -y, -z, -x \rangle$
(c) $\langle -y \cos(z), -z \cos(x), -x \cos(y) \rangle$ (d) $\langle 2yz, xyz, 3xy \rangle$
(e) $\langle 1 - 2z, 1 - 2x, 1 - 2y \rangle$

Partial Credit

You must show your work on the partial credit problems to receive credit!

11.(12 pts.) Let $\vec{F} = \langle y^2 + 1, 2xy + 2y + e^{3z}, 3ye^{3z} + 3z^2 \rangle$.

- (a) Find f such that $\nabla f = \vec{F}$.
(b) Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where C is any smooth curve beginning at $(1, 0, 0)$ and ending at $(0, 1, 0)$.

12.(12 pts.) Let E be the tetrahedron enclosed by the coordinate planes $x = 0$, $y = 0$, $z = 0$ and the plane $2x + y + z = 2$. Assume the density function is $\rho(x, y, z) = 1$. Write an iterated integral (with limits) for the moment of the solid E about the yz -plane. (You do NOT need to compute this iterated integral.)

13.(12 pts.) Use the transformation $x = \sqrt{3}u - v$, $y = \sqrt{3}u + v$ to evaluate the integral $\iint_R (x^2 - xy + y^2) dA$, where R is the region bounded by the ellipse $x^2 - xy + y^2 = 3$.

Name: _____

Instructor: ANSWERS

Math 20550, Last Year Exam 3
November 16, 2017

- The Honor Code is in effect for this examination. All work is to be your own.
- No calculators.
- The exam lasts for 1 hour and 15 minutes
- Be sure that your name is on every page in case pages become detached.
- Be sure that you have all 4 pages of the test.
- Each multiple choice question is 6 points. Each partial credit problem is 12 points.
You will receive 4 extra points.

PLEASE MARK YOUR ANSWERS WITH AN X, not a circle!

- | | | | | | |
|-----|-----|-----|-----|-----|-----|
| 1. | (a) | (b) | (c) | (●) | (e) |
| 2. | (a) | (●) | (c) | (d) | (e) |
| 3. | (a) | (●) | (c) | (d) | (e) |
| 4. | (a) | (b) | (c) | (●) | (e) |
| 5. | (●) | (b) | (c) | (d) | (e) |
| 6. | (a) | (b) | (c) | (●) | (e) |
| 7. | (a) | (b) | (c) | (●) | (e) |
| 8. | (a) | (●) | (c) | (d) | (e) |
| 9. | (a) | (b) | (●) | (d) | (e) |
| 10. | (a) | (b) | (c) | (●) | (e) |

Please do NOT write in this box.

Multiple Choice _____

11. _____

12. _____

13. _____

Extra Points. 4 _____

Total: _____

11.

(a) We seek a FUNCTION f such that $\nabla f = \vec{F}$, i.e., such that

$$(a) \quad f_x = y^2 + 1$$

$$(b) \quad f_y = 2xy + 2y + e^{3z}$$

$$(c) \quad f_z = 3ye^{3z} + 3z^2.$$

Integrating (a) with respect to x we get

$$(1) \quad f(x, y, z) = xy^2 + x + h(y, z)$$

where the functions h may depend on y and z only and does not depend on x .

We differentiate the function f from (1) with respect to y and using (b) set equal to $2xy + 2y + e^{3z}$:

$$2xy + h_y(y, z) = 2xy + 2y + e^{3z},$$

hence

$$h_y(y, z) = 2y + e^{3z}.$$

Integrating the above expression with respect to y we get

$$h(y, z) = y^2 + ye^{3z} + g(z),$$

where the functions g may depend on z only and does not depend on x, y . From (1) we obtain

$$(2) \quad f(x, y, z) = xy^2 + x + y^2 + ye^{3z} + g(z).$$

We differentiate the function f from (2) with respect to z and using (c) set equal to $3ye^{3z} + 3z^2$:

$$3ye^{3z} + g_z(z) = 3ye^{3z} + 3z^2,$$

hence

$$g_z(z) = 3z^2,$$

and

$$g(z) = z^3 + C.$$

We choose $C = 0$ and use (2) to obtain

$$f(x, y, z) = xy^2 + x + y^2 + ye^{3z} + z^3$$

(b) Finally, the fundamental theorem of line integrals tells us that

$$\int_C \vec{F} \cdot d\vec{r} = \int_C \nabla f \cdot d\vec{r} = f(0, 1, 0) - f(1, 0, 0) = 2 - 1 = 1$$

12. In the xy -plane the condition $2x + y + z = 2$ becomes $2x + y = 2$ so the x -intercept is $x = 1$.

Thus the base of E in the xy -plane is the region D bounded by the lines $x = 0$, $y = 0$, $y = 2 - 2x$, that we can write as $D = \{(x, y) | 0 \leq x \leq 1, 0 \leq y \leq 2 - 2x\}$.

The region E can be describes as the region that lies above D and below the plane $z = 2 - 2x - y$.

The moment is therefore

$$\iiint_E x \, dV = \int_0^1 \int_0^{2-2x} \int_0^{2-2x-y} x \, dz \, dy \, dx$$

13. We denote by $f(x, y)$ the function $f(x, y) = x^2 - xy + y^2$.

Thus we need to evaluate

$$\iint_R f(x, y) \, dA.$$

The ellipse $x^2 - xy + y^2 = 3$, which bounds the region R , when written in terms of u and v becomes

$$x^2 - xy + y^2 = (\sqrt{3}u - v)^2 - (\sqrt{3}u - v)(\sqrt{3}u + v) + (\sqrt{3}u + v)^2 = 3(u^2 + v^2) = 3$$

which is the circle S of radius 1. Hence under the change of variables the region R in the (x, y) -plane is transformed to the unit disk D in the (u, v) -plane.

The function $f(x, y)$ in terms of u and v is

$$f = x^2 - xy + y^2 = (\sqrt{3}u - v)^2 - (\sqrt{3}u - v)(\sqrt{3}u + v) + (\sqrt{3}u + v)^2 = 3(u^2 + v^2).$$

The Jacobian is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \sqrt{3} & -1 \\ \sqrt{3} & 1 \end{vmatrix} = 2\sqrt{3}$$

So

$$\begin{aligned} \iint_R (x^2 - xy + y^2) \, dx \, dy &= \iint_D 3(u^2 + v^2) |2\sqrt{3}| \, du \, dv \\ &= 6\sqrt{3} \int_0^{2\pi} \int_0^1 r^2 \, r \, dr \, d\theta = 6\sqrt{3} \int_0^{2\pi} \frac{1}{4} \, d\theta = 3\sqrt{3}\pi \end{aligned}$$