

# Math 43900 Problem Solving

## Fall 2021

### Lecture 10 Matrices

Andrei Jorza

Evan O'Dorney

These problems are taken from the textbook, from Engel's *Problem solving strategies*, from Ravi Vakil's Putnam seminar notes and from Po-Shen Loh's Putnam seminar notes.

## 1 Matrices

### Overview

The way matrices show up in problem solving problems involves the following three main themes:

1. algebraic manipulations of matrices (they can be multiplied and the operation is not commutative),
2. determinants and eigenvalues of matrices,
3. matrices as defining linear maps on vector spaces.

### Basic results

1. You can always add two  $m \times n$  matrices.
2. You can always multiply an  $m \times n$  matrix and an  $n \times p$  matrix to get an  $m \times p$  matrix.
3. The **trace** of a matrix  $\text{Tr} A$  is the sum of its diagonal terms. It has the property that  $\text{Tr}(A + B) = \text{Tr}(A) + \text{Tr}(B)$  and  $\text{Tr}(AB) = \text{Tr}(BA)$  for all matrices  $A$  and  $B$ .
4. The **determinant** of a matrix  $\det A$  is a polynomial expression in the entries of the matrix  $A$  and satisfies the following properties:

- (a) The determinant of  $(a_{ij})$  is  $\sum_{\sigma \in S_n} \varepsilon(\sigma) \prod_{i=1}^n a_{i, \sigma(i)}$ , where  $S_n$  is the group of permutations and  $\varepsilon(\sigma)$  is the sign. The sign  $\varepsilon$  is multiplicative and if  $\tau$  is a  $k$ -cycle then  $\varepsilon(\tau) = (-1)^{k-1}$ .
- (b) If in a matrix  $A = (a_{ij})$  you write  $A_{p,q}$  for the  $(n-1) \times (n-1)$  where you eliminate the  $p$ -th row and  $q$ -th column from  $A$  then

$$\det(A) = a_{11} \det A_{11} - a_{12} \det A_{12} + \cdots + (-1)^{n-1} a_{1,n} \det A_{1,n}$$

- (c)  $A$  is invertible if and only if  $\det A \neq 0$ .
- (d)  $\det(AB) = \det(A) \det(B)$  for all matrices  $A$  and  $B$ .
- (e) If you swap two rows or columns of a matrix  $A$  to obtain a matrix  $B$  then  $\det(B) = -\det(A)$ .
- (f) If in a matrix  $A$  you add a multiple of one row to a different row to get a matrix  $B$  then  $\det(B) = \det(A)$ . The same is true if you add a multiple of a column to a different column.

5. Suppose  $A$  is an  $n \times n$  matrix. If you can find a *nonzero* vector  $v$  (i.e., an  $n \times 1$  matrix consisting of a single column) and a scalar  $\alpha$  such that  $Av = \alpha v$  then  $\alpha$  is said to be an **eigenvalue** of  $A$  with **eigenvector**  $v$ .

6. If  $A$  is an  $n \times n$  matrix the **characteristic polynomial** of  $A$  is the monic degree  $n$  polynomial

$$P_A(X) = \det(XI_n - A)$$

(a) A scalar  $\alpha$  is an eigenvalue of  $A$  if and only if it is a root of  $P_A(X)$ . The roots of  $P_A(X)$  are **the** eigenvalues of  $A$  and are counted with multiplicity if they are not distinct. E.g.,  $I_n$  has  $n$  eigenvalues all equal to 1.

(b)  $P_A(X) = X^n - (\text{Tr } A)X^{n-1} + \dots + (-1)^n \det(A)$ .

(c) Since we know the relation between the coefficients of a polynomial and its roots we deduce that if  $\lambda_1, \dots, \lambda_n$  are the eigenvalues of  $A$  then

$$\text{Tr}(A) = \lambda_1 + \lambda_2 + \dots + \lambda_n$$

$$\det(A) = \lambda_1 \lambda_2 \dots \lambda_n$$

(d) The Cayley-Hamilton theorem: If you plug  $A$  into the polynomial  $P_A(X)$  you always get the 0 matrix,  $P_A(A) = O$ .

(e) If  $A$  and  $B$  are matrices then  $P_{AB}(X) = P_{BA}(X)$  as polynomials.

7. A big result in linear algebra says that for any matrix  $A$  (over  $\mathbb{C}$ ) you can find an invertible matrix  $S$  such that the conjugate  $SAS^{-1}$  has a very special shape: the **Jordan canonical form**. In fact the Jordan canonical form  $SAS^{-1}$  has the  $n$  eigenvalues on the diagonal but much more is true:  $SAS^{-1}$  is block diagonal and each block is of the form

$$\begin{pmatrix} \lambda & 1 & 0 & \dots \\ 0 & \lambda & 1 & \dots \\ & & \ddots & \ddots \\ 0 & \dots & 0 & \lambda \end{pmatrix}$$

with an eigenvalue  $\lambda$  on the diagonal and 1-s off diagonal. E.g., for a  $2 \times 2$  matrix the possible Jordan canonical forms are

$$\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \text{ for } \lambda_1 \neq \lambda_2 \text{ and } \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \text{ or } \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$$

8. (**VERY USEFUL**) Suppose  $A$  is an  $n \times n$  matrix and  $Q(X)$  is any polynomial. If the eigenvalues of  $A$  are  $\lambda_1, \dots, \lambda_n$  then the eigenvalues of  $Q(A)$  (also an  $n \times n$  matrix) are  $Q(\lambda_1), \dots, Q(\lambda_n)$ .

## 2 Problems

### 2.1 Determinants, traces, characteristic polynomials and eigenvalues

**Easier**

1. (Putnam 1978) Let  $a \neq b$  and  $p_1, \dots, p_n$  be real numbers, and let  $F(X) = (p_1 - X) \dots (p_n - X)$ . Let  $M$  be the  $n \times n$  matrix which has  $p_1, \dots, p_n$  on the diagonal,  $a$  above the diagonal, and  $b$  below the diagonal. Show that

$$\det M = \frac{bF(a) - aF(b)}{b - a}.$$

2. (Putnam 1969) Show that  $\det(|i - j|)_{1 \leq i, j \leq n} = (-1)^{n-1} (n-1) 2^{n-2}$ .

3. Let  $D_n$  be the  $(n-1) \times (n-1)$  determinant that has  $3, 4, \dots, n+1$  on the diagonal and 1's everywhere else. Show that  $\{D_n/n!\}$  is unbounded.

### Harder

- (Putnam 1984) Let  $M(x) = (m_{i,j})$  be the  $2n \times 2n$  matrix with entries  $m_{i,j} = x$  if  $i = j$ ,  $m_{i,j} = a$  if  $i \neq j$  and  $i + j$  is even, and  $m_{i,j} = b$  if  $i \neq j$  and  $i + j$  is odd. Compute  $\lim_{x \rightarrow a} \frac{\det M(x)}{(x - a)^{2n-2}}$ .
- (Putnam 1985) Let  $G = \{M_1, \dots, M_r\}$  be a finite set of  $n \times n$  matrices which form a group under matrix multiplication. Suppose  $\sum_{i=1}^r \text{Tr}(M_i) = 0$ . Show that  $\sum_{i=1}^r M_i = 0_{n \times n}$ .

## 2.2 Algebraic operations and linear algebra

### Easier

- Compute  $\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}^n$  and  $\begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}^n$  for all  $n$ .
- Suppose  $f(x) = a_0 + a_1x + a_2x^2 + \dots$  is a converging power series. Show that  $f(SAS^{-1}) = Sf(A)S^{-1}$ .

### Harder

- (Putnam 1986) Let  $A, B, C, D$  be  $n \times n$  matrices with complex entries such that:  $AB^t$  and  $CD^t$  are symmetric and  $AD^t - BC^t = I_n$ . Show that  $A^tD - C^tB = I_n$ .
- (Putnam 1987) Let  $M$  be a  $2n \times n$  matrix with complex entries such that whenever  $(z_1, \dots, z_{2n})M = 0_{1 \times n}$  with complex  $z_i$ , not all 0, then at least one  $z_i$  is not real. Show that for any real  $r_1, \dots, r_{2n}$  there exist complex  $z_1, \dots, z_n$  such that  $\text{Re}(M(z_1, \dots, z_n)^t) = (r_1, \dots, r_{2n})^t$ .

## 2.3 Extra problems

### Easier

- Show that you can never find two  $n \times n$  matrices  $A$  and  $B$  with real coefficients such that  $AB - BA = I_n$ .
- Consider an  $n \times (n + 1)$  matrix  $A = (a_{ij})$ . For a column  $k$  write  $A_k$  for the  $n \times n$  matrix you obtain from  $A$  by removing the  $k$ -th column. Show that

$$a_{11} \det A_1 - a_{12} \det A_2 + \dots + (-1)^{n+1} a_{1,n+1} \det A_{n+1} = 0$$

- Suppose  $P(X)$  is a polynomial and  $A$  is an  $n \times n$  matrix such that  $P(A) = 0$ . Show that the eigenvalues of  $A$  are among the roots of  $P(X)$ .
- This is an application of Exercise 19. Suppose  $X$  is an antisymmetric matrix, i.e., of the form  $X = -X^t$ . (Think  $\begin{pmatrix} & x \\ -x & \end{pmatrix}$ .) Show that every eigenvalue of  $X$  is of the form  $ai$  where  $i = \sqrt{-1}$  and  $a \in \mathbb{R}$ .
- Show that  $A^k = 0$  for some  $k \geq 0$  if and only if all the eigenvalues of  $A$  are 0 in which case  $A^n = 0$  as well.
- (Putnam 1994) Let  $A$  and  $B$  be 2 by 2 matrices with integer entries such that  $A, A+B, A+2B, A+3B$  and  $A+4B$  are all invertible matrices whose inverses have integer entries. Show that  $A+5B$  is invertible and that its inverse has integer entries.

16. Let  $p < m$  be positive integers. Show that

$$\det \begin{pmatrix} \binom{m}{0} & \binom{m}{1} & \cdots & \binom{m}{p} \\ \binom{m+1}{0} & \binom{m+1}{1} & \cdots & \binom{m+1}{p} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{m+p}{0} & \binom{m+p}{1} & \cdots & \binom{m+p}{p} \end{pmatrix} = 1.$$

17. Suppose  $(x_n)$  is a sequence defined by the linear recurrence  $x_{n+2} = ax_{n+1} + bx_n$  for all  $n \geq 0$ . Show that

$$\begin{pmatrix} x_{n+2} \\ x_{n+1} \end{pmatrix} = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_{n+1} \\ x_n \end{pmatrix}$$

and conclude that for  $n \geq 1$ ,  $x_n$  is the first entry of the matrix  $\begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix}^{n-1} \begin{pmatrix} x_1 \\ x_0 \end{pmatrix}$ .

18. A useful application of Exercise 6. Show that if  $f(x) = a_0 + a_1x + a_2x^2 + \cdots$  is an absolutely convergent power series then  $f\left(\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}\right) = \begin{pmatrix} f(\lambda_1) & 0 \\ 0 & f(\lambda_2) \end{pmatrix}$  and  $f\left(\begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}\right) = \begin{pmatrix} f(\lambda) & f'(\lambda) \\ 0 & f(\lambda) \end{pmatrix}$ .

19. If  $u$  and  $v$  are  $n \times 1$  column matrices write  $\langle u, v \rangle = u^t v$  for the dot product of the two vectors. If  $A$  is an  $n \times n$  matrix show that  $\langle u, Av \rangle = \langle A^t u, v \rangle$ . Show that  $\langle v, \bar{v} \rangle \geq 0$ , where  $\bar{v}$  is the complex conjugate of  $v$ .

20. If  $A = (a_{ij})$  show that  $\text{Tr}(A \cdot A^t) = \sum_{i,j} a_{ij}^2$ .

### Harder

21. Suppose  $A$  is an  $n \times n$  real matrix such that  $A^2 = A + I_n$ . Show that  $\det(A) < 2^n$ . In fact show that  $\det(A) \leq \left(\frac{1 + \sqrt{5}}{2}\right)^n$ .

22. Suppose  $X$  is a real matrix with  $X + X^t = I_n$ . Show that  $\det X \geq \frac{1}{2^n}$ .

23. Compute the determinant of the matrix  $(a_{ij})$  where  $a_{ii} = 2$  and if  $i \neq j$  then  $a_{ij} = (-1)^{i-j}$ .

24. Let  $A$  and  $B$  be  $3 \times 3$  matrices with real entries such that  $\det A = \det B = \det(A+B) = \det(A-B) = 0$ . Show that  $\det(xA + yB) = 0$  for all real numbers  $x, y$ .

25. Let  $n$  be an odd positive integer. Suppose  $A$  is an  $n \times n$  matrix whose square  $A^2$  is either  $0$  or  $I_n$ . Show that  $\det(A + I_n) \geq \det(A - I_n)$ .

26. Suppose  $A$  and  $B$  are commuting  $n \times n$  matrices with real entries such that  $\det(A + B) \geq 0$ . Show that  $\det(A^k + B^k) \geq 0$  for all  $k \geq 1$ .

27. (Putnam 1996) Show that there exists no complex matrix  $A$  such that  $\sin(A) = \begin{pmatrix} 1 & 1996 \\ 0 & 1 \end{pmatrix}$ .

28. Suppose  $A$  and  $B$  are  $n \times n$  real matrices such that  $\text{Tr}(A \cdot A^t + B \cdot B^t) = \text{Tr}(A \cdot B + A^t \cdot B^t)$ . Show that  $A = B^t$ .