

A Birman exact sequence for the Torelli subgroup of $\text{Aut}(F_n)$

Matthew Day* and Andrew Putman†

March 30, 2016

Abstract

We develop an analogue of the Birman exact sequence for the Torelli subgroup of $\text{Aut}(F_n)$. This builds on earlier work of the authors who studied an analogue of the Birman exact sequence for the entire group $\text{Aut}(F_n)$. These results play an important role in the authors' recent work on the second homology group of the Torelli group.

1 Introduction

The Birman exact sequence [2, 9] is a fundamental result that relates the mapping class groups of surfaces with differing numbers of boundary components. It is frequently used to understand the stabilizers in the mapping class group of simple closed curves on a surface. In [5], the authors constructed an analogous exact sequence for the automorphism group $\text{Aut}(F_n)$ of the free group F_n on n letters $\{x_1, \dots, x_n\}$. The *Torelli subgroup* of $\text{Aut}(F_n)$, denoted IA_n , is the kernel of the map $\text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z})$ obtained from the action of $\text{Aut}(F_n)$ on $F_n^{\text{ab}} \cong \mathbb{Z}^n$. In this paper, we construct a version of the Birman exact sequence for IA_n . This new exact sequence plays a key role in our recent paper [6] on $H_2(\text{IA}_n; \mathbb{Z})$.

Birman exact sequence. Let $F_{n,k}$ be the free group on the set $\{x_1, \dots, x_n, y_1, \dots, y_k\}$. For $z \in F_{n,k}$, let $\llbracket z \rrbracket$ denote the conjugacy class of z . Define

$$\mathcal{A}_{n,k} = \{f \in \text{Aut}(F_{n,k}) \mid \llbracket f(y_i) \rrbracket = \llbracket y_i \rrbracket \text{ for } 1 \leq i \leq k\}.$$

The map $F_{n,k} \rightarrow F_n$ whose kernel is the normal closure of $\{y_1, \dots, y_k\}$ induces a map $\pi: \mathcal{A}_{n,k} \rightarrow \text{Aut}(F_n)$. The inclusion $\text{Aut}(F_n) \hookrightarrow \mathcal{A}_{n,k}$ whose image consists of automorphisms that fix the y_i pointwise is a right inverse for π , so π is a split surjection. Let $\mathcal{K}_{n,k} = \ker(\pi)$, so we have a split short exact sequence

$$1 \longrightarrow \mathcal{K}_{n,k} \longrightarrow \mathcal{A}_{n,k} \xrightarrow{\pi} \text{Aut}(F_n) \longrightarrow 1. \quad (1)$$

This is the Birman exact sequence for $\text{Aut}(F_n)$ that was studied in [5]. In that paper, the authors proved that $\mathcal{K}_{n,k}$ is finitely generated but not finitely presentable, constructed a simple infinite presentation for it, and computed its abelianization. We say more about these results below.

Remark 1.1. In [5], slightly more general groups $\mathcal{A}_{n,k,l}$ and $\mathcal{K}_{n,k,l}$ were studied. To simplify our exposition, we decided to focus on the case $l = 0$ in this paper.

*Supported in part by NSF grant DMS-1206981

†Supported in part by NSF grant DMS-1255350 and the Alfred P. Sloan Foundation

Analogue for Torelli. Define $\text{IA}_{n,k}$ to be the Torelli subgroup of $\text{Aut}(F_{n,k})$. Set $\mathcal{A}_{n,k}^{\text{IA}} = \mathcal{A}_{n,k} \cap \text{IA}_{n,k}$ and $\mathcal{K}_{n,k}^{\text{IA}} = \mathcal{K}_{n,k} \cap \text{IA}_{n,k}$. The exact sequence (1) restricts to a split short exact sequence

$$1 \longrightarrow \mathcal{K}_{n,k}^{\text{IA}} \longrightarrow \mathcal{A}_{n,k}^{\text{IA}} \longrightarrow \text{IA}_n \longrightarrow 1. \quad (2)$$

This is our Birman exact sequence for IA_n . The purpose of this paper is to prove results for $\mathcal{K}_{n,k}^{\text{IA}}$ that are analogous to the results for $\mathcal{K}_{n,k}$ that we listed above.

Comparing the kernels. The group $\mathcal{K}_{n,k}^{\text{IA}}$ is the kernel of the restriction of the map $\text{Aut}(F_{n,k}) \rightarrow \text{GL}_n(\mathbb{Z})$ to $\mathcal{K}_{n,k}$. The image of this restriction is isomorphic to \mathbb{Z}^{nk} . Indeed, using the generating set for $\mathcal{K}_{n,k}$ constructed in [5] (see below), one can show that with respect to the basis $\{[x_1], \dots, [x_n], [y_1], \dots, [y_k]\}$ for \mathbb{Z}^{n+k} , it consists of matrices of the form

$$\begin{pmatrix} \mathbb{1}_n & 0 \\ A & \mathbb{1}_k \end{pmatrix},$$

where $\mathbb{1}_n$ and $\mathbb{1}_k$ are the $n \times n$ and $k \times k$ identity matrices and A is an arbitrary $k \times n$ integer matrix. We thus have a short exact sequence

$$1 \longrightarrow \mathcal{K}_{n,k}^{\text{IA}} \longrightarrow \mathcal{K}_{n,k} \longrightarrow \mathbb{Z}^{nk} \longrightarrow 1. \quad (3)$$

Unfortunately, it is difficult to use this exact sequence to deduce results about the combinatorial group theory of $\mathcal{K}_{n,k}^{\text{IA}}$ from analogous results for $\mathcal{K}_{n,k}$ (although in a sense we do this in the proof of Theorem B below). For instance, the authors proved in [5] that $\mathcal{K}_{n,k}$ is finitely generated, but this does not directly imply anything about generating sets for $\mathcal{K}_{n,k}^{\text{IA}}$.

Generators. We now turn to our theorems. Set $X = \{x_1, \dots, x_n\}$ and $Y = \{y_1, \dots, y_k\}$. For distinct $z, z' \in X \cup Y$, define $C_{z,z'} \in \text{Aut}(F_{n,k})$ via the formula

$$C_{z,z'}(s) = \begin{cases} z' s (z')^{-1} & \text{if } s = z, \\ s & \text{otherwise.} \end{cases} \quad (s \in X \cup Y).$$

Also, for $z \in X \cup Y$ and $\alpha = \pm 1$ and $v \in F_{n,k}$ in the subgroup generated by $(X \cup Y) \setminus \{z\}$, define $M_{z^\alpha, v} \in \text{Aut}(F_{n,k})$ via the formula

$$M_{z^\alpha, v}(s) = \begin{cases} vs & \text{if } s = z \text{ and } \alpha = 1, \\ sv^{-1} & \text{if } s = z \text{ and } \alpha = -1, \\ s & \text{otherwise.} \end{cases} \quad (s \in X \cup Y).$$

Observe that with this definition we have $M_{z^\alpha, v}(z^\alpha) = vz^\alpha$. The authors proved in [5] that $\mathcal{K}_{n,k}$ is generated by the finite set

$$\{M_{x,y} \mid x \in X, y \in Y\} \cup \{C_{y,z}, C_{z,y} \mid y \in Y, z \in (X \cup Y) \setminus \{y\}\}. \quad (4)$$

Remark 1.2. This is a little different from the generating set given in [5], which includes generators of the form $M_{x^{-1}, y}$ for $x \in X$ and $y \in Y$; however, these are unnecessary here since $M_{x^{-1}, y} = M_{x,y} C_{x,y}^{-1}$. The generators $C_{x,y}$ were not included in the generating set in [5]. We give the above form because it is a little more convenient for our purposes.

The analogue of this for $\mathcal{K}_{n,k}^{\text{IA}}$ is as follows.

Theorem A. *The group $\mathcal{K}_{n,k}^{\text{IA}}$ is generated by the finite set*

$$\{M_{x,[y,z]} \mid x \in X, y \in Y, z \in (X \cup Y) \setminus \{x, y\}\} \cup \{C_{y,z}, C_{z,y} \mid y \in Y, z \in (X \cup Y) \setminus \{y\}\}.$$

The Torelli kernel is not finitely presentable. Though $\mathcal{K}_{n,k}$ is finitely generated, the authors proved in [5] that it is not finitely presentable if $n \geq 2$ and $k \geq 1$; in fact, $H_2(\mathcal{K}_{n,k}; \mathbb{Q})$ is infinite dimensional. If $\mathcal{K}_{n,k}^{\text{IA}}$ were finitely presentable, then one could use the exact sequence (3) to build a finite presentation for $\mathcal{K}_{n,k}$. We deduce that $\mathcal{K}_{n,k}^{\text{IA}}$ is not finitely presentable. Our second main theorem strengthens this observation.

Theorem B. *If $n \geq 2$ and $k \geq 1$, then $H_2(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Q})$ is infinite dimensional. Consequently, $\mathcal{K}_{n,k}^{\text{IA}}$ is not finitely presentable.*

Abelianization. The authors proved in [5] that

$$H_1(\mathcal{K}_{n,k}) = \begin{cases} \mathbb{Z}^{k(k-1)} & \text{if } n = 0, \\ \mathbb{Z}^{2kn} & \text{if } n > 0. \end{cases}$$

These abelian quotients of $\mathcal{K}_{n,k}$ come from two sources.

- The restriction of the map $\text{Aut}(F_{n,k}) \rightarrow \text{GL}_{n+k}(\mathbb{Z})$ to $\mathcal{K}_{n,k}$, which has image \mathbb{Z}^{kn} (see the exact sequence (3)).
- The Johnson homomorphisms, which are homomorphisms

$$\tau: \text{IA}_{n,k} \rightarrow \text{Hom}(\mathbb{Z}^{n+k}, \wedge^2 \mathbb{Z}^{n+k})$$

constructed from the action of $\text{IA}_{n,k}$ on the second nilpotent truncation of $F_{n,k}$ (see §4 below). If $n = 0$, then $\mathcal{K}_{n,k} \subset \text{IA}_{n,k}$ and the restriction of τ to $\mathcal{K}_{n,k}$ has image $\mathbb{Z}^{k(k-1)}$; this provides the entire abelianization. If $n > 0$, then $\mathcal{K}_{n,k}$ does not lie in $\text{IA}_{n,k}$ and we cannot use the Johnson homomorphism directly; however, in [5] we construct a modified version of it which is defined on $\mathcal{K}_{n,k}$ and has image \mathbb{Z}^{kn} .

The analogue of these calculations for $\mathcal{K}_{n,k}^{\text{IA}}$ is as follows.

Theorem C. *The group $H_1(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Z})$ is free abelian of rank*

$$n(n-1)k + n \binom{k}{2} + 2nk + k(k-1).$$

The abelianization map is given by the restriction of the Johnson homomorphism to $\mathcal{K}_{n,k}^{\text{IA}}$.

Remark 1.3. Theorem C is related to the fact that the Johnson homomorphism gives the abelianization of $\text{IA}_{n,k}$, a theorem which was proved independently by Farb [8], Cohen–Pakianathan [4], and Kawazumi [11].

Finite L-presentation. Our final theorem gives an infinite presentation for the group $\mathcal{K}_{n,1}^{\text{IA}}$ by generators and relations. Though this result may appear technical, it is actually the most important theorem in this paper for our study in [6] of $H_2(\text{IA}_n; \mathbb{Z})$. To simplify our notation, we will write the generators of $F_{n,1}$ as $\{x_1, \dots, x_n, y\}$. The generators for our presentation will be the finite set

$$S_K := \{M_{x^\alpha, [y^\beta, z^\gamma]} \mid x \in X, z \in X \setminus \{x\}, \alpha, \beta, \gamma = \pm 1\} \cup \{C_{y,x}, C_{x,y} \mid x \in X\}.$$

This is larger than the generating set given by Theorem A; using S_K will simplify our relations. By Theorem B, the set of relations will have to be infinite. They will be generated from a finite list of relations by a simple recursive procedure which we will encode using the notion of an L-presentation, which was introduced by Bartholdi [1].

A *finite L-presentation* for a group G is a triple $\langle S \mid R^0 \mid E \rangle$ as follows.

- S is a finite generating set for G .
- R^0 is a finite subset of the free group $F(S)$ on S consisting of relations for G . It is not necessarily a complete set of relations.
- E is a finite subset of $\text{End}(F(S))$.

This triple must satisfy the following. Let $M \subset \text{End}(F(S))$ be the monoid generated by E . Define $R = \{f(r) \mid f \in M, r \in R^0\}$. Then we require that $G = \langle S \mid R \rangle$. Each element of E descends to an element of $\text{End}(G)$; we call the resulting subset $\tilde{E} \subset \text{End}(G)$ the *induced endomorphisms* of our L-presentation.

In this paper, the induced endomorphisms of our L-presentations will actually be automorphisms. Thus in the context of this paper one should think of an L-presentation as a group presentation incorporating certain symmetries of a group. Here is an example.

Example. Fix $\ell \geq 1$. Let $S = \{z_i \mid i \in \mathbb{Z}/\ell\}$ and $R^0 = \{z_0^2\}$. Let $\psi: F(S) \rightarrow F(S)$ be the homomorphism defined via the formula $\psi(z_i) = z_{i+1}$. Then $\langle S \mid R^0 \mid \{\psi\} \rangle$ is a finite L-presentation for the free product of ℓ copies of $\mathbb{Z}/2$.

We now return to the automorphism group of a free group. The group $\mathcal{K}_{n,1}$ is a normal subgroup of $\mathcal{A}_{n,1}$, so $\mathcal{A}_{n,1}$ acts on $\mathcal{K}_{n,1}$ by conjugation. In [5], the authors constructed a finite L-presentation for $\mathcal{K}_{n,1}$ whose set of induced endomorphisms generates

$$\mathcal{A}_{n,1} \subset \text{Aut}(\mathcal{K}_{n,1}) \subset \text{End}(\mathcal{K}_{n,1}).$$

The group $\mathcal{K}_{n,1}^{\text{IA}}$ is also a normal subgroup of $\mathcal{A}_{n,1}$, and hence $\mathcal{A}_{n,1}$ acts on $\mathcal{K}_{n,1}^{\text{IA}}$ by conjugation. Our final main theorem is as follows.

Theorem D. *For all $n \geq 2$, there exists a finite L-presentation $\mathcal{K}_{n,1}^{\text{IA}} = \langle S_K \mid R_K^0 \mid E_K \rangle$ whose set of induced endomorphisms generates $\mathcal{A}_{n,1} \subset \text{Aut}(\mathcal{K}_{n,1}^{\text{IA}}) \subset \text{End}(\mathcal{K}_{n,1}^{\text{IA}})$.*

See the tables in §6 for explicit lists enumerating R_K^0 and E_K .

Verifying the L-presentation. We obtained the list of relations in R_K^0 by starting with a guess of a presentation and then trying to run the following proof sketch. Every time it

failed, that failure revealed a relation we had missed. Let Γ_n be the group given by the purported presentation in Theorem D. There is a natural surjection $\Gamma_n \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$ that we want to prove is an isomorphism. As we will see in §6 below, we have a short exact sequence

$$1 \longrightarrow \mathcal{K}_{n,1}^{\text{IA}} \longrightarrow \mathcal{A}_{n,1} \xrightarrow{\rho} \mathbb{Z}^n \rtimes \text{Aut}(F_n) \longrightarrow 1, \quad (5)$$

where $\text{Aut}(F_n)$ acts on \mathbb{Z}^n via the natural surjection $\text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z})$. The heart of our proof is the construction of a similar extension Δ_n of $\mathbb{Z}^n \rtimes \text{Aut}(F_n)$ by Γ_n which fits into a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Gamma_n & \longrightarrow & \Delta_n & \longrightarrow & \mathbb{Z}^n \rtimes \text{Aut}(F_n) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow = \\ 1 & \longrightarrow & \mathcal{K}_{n,1}^{\text{IA}} & \longrightarrow & \mathcal{A}_{n,1} & \xrightarrow{\rho} & \mathbb{Z}^n \rtimes \text{Aut}(F_n) \longrightarrow 1. \end{array} \quad (6)$$

This construction is very involved; the exact sequences in (6) do not split, and constructing group extensions with nonabelian kernels is delicate. We will say more about how we do this in the next paragraph. In any case, once we have constructed (6) we can use a known presentation of $\mathcal{A}_{n,1}$ due to Jensen–Wahl [10] to show that the map $\Delta_n \rightarrow \mathcal{A}_{n,1}$ is an isomorphism. The five-lemma then implies that the map $\Gamma_n \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$ is an isomorphism, as desired.

The trouble with non-split extensions. If

$$1 \longrightarrow K \longrightarrow G \longrightarrow Q \longrightarrow 1 \quad (7)$$

is a group extension and presentations of Q and K are known, then it is straightforward to construct a presentation of G . However, when constructing the group Δ_n in (6) we have to confront a serious problem, namely we need to first verify that the desired extension exists. To put it another way, it is clear how to combine a known presentation of $\mathbb{Z}^n \rtimes \text{Aut}(F_n)$ with our purported presentation for Γ_n to form a group Δ_n which fits into a commutative diagram

$$\begin{array}{ccccccc} \Gamma_n & \longrightarrow & \Delta_n & \longrightarrow & \mathbb{Z}^n \rtimes \text{Aut}(F_n) & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow = \\ 1 & \longrightarrow & \mathcal{K}_{n,1}^{\text{IA}} & \longrightarrow & \mathcal{A}_{n,1} & \xrightarrow{\rho} & \mathbb{Z}^n \rtimes \text{Aut}(F_n) \longrightarrow 1. \end{array}$$

However, it is difficult to show that the map $\Gamma_n \rightarrow \Delta_n$ is injective. Standard techniques show that proving the existence of the extension (7) is equivalent to constructing a sort of “nonabelian K -valued 2-cocycle” on Q ; see [3, §IV.6]. Such a 2-cocycle is *not* determined by its values on generators for Q . This holds even in the simple case of a central extension; the general case is even worse. It is therefore very difficult to construct such a 2-cocycle using generators and relations.

But the extensions in (6) we are trying to understand are very special. While they do not split, there do exist “partial splittings”, namely homomorphisms $\iota_1: \mathbb{Z}^n \rightarrow \mathcal{A}_{n,1}$ and $\iota_2: \text{Aut}(F_n) \rightarrow \mathcal{A}_{n,1}$ such that $\rho \circ \iota_1 = \text{id}$ and $\rho \circ \iota_2 = \text{id}$. Letting $\Lambda_1 = \rho^{-1}(\mathbb{Z}^n)$ and $\Lambda_2 = \rho^{-1}(\text{Aut}(F_n))$ we therefore have $\Lambda_1 \cong \mathcal{K}_{n,1}^{\text{IA}} \rtimes \mathbb{Z}^n$ and $\Lambda_2 \cong \mathcal{K}_{n,1}^{\text{IA}} \rtimes \text{Aut}(F_n)$. The data

needed to combine Λ_1 and Λ_2 into a group $\mathcal{A}_{n,1}$ that fits into (5) is what we will call a “twisted bilinear map” from $\text{Aut}(F_n) \times \mathbb{Z}^n$ to $\mathcal{K}_{n,1}^{\text{IA}}$. The definition is complicated, so to give the flavor of it in this introduction we will discuss a simpler situation.

Splicing together direct products. Let A and B and K be abelian groups. We want to construct a not necessarily abelian group G with the following property.

- There is a short exact sequence

$$1 \longrightarrow K \longrightarrow G \xrightarrow{\rho} A \times B \longrightarrow 1$$

together with homomorphisms $\iota_1: A \rightarrow G$ and $\iota_2: B \rightarrow G$ such that $\rho^{-1}(A)$ and $\rho^{-1}(B)$ are the internal direct products $K \times \iota_1(A)$ and $K \times \iota_2(B)$, respectively.

Given this data, we can define a set map $\lambda: A \times B \rightarrow K$ via the formula

$$\lambda(a, b) = [\iota_1(a), \iota_2(b)] \quad (a \in A, b \in B);$$

here the bracket is the commutator bracket in G . It is easy to see that λ is bilinear. Conversely, given a bilinear map $\lambda: A \times B \rightarrow K$ we can construct a group G with the above properties by letting G consist of all triples $(k, b, a) \in K \times B \times A$ with the multiplication

$$(k, b, a)(k', b', a') = (k + k' + \phi(a, b'), b + b', a + a').$$

The bilinearity of ϕ is needed for this multiplication to be associative.

Adding the twisting. The groups we are interested in fit into semidirect products, so we will have to incorporate the various group actions into our bilinear maps. The key property of the resulting theory of twisted bilinear maps is that (unlike general 2-cocycles but like ordinary bilinear maps) they are determined by their values on generators. Letting Γ_n be the group in (6), we will therefore be able to use combinatorial group theory to construct an appropriate twisted bilinear map $\mathbb{Z}^n \times \text{Aut}(F_n) \rightarrow \Gamma_n$ that behaves like the twisted bilinear map $\mathbb{Z}^n \times \text{Aut}(F_n) \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$ that determines $\mathcal{A}_{n,1}$. This will allow us to construct the group Δ_n fitting into (6) and complete the proof of Theorem D.

Outline. We prove Theorem A in §2, Theorem B in §3, and Theorem C in §4. Preliminaries for the proof Theorem D are in §5, and the proof itself appears in §6. The proof of Theorem D depends on computer calculations that are described in §7.

2 Generators

In this section, we prove Theorem A. Letting $X = \{x_1, \dots, x_n\}$ and $Y = \{y_1, \dots, y_k\}$, recall that this theorem asserts that the set

$T := \{M_{x,[y,z]} \mid x \in X, y \in Y, z \in (X \cup Y) \setminus \{x, y\}\} \cup \{C_{y,z}, C_{z,y} \mid y \in Y, z \in (X \cup Y) \setminus \{y\}\}$
generates $\mathcal{K}_{n,k}^{\text{IA}}$.

$t \in T$	sts^{-1}	$s^{-1}ts$
$M_{x_a, [y_d, x_b]}$	$C_{x_a, y_d} M_{x_a, [y_d, x_b]} C_{x_a, y_d}^{-1}$	$C_{x_a, y_d}^{-1} M_{x_a, [y_d, x_b]} C_{x_a, y_d}$
$M_{x_b, [y_d, x_a]}$	$C_{x_a, y_d} M_{x_b, [y_d, x_a]} C_{x_a, y_d}^{-1}$	$C_{x_a, y_d}^{-1} M_{x_b, [y_d, x_a]} C_{x_a, y_d}$
$M_{x_a, [y_e, x_b]}$	$C_{x_a, y_d} M_{x_a, [y_e, x_b]} C_{x_a, y_d}^{-1}$	$C_{x_a, y_d}^{-1} M_{x_a, [y_e, x_b]} C_{x_a, y_d}$
$M_{x_b, [y_e, x_a]}$	$C_{x_b, y_d}^{-1} M_{x_b, [y_e, x_a]} C_{x_b, y_d} M_{x_b, [y_e, y_d]}$	$C_{x_b, y_d} M_{x_b, [y_e, x_a]} M_{x_b, [y_d, y_e]} C_{x_b, y_d}^{-1}$
$M_{x_a, [y_d, y_e]}$	$C_{x_a, y_d} M_{x_a, [y_d, y_e]} C_{x_a, y_d}^{-1}$	$C_{x_a, y_d}^{-1} M_{x_a, [y_d, y_e]} C_{x_a, y_d}$
$M_{x_a, [y_e, y_f]}$	$C_{x_a, y_d} M_{x_a, [y_e, y_f]} C_{x_a, y_d}^{-1}$	$C_{x_a, y_d}^{-1} M_{x_a, [y_e, y_f]} C_{x_a, y_d}$
C_{y_d, y_e}	$C_{x_a, y_d} M_{x_a, [y_d, y_e]} C_{x_a, y_d}^{-1} C_{y_d, y_e}$	$M_{x_a, [y_e, y_d]} C_{y_d, y_e}$
C_{y_d, x_a}	$C_{x_a, y_d} C_{y_d, x_a}$	$C_{x_a, y_d}^{-1} C_{y_d, x_a}$
C_{y_d, x_b}	$C_{y_d, x_b} C_{x_a, y_d} C_{y_d, x_b}^{-1} M_{x_a, [y_d, x_b]} C_{y_d, x_b} C_{x_a, y_d}$	$C_{x_a, y_d} C_{y_d, x_b}^{-1} M_{x_a, [y_d, x_b]} C_{y_d, x_b}$
C_{y_e, x_a}	$C_{y_e, x_a} C_{y_e, y_d}$	$C_{y_e, x_a} C_{y_e, y_d}^{-1}$
C_{x_a, y_e}	$C_{x_a, y_e} C_{x_a, y_d} M_{x_a, [y_e, y_d]} C_{x_a, y_d}^{-1}$	$C_{x_a, y_e} M_{x_a, [y_d, y_e]}$

Table 1: Fix $s = M_{x_a, y_d}$. This table shows how to write sts^{-1} and $s^{-1}ts$ as a word in T for all $t \in T$. Basis elements with distinct subscripts are assumed to be distinct. If a formula is not listed, then $sts^{-1} = s^{-1}ts = t$. All these formulas can be easily proved by checking the effect of the indicated automorphisms on a basis for the free group.

Proof of Theorem A. The key is the exact sequence

$$1 \longrightarrow \mathcal{K}_{n,k}^{\text{IA}} \longrightarrow \mathcal{K}_{n,k} \xrightarrow{\rho} \mathbb{Z}^{nk} \longrightarrow 1$$

discussed in the introduction (see (3)). Define

$$S_1 = \{M_{x,y} \mid x \in X, y \in Y\} \quad \text{and} \quad S_2 = \{C_{y,z}, C_{z,y} \mid y \in Y, z \in (X \cup Y) \setminus \{y\}\}.$$

As we discussed in the introduction (see the equation (4) and the remark following it), the authors proved in [5] that $S_1 \cup S_2$ generates $\mathcal{K}_{n,k}$. We have $S_2 \subset \mathcal{K}_{n,k}^{\text{IA}} = \ker(\rho)$. Also, ρ maps the elements of S_1 to a basis of \mathbb{Z}^{nk} . We therefore see that \mathbb{Z}^{nk} is the quotient of $\mathcal{K}_{n,k}$ by the normal closure of the set $S'_1 \cup S_2$, where

$$\begin{aligned} S'_1 &= \{[s, s'] \mid s, s' \in S_1\} \\ &= \{[M_{x,y}, M_{x,y'}] \mid x \in X, y, y' \in Y, y \neq y'\} \\ &\quad \cup \{[M_{x,y}, M_{x',y'}] \mid x, x' \in X, y, y' \in Y, x \neq x'\} \\ &= \{M_{x, [y, y']} \mid x \in X, y, y' \in Y, y \neq y'\}. \end{aligned}$$

Here we are using the fact that $[M_{x,y}, M_{x',y}] = 1$ for $x, x' \in X$ and $y \in Y$ with $x \neq x'$. Since $S'_1 \cup S_2 \subset T$, we conclude that T normally generates $\mathcal{K}_{n,k}^{\text{IA}}$.

Letting $G \subset \mathcal{K}_{n,k}^{\text{IA}}$ be the subgroup generated by T , it is therefore enough to prove that G is a normal subgroup. To do this, it is enough to prove that for $s \in S_1 \cup S_2$ and $t \in T$, we have $sts^{-1} \in G$ and $s^{-1}ts \in G$. In fact, since $S_2 \subset T$ it is enough to do this for $s \in S_1$. The identities that show this are in Table 1. \square

3 The Torelli kernel is not finitely presentable

In this section, we prove Theorem B. Recall that this theorem asserts that $H_2(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Q})$ is infinite dimensional when $n \geq 2$ and $k \geq 1$.

Proof of Theorem B. Consider the Hochschild–Serre spectral sequence associated to the short exact sequence

$$1 \longrightarrow \mathcal{K}_{n,k}^{\text{IA}} \longrightarrow \mathcal{K}_{n,k} \xrightarrow{\rho} \mathbb{Z}^{nk} \longrightarrow 1$$

discussed in the introduction (see (3)). It is of the form

$$E_{pq}^2 = H_p(\mathbb{Z}^{nk}; H_q(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Q})) \Rightarrow H_{p+q}(\mathcal{K}_{n,k}; \mathbb{Q}).$$

The authors proved in [5] that $H_2(\mathcal{K}_{n,k}; \mathbb{Q})$ is infinite dimensional, so at least one of E_{20}^2 and E_{11}^2 and E_{02}^2 must be infinite dimensional. Clearly $E_{20}^2 = H_2(\mathbb{Z}^{nk}; \mathbb{Q})$ is finite dimensional. Also, Theorem A implies that $H_1(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Q})$ is finite-dimensional, so $E_{11}^2 = H_1(\mathbb{Z}^{nk}; H_1(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Q}))$ is finite dimensional. We conclude that $E_{02}^2 = H_0(\mathbb{Z}^{nk}; H_2(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Q}))$ is infinite dimensional, so $H_2(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Q})$ is infinite dimensional, as desired. \square

4 Abelianization

In this section, we prove Theorem C. Recall that this theorem asserts that $H_1(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Z})$ is free abelian and that the abelianization map is given by the restriction of the Johnson homomorphism to $\mathcal{K}_{n,k}^{\text{IA}}$. The theorem also gives the rank of the abelianization of $\mathcal{K}_{n,k}^{\text{IA}}$ as a polynomial in n and k .

Proof of Theorem C. We begin by recalling the definition of the Johnson homomorphism; see [12] for more details and references. Let $\pi: [F_{n,k}, F_{n,k}] \rightarrow \bigwedge^2 \mathbb{Z}^{n+k}$ be the projection whose kernel is $[F_{n,k}, [F_{n,k}, F_{n,k}]]$. This map satisfies $\pi([z, z']) = [z] \wedge [z']$ for $z, z' \in F_{n,k}$; here $[z], [z'] \in \mathbb{Z}^{n+k}$ are the images of z and z' in the abelianization of $F_{n,k}$. The Johnson homomorphism is then a homomorphism

$$\tau: \text{IA}_{n,k} \rightarrow \text{Hom}(\mathbb{Z}^{n+k}, \bigwedge^2 \mathbb{Z}^{n+k})$$

that satisfies the formula

$$\tau(f)([z]) = \pi(f(z)z^{-1}) \quad (f \in \text{IA}_{n,k}, z \in F_{n,k}).$$

Letting $X = \{x_1, \dots, x_n\}$ and $Y = \{y_1, \dots, y_k\}$, the Johnson homomorphism has the following effect on the basic elements of $\text{IA}_{n,k}$ defined in the introduction.

- For distinct $z, z' \in X \cup Y$, we have

$$\tau(C_{z,z'})([w]) = \begin{cases} [z] \wedge [z'] & \text{if } w = z, \\ 0 & \text{otherwise} \end{cases} \quad (w \in X \cup Y).$$

- For distinct $z, z', z'' \in X \cup Y$, we have

$$\tau(M_{z,[z',z'']})([w]) = \begin{cases} [z'] \wedge [z''] & \text{if } w = z, \\ 0 & \text{otherwise} \end{cases} \quad (w \in X \cup Y).$$

Set

$$\begin{aligned} T = & \{M_{x,[y,x']} \mid x \in X, y \in Y, x' \in X \setminus \{x\}\} \\ & \cup \{M_{x,[y_a,y_b]} \mid x \in X, 1 \leq a < b \leq k\} \\ & \cup \{C_{y,z}, C_{z,y} \mid y \in Y, z \in (X \cup Y) \setminus \{y\}\}. \end{aligned}$$

Since $M_{x,[y_b,y_a]} = M_{x,[y_a,y_b]}^{-1}$ for $x \in X$ and $1 \leq a < b \leq k$, Theorem A implies that T generates $\mathcal{K}_{n,k}^{\text{IA}}$. Examining the above formulas, we see that τ takes T injectively to a linearly independent subset of the free abelian group $\text{Hom}(\mathbb{Z}^{n+k}, \bigwedge^2 \mathbb{Z}^{n+k})$. This implies that if u is an element of the free group $F(T)$ on T which maps to a relation in $\mathcal{K}_{n,k}^{\text{IA}}$, then $u \in [F(T), F(T)]$ (otherwise, τ would take the image of u in $\mathcal{K}_{n,k}^{\text{IA}}$ to a nontrivial element of $\text{Hom}(\mathbb{Z}^{n+k}, \bigwedge^2 \mathbb{Z}^{n+k})$). We conclude that τ induces the abelianization of $\mathcal{K}_{n,k}^{\text{IA}}$ and that $H_1(\mathcal{K}_{n,k}^{\text{IA}}; \mathbb{Z}) = \mathbb{Z}^{|T|}$. This is free abelian of rank

$$|T| = n(n-1)k + n \binom{k}{2} + 2nk + k(k-1). \quad \square$$

5 Preliminaries for the proof of Theorem D

The rest of this paper is devoted to proving Theorem D, which gives a finite L-presentation for $\mathcal{K}_{n,1}^{\text{IA}}$. This section contains three subsections of preliminaries: §5.1 constructs a needed exact sequence, §5.2 discusses twisted bilinear maps, and §5.3 discusses presentations for some related groups.

To simplify our notation, we will set $y = y_1$, so $\{x_1, \dots, x_n, y\}$ is the basis for $F_{n,1}$. When writing matrices in $\text{GL}_{n+1}(\mathbb{Z})$, we will always use the basis $\{[x_1], \dots, [x_n], [y]\}$ for \mathbb{Z}^{n+1} .

5.1 Relating the two kernels

This is the first of three preliminary sections for the proof of Theorem D. In it, we construct the exact sequence

$$1 \longrightarrow \mathcal{K}_{n,1}^{\text{IA}} \longrightarrow \mathcal{A}_{n,1} \xrightarrow{\rho} \mathbb{Z}^n \rtimes \text{Aut}(F_n) \longrightarrow 1$$

discussed in the introduction (see Lemma 5.3 below). We first address an irritating technical point. Throughout this paper, all group actions are left actions. In particular, elements of \mathbb{Z}^n will be regarded as column vectors and matrices in $\text{GL}_n(\mathbb{Z})$ act on these column vectors on the left (we have already silently used this convention when we wrote matrices). However, it turns out that the action of $\text{Aut}(F_n)$ on \mathbb{Z}^n in the semidirect product appearing in the above exact sequence is induced by the natural *right* action of $\text{GL}_n(\mathbb{Z})$ on row vectors. We do not wish to mix up right and left actions, so we convert this into a left action and define $\mathbb{Z}^n \rtimes_r \text{GL}_n(\mathbb{Z})$ to be the semidirect product associated to the action of $\text{GL}_n(\mathbb{Z})$ on \mathbb{Z}^n defined by the formula

$$M \cdot z = (M^{-1})^t z \quad (M \in \text{GL}_n(\mathbb{Z}), z \in \mathbb{Z}^n).$$

To understand this formula, observe that $(M^{-1})^t z$ is the transpose of $(z^t)M^{-1}$; the inverse appears because we are converting a right action into a left action. We then have the following.

Lemma 5.1. *The stabilizer subgroup $(\mathrm{GL}_{n+1}(\mathbb{Z}))_{[y]}$ is isomorphic to $\mathbb{Z}^n \rtimes_r \mathrm{GL}_n(\mathbb{Z})$.*

Proof. We define a homomorphism $\psi: (\mathrm{GL}_{n+1}(\mathbb{Z}))_{[y]} \rightarrow \mathbb{Z}^n \rtimes_r \mathrm{GL}_n(\mathbb{Z})$ as follows. Consider $M \in (\mathrm{GL}_{n+1}(\mathbb{Z}))_{[y]}$. There exist $\widehat{M} \in \mathrm{GL}_n(\mathbb{Z})$ and $\overline{M} \in \mathbb{Z}^n$ such that

$$M = \left(\begin{array}{c|c} \widehat{M} & 0 \\ \hline \overline{M}^t & 1 \end{array} \right);$$

here we are using our convention that elements of \mathbb{Z}^n are column vectors, so the transpose \overline{M}^t of $\overline{M} \in \mathbb{Z}^n$ is a row. We then define $\psi(M) = \left(\left(\widehat{M}^{-1} \right)^t \overline{M}, \widehat{M} \right)$. To see that this is a homomorphism, observe that for $M_1, M_2 \in (\mathrm{GL}_{n+1}(\mathbb{Z}))_{[y]}$ we have

$$M_1 M_2 = \left(\begin{array}{c|c} \widehat{M}_1 & 0 \\ \hline \overline{M}_1^t & 1 \end{array} \right) \left(\begin{array}{c|c} \widehat{M}_2 & 0 \\ \hline \overline{M}_2^t & 1 \end{array} \right) = \left(\begin{array}{c|c} \widehat{M}_1 \widehat{M}_2 & 0 \\ \hline \overline{M}_1^t \widehat{M}_2 + \overline{M}_2^t & 1 \end{array} \right) = \left(\begin{array}{c|c} \widehat{M}_1 \widehat{M}_2 & 0 \\ \hline (\widehat{M}_2^t \overline{M}_1 + \overline{M}_2^t) & 1 \end{array} \right),$$

and hence

$$\begin{aligned} \psi(M_1) \psi(M_2) &= \left(\left(\widehat{M}_1^{-1} \right)^t \overline{M}_1, \widehat{M}_1 \right) \left(\left(\widehat{M}_2^{-1} \right)^t \overline{M}_2, \widehat{M}_2 \right) \\ &= \left(\left(\widehat{M}_1^{-1} \right)^t \overline{M}_1 + \left(\widehat{M}_1^{-1} \right)^t \left(\widehat{M}_2^{-1} \right)^t \overline{M}_2, \widehat{M}_1 \widehat{M}_2 \right) \\ &= \left(\left(\widehat{M}_1^{-1} \right)^t \left(\widehat{M}_2^{-1} \right)^t \left(\widehat{M}_2^t \overline{M}_1 + \overline{M}_2 \right), \widehat{M}_1 \widehat{M}_2 \right) \\ &= \psi(M_1 M_2). \end{aligned}$$

That ψ is a bijection is obvious. □

Remark 5.2. There is an isomorphism between $\mathbb{Z}^n \rtimes_r \mathrm{GL}_n(\mathbb{Z})$ and the semidirect product of \mathbb{Z}^n and $\mathrm{GL}_n(\mathbb{Z})$ with respect to the standard left action of $\mathrm{GL}_n(\mathbb{Z})$ on \mathbb{Z}^n . However, this isomorphism acts as the inverse transpose on the $\mathrm{GL}_n(\mathbb{Z})$ factor, and to keep our formulas from getting out of hand we want to not change this factor. Throughout this paper, we will use the explicit isomorphism described in the proof of Lemma 5.1.

Define $\mathbb{Z}^n \rtimes_r \mathrm{Aut}(F_n)$ to be the semidirect product induced by the action of $\mathrm{Aut}(F_n)$ on \mathbb{Z}^n obtained by composing the projection $\mathrm{Aut}(F_n) \rightarrow \mathrm{GL}_n(\mathbb{Z})$ with the action of $\mathrm{GL}_n(\mathbb{Z})$ on \mathbb{Z}^n discussed above. We then have the following lemma, which is the main result of this section.

Lemma 5.3. *There is a short exact sequence*

$$1 \longrightarrow \mathcal{K}_{n,1}^{IA} \longrightarrow \mathcal{A}_{n,1} \xrightarrow{\rho} \mathbb{Z}^n \rtimes_r \mathrm{Aut}(F_n) \longrightarrow 1.$$

Also, there exist homomorphisms $\iota_1: \mathrm{Aut}(F_n) \rightarrow \mathcal{A}_{n,1}$ and $\iota_2: \mathbb{Z}^n \rightarrow \mathcal{A}_{n,1}$ such that $\rho \circ \iota_1$ and $\rho \circ \iota_2$ are the standard inclusions of $\mathrm{Aut}(F_n)$ and \mathbb{Z}^n into $\mathbb{Z}^n \rtimes_r \mathrm{Aut}(F_n)$ respectively.

Proof. Let $\pi_1: \mathcal{A}_{n,1} \rightarrow \text{Aut}(F_n)$ be the map induced by the projection $F_{n,1} \rightarrow F_n$ whose kernel is normally generated by y . Also, let $\pi_2: \mathcal{A}_{n,1} \rightarrow \mathbb{Z}^n \rtimes_r \text{GL}_n(\mathbb{Z})$ be the composition

$$\mathcal{A}_{n,1} \longrightarrow (\text{GL}_{n+1}(\mathbb{Z}))_{[y]} \xrightarrow{\cong} \mathbb{Z}^n \rtimes_r \text{GL}_n(\mathbb{Z}),$$

where the second map is the isomorphism given by Lemma 5.1. By definition, $\mathcal{K}_{n,1} = \ker(\pi_1)$ and $\mathcal{A}_{n,1}^{\text{IA}} = \ker(\pi_2)$. Recalling that $\mathcal{K}_{n,1}^{\text{IA}} = \mathcal{K}_{n,1} \cap \mathcal{A}_{n,1}^{\text{IA}}$, it follows that $\mathcal{K}_{n,1}^{\text{IA}} = \ker(\rho)$, where ρ is the composition

$$\mathcal{A}_{n,1} \xrightarrow{\pi_1 \oplus \pi_2} (\text{Aut}(F_n)) \oplus (\mathbb{Z}^n \rtimes_r \text{GL}_n(\mathbb{Z})).$$

Let $\eta: \text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z})$ be the natural projection. The image of ρ is contained in the subgroup

$$\{(f, (z, \eta(f))) \mid f \in \text{Aut}(F_n), z \in \mathbb{Z}^n\} \subset (\text{Aut}(F_n)) \oplus (\mathbb{Z}^n \rtimes_r \text{GL}_n(\mathbb{Z})),$$

which is clearly isomorphic to $\mathbb{Z}^n \rtimes_r \text{Aut}(F_n)$. We can therefore regard ρ as a homomorphism $\rho: \mathcal{A}_{n,1} \rightarrow \mathbb{Z}^n \rtimes_r \text{Aut}(F_n)$, and we have an exact sequence

$$1 \longrightarrow \mathcal{K}_{n,1}^{\text{IA}} \longrightarrow \mathcal{A}_{n,1} \xrightarrow{\rho} \mathbb{Z}^n \rtimes_r \text{Aut}(F_n).$$

Let $\iota_1: \text{Aut}(F_n) \rightarrow \mathcal{A}_{n,1}$ be the evident inclusion whose image is the stabilizer subgroup $(\mathcal{A}_{n,1})_y$ and let $\iota_2: \mathbb{Z}^n \rightarrow \mathcal{A}_{n,1}$ be the map defined via the formula

$$\iota_2(z_1, \dots, z_n) = M_{x_1, y}^{z_1} M_{x_2, y}^{z_2} \cdots M_{x_n, y}^{z_n} \quad (z_1, \dots, z_n \in \mathbb{Z}),$$

where the automorphisms $M_{x_i, y}$ are as in the introduction. The map ι_2 is a homomorphism because the $M_{x_i, y}$ commute. It is clear that $\rho \circ \iota_1 = \text{id}$ and $\rho \circ \iota_2 = \text{id}$. This implies that ρ is surjective, and the lemma follows. \square

5.2 Twisted bilinear maps and group extensions

This is the second section containing preliminaries for the proof of Theorem D. In it, we discuss the theory of twisted bilinear maps alluded to in the introduction. Throughout this section, let A and B and K be groups equipped with the following left actions.

- The group A acts on B ; for $a \in A$ and $b \in B$, we will write ${}^a b$ for the image of b under the action of a .
- The groups A and B both act on K . For $k \in K$ and $a \in A$ and $b \in B$, we will write $\alpha_a(k)$ and $\beta_b(k)$ for the images of k under the actions of a and b , respectively.

A *twisted bilinear map* from $A \times B$ to K is a set map $\lambda: A \times B \rightarrow K$ satisfying the following three properties.

- TB1. For all $a \in A$ and $b_1, b_2 \in B$, we have $\lambda(a, b_1 b_2) = \lambda(a, b_1) \cdot \beta_{a b_1}(\lambda(a, b_2))$.
 TB2. For all $a_1, a_2 \in A$ and $b \in B$, we have $\lambda(a_1 a_2, b) = \alpha_{a_1}(\lambda(a_2, b)) \cdot \lambda(a_1, {}^{a_2} b)$.
 TB3. For all $a \in A$ and $b \in B$ and $k \in K$, we have $\lambda(a, b) \cdot \beta_{a b}(\alpha_a(k)) \cdot \lambda(a, b)^{-1} = \alpha_a(\beta_b(k))$.

Observe that this reduces to the definition of a bilinear map if all the actions are trivial and all the groups are abelian. The key example is as follows.

Example. Consider a short exact sequence of groups

$$1 \longrightarrow K \longrightarrow G \xrightarrow{\rho} B \rtimes A \longrightarrow 1$$

together with homomorphisms $\iota_1: A \rightarrow G$ and $\iota_2: B \rightarrow G$ such that $\rho \circ \iota_1$ and $\rho \circ \iota_2$ are the standard inclusions of A and B in $B \rtimes A$ respectively. Observe that this implies that $\rho^{-1}(A)$ and $\rho^{-1}(B)$ are the internal semidirect products $K \rtimes_{\iota_1} A$ and $K \rtimes_{\iota_2} B$, respectively. Let ${}^a b$ be the action $a \cdot b$ defining the semidirect product $B \rtimes A$, and define actions of A and B on K by

$$\alpha_a(k) = \iota_1(a) \cdot k \cdot \iota_1(a)^{-1} \quad \text{and} \quad \beta_b(k) = \iota_2(b) \cdot k \cdot \iota_2(b)^{-1}$$

for all $a \in A$ and $b \in B$ and $k \in K$. Define a set map $\lambda: A \times B \rightarrow K$ via the formula

$$\lambda(a, b) = \iota_1(a) \cdot \iota_2(b) \cdot \iota_1(a)^{-1} \iota_2({}^a b)^{-1}.$$

Note that this is a kind of “twisted commutator” map; it reduces to the commutator bracket if the action of A on B is trivial. Given these definitions, an easy algebraic juggle shows that λ is a twisted bilinear map. We will say that λ is the twisted bilinear map *associated* to G and ι_1 and ι_2 .

Remark 5.4. We take a moment to explain the different aspects of the definition of a twisted bilinear map. For a fixed $a \in A$, we can twist the action of B on K by the action of A on B by a , to get an action $b \cdot k = \beta_{a_b}(k)$. Property TB1 simply states that the function $\lambda(a, \cdot): B \rightarrow K$ is a crossed homomorphism with respect to this action twisted by a .

Property TB2 is similar, but involves two kinds of twisting. The set of functions $B \rightarrow K$ is a group with the pointwise product. The group A acts on this group in two ways. The first is by post-composition: for $a \in A$ and $f: B \rightarrow K$, define $a \cdot f$ by $(a \cdot f)(b) = \alpha_a(f(b))$. The second is by pre-composition, and is a right action: for a, f as above, define $f \cdot a$ by $(f \cdot a)(b) = f({}^a b)$. So property TB2 says that λ is like a crossed homomorphism from A to the group of functions $B \rightarrow K$, but simultaneously twisted by both of these actions.

Property TB3 can be explained in the context of Example 5.2. As usual in group extensions, there is a well defined *outer* action of $B \rtimes A$ on K : to act on k by (b, a) , lift (b, a) to G and conjugate k by this lift. The conjugate of k depends on the choice of lift, but the resulting map $B \rtimes A \rightarrow \text{Out}(K)$ is well defined. Using our maps ι_1 and ι_2 , we have two ways to build lifts. By the definition of the product in $B \rtimes A$, we have

$$(1, a)(b, 1) = ({}^a b, a) = ({}^a b, 1)(1, a).$$

So we may view the outer action of $({}^a b, a)$ on K as coming from conjugation either by $\iota_1(a)\iota_2(b)$ or by $\iota_2({}^a b)\iota_1(a)$. These conjugations are given by $k \mapsto \alpha_a(\beta_b(k))$ and $k \mapsto \beta_{a_b}(\alpha_a(k))$ respectively. Since they define the same outer automorphism, they differ by conjugation by some element; TB3 says that $\lambda(a, b)$ is such an element.

The following theorem shows that every twisted bilinear map is associated to some group extension.

Theorem 5.5. *Let the groups and actions be as above and let $\lambda: A \times B \rightarrow K$ be a twisted bilinear map. Then λ is the twisted bilinear map associated to some group G , some short exact sequence*

$$1 \longrightarrow K \longrightarrow G \xrightarrow{\rho} B \rtimes A \longrightarrow 1,$$

and some homomorphisms $\iota_1: A \rightarrow G$ and $\iota_2: B \rightarrow G$.

Proof. Set $Q = B \rtimes A$. We will construct G using the theory of nonabelian group extensions sketched in [3, §IV.6] and proven in detail in [7]. This machine needs two inputs.

- The first is a set map $\phi: Q \rightarrow \text{Aut}(K)$ satisfying $\phi(1) = \text{id}$. For $q \in Q$, we define $\phi(q) \in \text{Aut}(K)$ as follows. We can uniquely write $q = ba$ with $b \in B$ and $a \in A$. We then define

$$\phi(q)(k) = \beta_b(\alpha_a(k)) \quad (k \in K).$$

- The second is a set map $\gamma: Q \times Q \rightarrow K$ satisfying $\gamma(1, q) = \gamma(q, 1) = 1$ for all $q \in Q$, which we define as follows. Consider $q_1, q_2 \in Q$. We can uniquely write $q_1 = b_1 a_1$ and $q_2 = b_2 a_2$ with $b_1, b_2 \in B$ and $a_1, a_2 \in A$. We then define

$$\gamma(q_1, q_2) = \beta_{b_1}(\lambda(a_1, b_2)).$$

We remark that these pieces of data are not homomorphisms. They must satisfy two key identities which we will verify below in Claims 1 and 2. We postpone these verifications momentarily to explain the output of the machine.

Let G be the set of pairs (k, q) with $K \in K$ and $q \in Q$. Define a multiplication in G via the formula

$$(k_1, q_1)(k_2, q_2) = (k_1 \cdot \phi(q_1)(k_2) \cdot \gamma(q_1, q_2), q_1 q_2).$$

The machine says that this G is a group (the purpose of the postponed identities is to show that the above multiplication is associative). It clearly lies in a short exact sequence

$$1 \longrightarrow K \longrightarrow G \xrightarrow{\rho} B \rtimes A \longrightarrow 1,$$

and we can define the desired homomorphisms $\iota_1: A \rightarrow G$ and $\iota_2: B \rightarrow G$ via the formulas $\iota_1(a) = (1, (1, a))$ and $\iota_2(b) = (1, (b, 1))$. An easy calculation shows that the conclusions of the theorem are satisfied.

It remains to verify the two needed identities, which are as follows.

Claim 1. *For $q_1, q_2 \in Q$ and $k \in K$, we have*

$$\phi(q_1)(\phi(q_2)(k)) = \gamma(q_1, q_2) \cdot \phi(q_1 q_2)(k) \cdot \gamma(q_1, q_2)^{-1}.$$

For $i = 1, 2$, write $q_i = b_i a_i$ with $b_i \in B$ and $a_i \in A$, so $q_1 q_2 = (b_1 a_1 b_2)(a_1 a_2)$. We then have

$$\begin{aligned} \gamma(q_1, q_2) \cdot \phi(q_1 q_2)(k) \cdot \gamma(q_1, q_2)^{-1} &= \beta_{b_1}(\lambda(a_1, b_2)) \cdot \beta_{b_1 a_1 b_2}(\alpha_{a_1 a_2}(k)) \cdot \beta_{b_1}(\lambda(a_1, b_2))^{-1} \\ &= \beta_{b_1}(\lambda(a_1, b_2) \cdot \beta_{a_1 b_2}(\alpha_{a_1}(\alpha_{a_2}(k)))) \cdot \lambda(a_1, b_2)^{-1}. \end{aligned}$$

Property TB3 of a twisted bilinear map says that this equals

$$\beta_{b_1}(\alpha_{a_1}(\beta_{b_2}(\alpha_{a_2}(k)))) = \phi(q_1)(\phi(q_2)(k)),$$

as claimed.

Claim 2. For $q_1, q_2, q_3 \in Q$, we have $\gamma(q_1, q_2) \cdot \gamma(q_1 q_2, q_3) = \phi(q_1)(\gamma(q_2, q_3)) \cdot \gamma(q_1, q_2 q_3)$.

For $i = 1, 2, 3$, write $q_i = b_i a_i$ with $b_i \in B$ and $a_i \in A$. We begin by examining the left side of the purported equality. Since $q_1 q_2 = (b_1 a_1 b_2)(a_1 a_2)$, it equals

$$\begin{aligned} \gamma(q_1, q_2) \cdot \gamma(q_1 q_2, q_3) &= \beta_{b_1}(\lambda(a_1, b_2)) \cdot \beta_{b_1 a_1 b_2}(\lambda(a_1 a_2, b_3)) \\ &= \beta_{b_1}(\lambda(a_1, b_2) \cdot \beta_{a_1 b_2}(\lambda(a_1 a_2, b_3))). \end{aligned}$$

Using property TB2 of a twisted bilinear map, this equals

$$\beta_{b_1}(\lambda(a_1, b_2) \cdot \beta_{a_1 b_2}(\alpha_{a_1}(\lambda(a_2, b_3)) \cdot \lambda(a_1, a_2 b_3))) \quad (8)$$

Property TB3 of a twisted bilinear map with $a = a_1$ and $b = b_2$ and $k = \lambda(a_2, b_3)$ says that

$$\lambda(a_1, b_2) \cdot \beta_{a_1 b_2}(\alpha_{a_1}(\lambda(a_2, b_3))) = \alpha_{a_1}(\beta_{b_2}(\lambda(a_2, b_3))) \cdot \lambda(a_1, b_2).$$

Plugging this into (8) (and remembering to distribute the $\beta_{a_1 b_2}$ over the last term), we get

$$\beta_{b_1}(\alpha_{a_1}(\beta_{b_2}(\lambda(a_2, b_3))) \cdot \lambda(a_1, b_2) \cdot \beta_{a_1 b_2}(\lambda(a_1, a_2 b_3))). \quad (9)$$

We now turn to the right hand side of the purported equality. Since $q_2 q_3 = (b_2 a_2 b_3)(a_2 a_3)$, it equals

$$\beta_{b_1}(\alpha_{a_1}(\beta_{b_2}(\lambda(a_2, b_3)))) \cdot \beta_{b_1}(\lambda(a_1, b_2 a_2 b_3)) = \beta_{b_1}(\alpha_{a_1}(\beta_{b_2}(\lambda(a_2, b_3))) \cdot \lambda(a_1, b_2 a_2 b_3)).$$

Property TB1 of a twisted bilinear map says that this equals

$$\beta_{b_1}(\alpha_{a_1}(\beta_{b_2}(\lambda(a_2, b_3))) \cdot \lambda(a_1, b_2) \cdot \beta_{a_1 b_2}(\lambda(a_1, a_2 b_3))). \quad (10)$$

Since (9) and (10) are equal, the claim follows. \square

5.3 Presentations of $\text{Aut}(F_n)$ and $\mathcal{A}_{n,1}$

This is the third and final section of preliminaries for the proof of Theorem D. In it, we give presentations for the groups $\text{Aut}(F_n)$ and $\mathcal{A}_{n,1}$.

We begin with $\text{Aut}(F_n)$. Recall that $X = \{x_1, \dots, x_n\}$ is the standard basis for F_n . The presentation for $\text{Aut}(F_n)$ we will use has three classes of generators.

- For $\alpha \in \{1, -1\}$ and distinct $x_a, x_b \in X$, we need the automorphisms $M_{x_a^\alpha, x_b}$ defined in the introduction. Recall that their characteristic properties are that

$$M_{x_a^\alpha, x_b}(x_a^\alpha) = x_b x_a^\alpha \quad \text{and} \quad M_{x_a^\alpha, x_b}(x_c) = x_c$$

for $x_c \in X$ with $x_c \neq x_a$. The elements $M_{x_a^\alpha, x_b}$ will be called *transvections*.

- For distinct $x_a, x_b \in X$, we will need the automorphisms $P_{a,b}$ defined via the formula

$$P_{a,b}(x_c) = \begin{cases} x_b & \text{if } c = a, \\ x_a & \text{if } c = b, \\ x_c & \text{otherwise} \end{cases} \quad (1 \leq c \leq n).$$

The elements $P_{a,b}$ will be called *swaps*.

-
- N1. relations for the subgroup generated by inversions and swaps, a signed permutation group:
- $I_a^2 = 1$ and $[I_a, I_b] = 1$,
 - $P_{a,b}^2 = 1$, $[P_{a,b}, P_{c,d}] = 1$, and $P_{a,b}P_{b,c}P_{a,b}^{-1} = P_{a,c}$,
 - $P_{a,b}I_aP_{a,b}^{-1} = I_b$ and $[P_{a,b}, I_c] = 1$;
- N2. relations for conjugating transvections by inversions and swaps, coming from the natural action of inversions and swaps on $\{x_1, \dots, x_n\}$:
- $P_{a,b}M_{x_c^\gamma, x_d}P_{a,b}^{-1} = M_{P_{a,b}(x_c^\gamma), P_{a,b}(x_d)}$ even if $\{a, b\} \cap \{c, d\} \neq \emptyset$,
 - $I_aM_{x_c^\gamma, x_d}I_a^{-1} = M_{I_a(x_c^\gamma), I_a(x_d)}$ even if $a \in \{c, d\}$;
- N3. $M_{x_a^\alpha, x_b}^\beta M_{x_b^\beta, x_a}^\alpha M_{x_a^\alpha, x_b}^{-\beta} = I_bP_{a,b}$;
- N4. $[M_{x_a^\alpha, x_b}^\alpha, M_{x_c^\gamma, x_d}^\gamma] = 1$ with a, b, c, d not necessarily all distinct, such that $a \neq b$, $c \neq d$, $x_a^\alpha \notin \{x_c^\gamma, x_d, x_d^{-1}\}$ and $x_c^\gamma \notin \{x_b, x_b^{-1}\}$;
- N5. $M_{x_b^\beta, x_a}^\alpha M_{x_c^\gamma, x_b}^\beta = M_{x_c^\gamma, x_b}^\beta M_{x_b^\beta, x_a}^\alpha M_{x_c^\gamma, x_a}^\alpha$.
-

Table 2: Nielsen’s relations for $\text{Aut}(F_n)$ consist of the set R_A of relations listed above. The letters a, b, c, d are elements of $\{1, \dots, n\}$ (assumed distinct unless otherwise stated) and $\alpha, \beta, \gamma, \in \{1, -1\}$.

- For $x_a \in X$, we will need the automorphisms I_a defined via the formula

$$I_a(x_b) = \begin{cases} x_b^{-1} & \text{if } b = a, \\ x_b & \text{otherwise} \end{cases} \quad (1 \leq b \leq n).$$

The elements I_a will be called *inversions*.

Let S_A be the set consisting of the above generators. The set S_A does not contain elements of the form $M_{x_a^\alpha, x_b^{-1}}$, but we will frequently use $M_{x_a^\alpha, x_b^{-1}}$ as an alternate notation for $M_{x_a^\alpha, x_b}^{-1}$. We then have the following theorem of Nielsen.

Theorem 5.6 (Nielsen [13]). *The group $\text{Aut}(F_n)$ has the presentation $\langle S_A \mid R_A \rangle$, where R_A is given in Table 2.*

We now turn to $\mathcal{A}_{n,1}$. Recall that this is a subgroup of $\text{Aut}(F_{n,1})$, where $F_{n,1}$ is the free group on $\{x_1, \dots, x_n, y\}$. We will use a presentation that is due to Jensen–Wahl [10]. See [5, Theorem 5.2] for a small correction to Jensen–Wahl’s original statement.

Theorem 5.7 (Jensen–Wahl [10]). *The group $\mathcal{A}_{n,1}$ has the presentation whose generators are the union of S_A with the set*

$$\{M_{x_a^\alpha, y}, C_{y, x_a} \mid x_a \in X, \alpha \in \{1, -1\}\}$$

and whose relations are those appearing in Table 3.

-
- Q1. Nielsen's relations among S_A from Table 2,
Q2. Commuting relations:
– $[M_{x_a^\alpha, y}, M_{x_b^\beta, y}] = 1$ if $x_a^\alpha \neq x_b^\beta$,
– $[M_{x_a^\alpha, x_b}, M_{x_c^\gamma, y}] = 1$ if $x_a^\alpha \neq x_c^\gamma$,
– $[M_{x_a^\alpha, x_b}, C_{y, x_c}] = 1$ if $c \neq a$.
- Q3. The obvious analogues of the N2 relations from Table 2 giving the effect of conjugating C_{y, x_a} and $M_{x_a^\alpha, y}$ by swaps and inversions,
Q4. $M_{x_a^\alpha, x_b}^{-\beta} M_{x_b^\beta, y} M_{x_a^\alpha, x_b}^\beta = M_{x_a^\alpha, y} M_{x_b^\beta, y}$, and
Q5. $C_{y, x_a}^{-\alpha} M_{x_a^{-\alpha}, y} C_{y, x_a}^\alpha = M_{x_a^\alpha, y}^{-1}$.
-

Table 3: *Jensen–Wahl's relations for $\mathcal{A}_{n,1}$ consist of the relations above. The letters a, b, c are elements of $\{1, \dots, n\}$ (assumed distinct unless otherwise stated) and $\alpha, \beta, \gamma, \in \{1, -1\}$.*

6 A finite L-presentation for $\mathcal{K}_{n,1}^{\text{IA}}$

This section contains the proof of Theorem D, which asserts that $\mathcal{K}_{n,1}^{\text{IA}}$ has a finite L-presentation. We begin in §6.1 by describing the generators, relations, and endomorphisms which make up our L-presentation. Next, in §6.2 we construct the data needed to use the theory of twisted bilinear maps to construct an appropriate extension of our purported presentation for $\mathcal{K}_{n,1}^{\text{IA}}$. Finally, in §6.3 we prove that our presentation is complete.

The proofs of several of our lemmas will depend on computer calculations. These computer calculations will be discussed in §7.

Just like in §5, we will denote the free basis for $F_{n,1}$ by $\{x_1, \dots, x_n, y\}$.

6.1 Statement of L-presentation

In this section, we will describe the generators, relations, and endomorphisms that make up the finite L-presentation for $\mathcal{K}_{n,1}^{\text{IA}}$ whose existence is asserted by Theorem D. To help us keep track of the role that our symbols are playing, we will change the font for the generators of $\mathcal{K}_{n,1}^{\text{IA}}$ and use

$$S_K = \{\mathfrak{C}_{y, x_a}, \mathfrak{C}_{x_a, y} \mid x_a \in X\} \cup \{\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mid x_a, x_b \in X, x_a \neq x_b, \alpha, \beta, \epsilon \in \{1, -1\}\}$$

as our generating set. We remark that elements of the form $\mathfrak{M}_{x_a^\alpha, [x_b^\beta, y^\epsilon]}$ are not included in S_K ; however, we will frequently use the symbol $\mathfrak{M}_{x_a^\alpha, [x_b^\beta, y^\epsilon]}$ as a synonym for $\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]}^{-1}$, which is the inverse of an element of S_K .

Next, we explain our endomorphisms E_K for our L-presentation. These endomorphisms are indexed by the generators for $\mathcal{A}_{n,1}$ given by Theorem 5.7 that do not lie in $\mathcal{K}_{n,1}^{\text{IA}}$. More

precisely, define

$$S_Q = \{P_{a,b}, I_a, M_{x_a,y}, M_{x_a^\alpha,x_b} \mid x_a, x_b \in X, x_a \neq x_b, \alpha \in \{1, -1\}\}.$$

We use $S_Q^{\pm 1}$ to denote $S_Q \cup \{s^{-1} \mid s \in S_Q\}$ (and similarly for other sets). We remark that in S_Q , we regard $P_{a,b}$ and $P_{b,a}$ as being the same element.

We now define a function $\phi: S_Q^{\pm 1} \rightarrow \text{End}(F(S_K))$ (see Lemma 6.1 below for an elucidation of the purpose of this definition). By the universal property of a free group, to do this it is enough to give $\phi(s)(t)$ for each choice of $s \in S_Q^{\pm 1}$ and $t \in S_K$. Once we have given these formulas, we define E_K to be the image of the map ϕ . In our defining formulas, we use x_a, x_b, x_c, \dots for elements of X and $\alpha, \beta, \gamma, \epsilon, \dots$ for elements of $\{1, -1\}$. Elements with distinct subscripts are assumed to be distinct unless noted.

The action of swaps and inversions through ϕ is by acting on the elements of X indexing our generators: if s is a swap or inversion, then

$$\phi(s)(\mathfrak{C}_{x_a,y}) = \mathfrak{C}_{s(x_a),y}, \quad \phi(s)(\mathfrak{C}_{y,x_a}) = \mathfrak{C}_{y,s(x_a)}, \quad \text{and} \quad \phi(s)(\mathfrak{M}_{x_a^\alpha,[y^\epsilon,x_b^\beta]}) = \mathfrak{M}_{s(x_a^\alpha),[y^\epsilon,s(x_b^\beta)]}.$$

Here we interpret

$$\mathfrak{C}_{x_a^{-1},y} = \mathfrak{C}_{x_a,y} \quad \text{and} \quad \mathfrak{C}_{y,x_a^{-1}} = \mathfrak{C}_{y,x_a}^{-1}.$$

Furthermore, for s a swap or inversion we define $\phi(s^{-1})(t) = \phi(s)(t)$ for any $t \in S_K$.

If $s \in S_Q$ is of the form $M_{x_a^\alpha,y}$, we define $\phi(s)(t) = \phi(s^{-1})(t) = t$ if $t \in S_K$ is anything of the form

$$\mathfrak{C}_{x_a,y}, \quad \mathfrak{C}_{x_b,y}, \quad \mathfrak{M}_{x_a^{-\alpha},[y^\epsilon,x_b^\beta]}, \quad \text{or} \quad \mathfrak{M}_{x_b^\beta,[y^\epsilon,x_c^\gamma]}.$$

If $s \in S_Q$ is of the form $M_{x_a^\alpha,x_b}$, we define $\phi(s)(t) = \phi(s^{-1})(t) = t$ if $t \in S_K$ is anything of the form

$$\mathfrak{C}_{x_c,y}, \quad \mathfrak{C}_{y,x_b}, \quad \mathfrak{C}_{y,x_c}, \quad \mathfrak{M}_{x_a^{-\alpha},[y^\epsilon,x_b^\beta]}, \quad \mathfrak{M}_{x_a^{-\alpha},[y^\epsilon,x_c^\gamma]}, \quad \mathfrak{M}_{x_c^\gamma,[y^\epsilon,x_b^\beta]}, \quad \text{or} \quad \mathfrak{M}_{x_c^\gamma,[y^\epsilon,x_d^\delta]}.$$

The other cases for $\phi(s)(t) \in F(S_K)$, for $s \in S_Q^{\pm 1}$ and $t \in S_K$, are given in Table 4.

The key property of the map ϕ is as follows.

Lemma 6.1. *Let $\Psi: F(S_K) \rightarrow \mathcal{K}_{n,1}^{IA}$ be the natural surjection. Then regarding S_Q as a subset of $\mathcal{A}_{n,1}$, we have*

$$\Psi(\phi(s)(t)) = s\Psi(t)s^{-1} \quad (s \in S_Q, t \in S_K).$$

Proof. This is a computer calculation which is described in Lemma 7.3 below. \square

We can now give a statement of our L-presentation. The following theorem (which will be proven in §6.3) is a more precise version of Theorem D.

Theorem 6.2. *Let S_K and S_Q and ϕ be as above and let R_K^0 be the set of relations in Table 5. Then the group $\mathcal{K}_{n,1}^{IA}$ has the finite L-presentation $\mathcal{K}_{n,1}^{IA} = \langle S_K \mid R_K^0 \mid \phi(S_Q^{\pm 1}) \rangle$.*

$s \in S_Q^{\pm 1}$	$t \in S_K$	$\phi(s)(t)$
$M_{x_a, y}^\epsilon$	\mathfrak{C}_{y, x_b}	$\mathfrak{C}_{y, x_b} \mathfrak{M}_{x_a, [y^{-\epsilon}, x_b^{-1}]}$
	\mathfrak{C}_{y, x_a}	$\mathfrak{C}_{x_a, y}^\epsilon \mathfrak{C}_{y, x_a}$
	$\mathfrak{M}_{x_a, [y^\zeta, x_b^\beta]}$	$\mathfrak{C}_{x_a, y}^\epsilon \mathfrak{M}_{x_a, [y^\zeta, x_b^\beta]} \mathfrak{C}_{x_a, y}^{-\epsilon}$
	$\mathfrak{M}_{x_b^\beta, [y^\zeta, x_a]}$	$\mathfrak{C}_{x_a, y}^\epsilon \mathfrak{M}_{x_b^\beta, [y^\zeta, x_a]} \mathfrak{C}_{x_a, y}^{-\epsilon}$
$M_{x_a^\alpha, x_b}^\beta$	\mathfrak{C}_{y, x_a}	$(\mathfrak{C}_{y, x_a}^\alpha \mathfrak{C}_{y, x_b}^\beta)^\alpha$
	$\mathfrak{C}_{x_a, y}$	$\mathfrak{C}_{x_a, y} \mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y]}$
	$\mathfrak{C}_{x_b, y}$	$\mathfrak{C}_{x_b, y} \mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y^{-1}]}$
	$\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_c^\zeta]}$	$\mathfrak{C}_{y, x_c}^\gamma \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^{-\beta}]} \mathfrak{C}_{y, x_c}^{-\gamma} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_c^\zeta]} \mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y^\epsilon]}$
	$\mathfrak{M}_{x_b^\beta, [y^\epsilon, x_c^\zeta]}$	$\mathfrak{M}_{x_b^\beta, [y^\epsilon, x_c^\zeta]} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^{-\beta}]} \mathfrak{M}_{x_a^\alpha, [x_c^\zeta, y^\epsilon]} \mathfrak{C}_{y^\epsilon, x_c^\zeta} \mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y^\epsilon]} \mathfrak{C}_{y^\epsilon, x_c}^{-\gamma}$
	$\mathfrak{M}_{x_b^{-\beta}, [y^\epsilon, x_c^\zeta]}$	$\mathfrak{M}_{x_b^{-\beta}, [y^\epsilon, x_c^\zeta]} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_c^\zeta]}$
	$\mathfrak{M}_{x_c^\zeta, [y^\epsilon, x_a^\alpha]}$	$\mathfrak{C}_{y, x_a}^\alpha \mathfrak{M}_{x_c^\zeta, [y^\epsilon, x_b^\beta]} \mathfrak{C}_{y, x_a}^{-\alpha} \mathfrak{M}_{x_c^\zeta, [y^\epsilon, x_a^\alpha]}$
	$\mathfrak{M}_{x_c^\zeta, [y^\epsilon, x_a^{-\alpha}]}$	$\mathfrak{C}_{y, x_b}^{-\beta} \mathfrak{M}_{x_c^\zeta, [y^\epsilon, x_a^{-\alpha}]} \mathfrak{M}_{x_c^\zeta, [x_b^\beta, y^\epsilon]} \mathfrak{C}_{y, x_b}^\beta$
	$\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]}$	$\mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y^\epsilon]}$
	$\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^{-\beta}]}$	$\mathfrak{C}_{y, x_b}^{-\beta} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^{-\beta}]} \mathfrak{C}_{y, x_b}^\beta$
	$\mathfrak{M}_{x_b^\beta, [y^\epsilon, x_a^\alpha]}$	$\mathfrak{M}_{x_b^{-\beta}, [y^{-\epsilon}, x_a^\alpha]} \mathfrak{C}_{x_b, y}^\epsilon \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^{-\beta}]} \mathfrak{C}_{x_a, y}^{-\epsilon}$
	$\mathfrak{M}_{x_b^{-\beta}, [y^\epsilon, x_a^\alpha]}$	$\mathfrak{C}_{y, x_a}^\alpha \mathfrak{C}_{y, x_b}^\beta \mathfrak{C}_{x_b, y}^{-\epsilon} \mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y^\epsilon]} \mathfrak{C}_{y, x_b}^{-\beta} \mathfrak{M}_{x_b^{-\beta}, [x_a^{-\alpha}, y^\epsilon]} \mathfrak{C}_{y, x_a}^{-\alpha} \mathfrak{C}_{x_a, y}^\epsilon$
	$\mathfrak{M}_{x_b^\beta, [y^\epsilon, x_a^{-\alpha}]}$	$\mathfrak{C}_{y, x_b}^{-\beta} \mathfrak{C}_{y, x_a}^{-\alpha} \mathfrak{C}_{x_a, y}^\epsilon \mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y^\epsilon]} \mathfrak{C}_{x_b, y}^{-\beta} \mathfrak{M}_{x_b^{-\beta}, [x_a^\alpha, y^{-\epsilon}]} \mathfrak{C}_{y, x_a}^\alpha \mathfrak{C}_{y, x_b}^\beta$
	$\mathfrak{M}_{x_b^{-\beta}, [y^\epsilon, x_a^{-\alpha}]}$	$\mathfrak{C}_{y, x_b}^{-\beta} \mathfrak{C}_{y, x_a}^{-\alpha} \mathfrak{C}_{x_a, y}^{-\epsilon} \mathfrak{M}_{y, x_a}^\alpha \mathfrak{M}_{x_b^{-\beta}, [y^\epsilon, x_a^{-\alpha}]} \mathfrak{C}_{y, x_b}^\beta \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^{-\beta}]} \mathfrak{C}_{x_b, y}^\epsilon$

Table 4: The defining formulas for ϕ . The letters a, b, c are distinct elements of $\{1, \dots, n\}$ $\alpha, \beta, \gamma, \zeta \in \{1, -1\}$.

-
- R1. $[\mathfrak{C}_{x_a, y}, \mathfrak{C}_{x_b, y}] = 1$;
- R2. $[\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_c^\zeta]}, \mathfrak{M}_{x_b^\beta, [y, x_d^\delta]}] = 1$, possibly with $x_a^\alpha = x_b^{-\beta}$ or $x_c = x_d$ (or both), as long as $x_a^\alpha \neq x_b^\beta$, $x_a \neq x_d$ and $x_b \neq x_c$;
- R3. $[\mathfrak{C}_{x_a, y}, \mathfrak{M}_{x_b^\beta, [y^\epsilon, x_c^\zeta]}] = 1$;
- R4. $\mathfrak{C}_{y, x_b}^{-\beta} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{C}_{y, x_b}^\beta = \mathfrak{M}_{x_a^\alpha, [x_b^{-\beta}, y^\epsilon]}$;
- R5. $\mathfrak{C}_{x_b, y}^{-\epsilon} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{C}_{x_b, y}^\epsilon = \mathfrak{M}_{x_a^\alpha, [x_b^\beta, y^{-\epsilon}]}$;
- R6. $\mathfrak{C}_{x_a, y}^\epsilon \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{C}_{x_a, y}^{-\epsilon} = \mathfrak{M}_{x_a^\alpha, [x_b^\beta, y^{-\epsilon}]}$;
- R7. $\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{M}_{x_a^{-\alpha}, [y^\epsilon, x_b^\beta]} = \mathfrak{C}_{y, x_b}^\beta \mathfrak{C}_{x_a, y}^{-\epsilon} \mathfrak{C}_{y, x_b}^{-\beta} \mathfrak{C}_{x_a, y}^\epsilon$;
- R8. $\mathfrak{M}_{x_b^\beta, [y^{-\epsilon}, x_c^\zeta]} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{M}_{x_b^\beta, [x_c^\zeta, y^{-\epsilon}]} = \mathfrak{M}_{x_a^\alpha, [x_c^\zeta, y^{-\epsilon}]} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{M}_{x_a^\alpha, [x_c^\zeta, y^\epsilon]}$;
- R9. $\mathfrak{C}_{x_b, y}^{-\epsilon} \mathfrak{C}_{y, x_c}^\gamma \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{C}_{y, x_c}^{-\gamma} \mathfrak{C}_{x_b, y}^\epsilon$
 $= \mathfrak{M}_{x_a^\alpha, [x_b^\beta, y^{-\epsilon}]} \mathfrak{C}_{y, x_c}^\gamma \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{C}_{y, x_c}^{-\gamma} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_c^\zeta]} \mathfrak{M}_{x_a^\alpha, [x_b^\beta, y^\epsilon]} \mathfrak{M}_{x_a^\alpha, [x_c^\zeta, y^\epsilon]}$;
- R10. $\mathfrak{C}_{x_c, y}^{-\epsilon} \mathfrak{C}_{y, x_c}^\gamma \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{C}_{y, x_c}^{-\gamma} \mathfrak{C}_{x_c, y}^\epsilon$
 $= \mathfrak{M}_{x_a^\alpha, [y^{-\epsilon}, x_b^\beta]} \mathfrak{M}_{x_a^\alpha, [x_c^\zeta, y^{-\epsilon}]} \mathfrak{C}_{y, x_c}^\gamma \mathfrak{M}_{x_a^\alpha, [x_b^\beta, y^{-\epsilon}]} \mathfrak{C}_{y, x_c}^{-\gamma} \mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mathfrak{M}_{x_a^\alpha, [y^{-\epsilon}, x_c^\zeta]}$.
-

Table 5: The relations R_K^0 for the L -presentation of $\mathcal{K}_{n,1}^{IA}$. The letters a, b, c, d are elements of $\{1, \dots, n\}$ (assumed distinct unless otherwise stated) and $\alpha, \beta, \gamma, \delta, \epsilon \in \{1, -1\}$.

6.2 Constructing the extension

Let $\Gamma_n = \langle S_K \mid R_K^0 \mid \phi(S_Q^{\pm 1}) \rangle$ be the group with the presentation described in Theorem 6.2. There is thus a surjection $\Psi: \Gamma_n \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$ which Theorem 6.2 claims is an isomorphism. Lemma 5.3 says that there exists a short exact sequence

$$1 \longrightarrow \mathcal{K}_{n,1}^{\text{IA}} \longrightarrow \mathcal{A}_{n,1} \xrightarrow{\rho} \mathbb{Z}^n \rtimes_r \text{Aut}(F_n) \longrightarrow 1 \quad (11)$$

together with homomorphisms $\iota_1: \text{Aut}(F_n) \rightarrow \mathcal{A}_{n,1}$ and $\iota_2: \mathbb{Z}^n \rightarrow \mathcal{A}_{n,1}$ such that $\rho \circ \iota_1 = \text{id}$ and $\rho \circ \iota_2 = \text{id}$. The purpose of this section is to construct the data needed to apply Theorem 5.5 and deduce that there exists a similar extension involving Γ_n instead of $\mathcal{K}_{n,1}^{\text{IA}}$.

For $f \in \text{Aut}(F_n)$ and $z \in \mathbb{Z}^n$, define homomorphisms $\bar{\alpha}_f: \mathcal{K}_{n,1}^{\text{IA}} \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$ and $\bar{\beta}_z: \mathcal{K}_{n,1}^{\text{IA}} \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$ via the formulas

$$\bar{\alpha}_f(x) = \iota_1(f)x\iota_1(f)^{-1} \quad \text{and} \quad \bar{\beta}_z(x) = \iota_2(z)x\iota_2(z)^{-1} \quad (x \in \mathcal{K}_{n,1}^{\text{IA}}).$$

These define actions of $\text{Aut}(F_n)$ and \mathbb{Z}^n on $\mathcal{K}_{n,1}^{\text{IA}}$. Using the construction described in Example 5.2, we obtain from (11) a twisted bilinear map $\bar{\lambda}: \text{Aut}(F_n) \times \mathbb{Z}^n \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$. We must lift all of this data to Γ_n . This is accomplished in the following three lemmas. For a set S , let S^* denote the free monoid on S , so S^* consists of words in S .

Lemma 6.3. *There exists an action of $\text{Aut}(F_n)$ on Γ_n with the following property. For $f \in \text{Aut}(F_n)$, denote by $\alpha_f: \Gamma_n \rightarrow \Gamma_n$ the associated automorphism. Then*

$$\Psi(\alpha_f(x)) = \bar{\alpha}_f(\Psi(x)) \quad (f \in \text{Aut}(F_n), x \in \Gamma_n).$$

Proof. Let $\text{Aut}(F_n) = \langle S_A \mid R_A \rangle$ be the presentation given by Theorem 5.6. We have $S_A \subset S_Q$, so the map $\phi: S_Q^{\pm 1} \rightarrow \text{End}(F(S_K))$ used in the construction of the L-presentation for Γ_n restricts to a set map $S_A^{\pm 1} \rightarrow \text{End}(F(S_K))$. By the definition of an L-presentation, the image of this set map preserves the relations between elements of S_K that make up Γ_n , so we get a set map $S_A^{\pm 1} \rightarrow \text{End}(\Gamma_n)$. By the universal property of the free monoid, this induces a monoid homomorphism $\zeta: (S_A^{\pm 1})^* \rightarrow \text{End}(\Gamma_n)$. Computer calculations described in Lemma 7.4 below show that $\zeta(s)(\zeta(s^{-1})(t)) = t$ for all $s \in S_A^{\pm 1}$ and $t \in S_K$, which implies that the image of ζ is contained in $\text{Aut}(\Gamma_n)$ and that ζ descends to a group homomorphism $\eta: F(S_A) \rightarrow \text{Aut}(\Gamma_n)$. Further computer calculations described in Lemma 7.5 below show that $\eta(r)(t) = t$ for all $r \in R_A$ and all $t \in S_K$. This implies that η descends to a group homomorphism $\text{Aut}(F_n) \rightarrow \text{Aut}(\Gamma_n)$. This is the desired action; the claimed naturality property follows from Lemma 6.1. \square

Lemma 6.4. *There exists an action of \mathbb{Z}^n on Γ_n with the following property. For $z \in \mathbb{Z}^n$, denote by $\beta_z: \Gamma_n \rightarrow \Gamma_n$ the associated automorphism. Then*

$$\Psi(\beta_z(x)) = \bar{\beta}_z(\Psi(x)) \quad (z \in \mathbb{Z}^n, x \in \Gamma_n).$$

Proof. Set $S_Z = \{M_{x_1,y}, \dots, M_{x_n,y}\}$ and $R_Z = \{[M_{x_i,y}, M_{x_j,y}] \mid 1 \leq i < j \leq n\}$, so $S_Z \subset S_Q$ and $\mathbb{Z}^n = \langle S_Z \mid R_Z \rangle$. Just like in the proof of Lemma 6.3, the map $\phi: S_Q^{\pm 1} \rightarrow$

$f \in S_A^{\pm 1}$	$z \in S_Z^{\pm 1}$	$\lambda(f, z) \in F(S_K)$
$I_a^{\pm 1}$	$M_{x_a, y}^\epsilon$	$\mathfrak{C}_{x_a, y}^\epsilon$
M_{x_a, x_b}^β	$M_{x_a, y}^\epsilon$	$\mathfrak{M}_{x_a, [y^{-\epsilon}, x_b^{-\beta}]}$
M_{x_a, x_b}	$M_{x_b, y}^\epsilon$	$\mathfrak{M}_{x_a, [y^\epsilon, x_b^{-1}]}$
$M_{x_a^{-1}, x_b}$	$M_{x_b, y}^\epsilon$	$(\mathfrak{M}_{x_a^{-1}, [y, x_b^{-1}]} \mathfrak{C}_{x_a, y}^{-1})^\epsilon$
$M_{x_a^{-1}, x_b}^{-1}$	$M_{x_b, y}^\epsilon$	$\mathfrak{C}_{x_a, y}^\epsilon$

Table 6: The effect of $\lambda(\cdot, \cdot)$ on generators. For $f \in S_A^{\pm 1}$ and $z \in S_Z^{\pm 1}$ such that there is no entry in the above table, we have $\lambda(f, z) = 1$.

$\text{End}(F(S_K))$ restricts to a set map $S_Z^{\pm 1} \rightarrow \text{End}(F(S_K))$ which induces a monoid homomorphism $\zeta: (S_Z^{\pm 1})^* \rightarrow \text{End}(\Gamma_n)$. Computer calculations described in Lemma 7.6 below show that $\zeta(s)(\zeta(s^{-1})(t)) = t$ for all $s \in S_Z^{\pm 1}$ and $t \in S_K$, so ζ induces a group homomorphism $\eta: F(S_Z) \rightarrow \text{Aut}(\Gamma_n)$. Further computer calculations described in Lemma 7.7 below show that $\eta(r)(t) = t$ for all $r \in R_Z$ and all $t \in S_K$. This implies that η descends to a group homomorphism $\mathbb{Z}^n \rightarrow \text{Aut}(\Gamma_n)$. This is the desired action; the claimed naturality property follows from Lemma 6.1. \square

Lemma 6.5. *With respect to the action of $\text{Aut}(F_n)$ on \mathbb{Z}^n introduced in §5.1 and the actions of \mathbb{Z}^n and $\text{Aut}(F_n)$ on Γ_n given by Lemmas 6.3 and 6.4, there exists a twisted bilinear map $\lambda: \text{Aut}(F_n) \times \mathbb{Z}^n \rightarrow \Gamma_n$ such that*

$$\Psi(\lambda(f, z)) = \bar{\lambda}(f, z) \quad (f \in \text{Aut}(F_n), z \in \mathbb{Z}^n).$$

Proof. Let α_f and β_z be as in Lemmas 6.3 and 6.4, respectively. Let $\text{Aut}(F_n) = \langle S_A \mid R_A \rangle$ be the presentation given by Theorem 5.6. Also, let $S_Z = \{M_{x_1, y}, \dots, M_{x_n, y}\}$ and $R_Z = \{[M_{x_i, y}, M_{x_j, y}] \mid 1 \leq i < j \leq n\}$, so $\mathbb{Z}^n = \langle S_Z \mid R_Z \rangle$. We claim that it is enough to construct a twisted bilinear map $\lambda: \text{Aut}(F_n) \times \mathbb{Z}^n \rightarrow \Gamma_n$ such that

$$\Psi(\lambda(f, z)) = \bar{\lambda}(f, z) \quad (f \in S_A^{\pm 1}, z \in S_Z^{\pm 1}). \quad (12)$$

Indeed, the axioms of a twisted bilinear map show that λ is determined by its values on generators: property TB2 says that $\lambda(a_1 a_2, b) = \alpha_{a_1}(\lambda(a_2, b)) \cdot \lambda(a_1, {}^{a_2}b)$ for all $a_1, a_2 \in \text{Aut}(F_n)$ and $b \in \mathbb{Z}^n$, so the values of λ are determined by the values of $\lambda(f, z)$ for $f \in S_A^{\pm 1}$ and $z \in \mathbb{Z}^n$, and then property TB1 says that $\lambda(a, b_1 b_2) = \lambda(a, b_1) \cdot \beta_{a b_1}(\lambda(a, b_2))$ for all $a \in \text{Aut}(F_n)$ and $b_1, b_2 \in \mathbb{Z}^n$, so the values of λ are determined by the values of $\lambda(f, z)$ for $f \in S_A^{\pm 1}$ and $z \in S_Z^{\pm 1}$. An analogous fact holds for $\bar{\lambda}$, whence the claim.

We will construct λ such that $\lambda(f, z)$ is as in Table 6 for $f \in S_A^{\pm 1}$ and $z \in S_Z^{\pm 1}$. It is easy to check that these values satisfy (12). We will do this in four steps. For a set S , let S^* be the free monoid on S , so S^* consists of words in S .

- First, for $f \in S_A^{\pm 1}$ we will use the “expansion rule” TB1 to construct a map $\tilde{\lambda}_1(f, \cdot)$ from $(S_Z^{\pm 1})^*$ to Γ_n with $\tilde{\lambda}_1(f, z)$ equal to the value of $\lambda(f, z)$ from Table 6 for $z \in S_Z^{\pm 1}$.
- Next, we will show that $\tilde{\lambda}_1(f, \cdot)$ descends to a map $\lambda_1(f, \cdot)$ from \mathbb{Z}^n to Γ_n .

- Next, for $z \in \mathbb{Z}^n$ we will use the “expansion rule” TB2 to construct a map $\tilde{\lambda}_2(\cdot, z)$ from $(S_A^{\pm 1})^*$ to Γ_n with $\tilde{\lambda}_2(f, z) = \tilde{\lambda}_1(f, z)$ for $f \in S_A^{\pm 1}$.
- Finally, we will show that $\tilde{\lambda}_2(f, \cdot)$ descends to a map $\lambda_2(f, \cdot)$ from $\text{Aut}(F_n)$ to Γ_n .

The desired twisted bilinear map will then be defined by $\lambda(f, z) = \lambda_2(f, z)$. It will follow from the various intermediate steps in our construction that $\lambda(\cdot, \cdot)$ is a twisted bilinear map.

As notation, for $w \in (S_A^{\pm 1})^*$, let \hat{w} denote the image of w in $\text{Aut}(F_n)$. Similarly, for $w \in (S_Z^{\pm 1})^*$, let \hat{w} denote the image of w in \mathbb{Z}^n .

We now construct $\tilde{\lambda}_1$. For $f \in S_A^{\pm 1}$ and $w \in (S_Z^{\pm 1})^*$, we define $\tilde{\lambda}_1(f, w) \in \Gamma_n$ by induction on the length of w . If $w = 1$ (i.e. w has length 0), then we define $\tilde{\lambda}_1(f, w) = 1$. If $w \in S_Z^{\pm 1}$ (i.e. w has length 1), then we define $\tilde{\lambda}_1(f, w)$ to be the value of $\lambda(f, w)$ from Table 6. Finally, if w has length at least 2 and $\tilde{\lambda}_1(f, \cdot)$ has been defined for all shorter words, then write $w = sw'$ with $s \in S_Z^{\pm 1}$ and define

$$\tilde{\lambda}_1(f, w) = \tilde{\lambda}_1(f, s) \cdot \beta_{\hat{s}} \left(\tilde{\lambda}_1(f, w') \right).$$

This formula should remind the reader of property TB1 from the definition of a twisted bilinear map, as should the following claim.

Claim 1. $\tilde{\lambda}_1(f, ww') = \tilde{\lambda}_1(f, w) \cdot \beta_{\hat{w}} \left(\tilde{\lambda}_1(f, w') \right)$ for $f \in S_A^{\pm 1}$ and $w, w' \in (S_Z^{\pm 1})^*$.

Proof of claim. The proof is by induction on the length of w . For w of length 0, this is trivial, and for w of length 1, it holds by definition. Now assume that w has length at least 2 and that the desired formula holds whenever w has smaller length. Write $w = w_1 w_2$, where w_1 and w_2 are shorter words than w . Applying our inductive hypothesis twice, we see that

$$\begin{aligned} \tilde{\lambda}_1(f, ww') &= \tilde{\lambda}_1(f, w_1 w_2 w') = \tilde{\lambda}_1(f, w_1) \cdot \beta_{\hat{w}_1} \left(\tilde{\lambda}_1(f, w_2 w') \right) \\ &= \tilde{\lambda}_1(f, w_1) \cdot \beta_{\hat{w}_1} \left(\tilde{\lambda}_1(f, w_2) \cdot \beta_{\hat{w}_2} \left(\tilde{\lambda}_1(f, w') \right) \right) \\ &= \tilde{\lambda}_1(f, w_1) \cdot \beta_{\hat{w}_1} \left(\tilde{\lambda}_1(f, w_2) \right) \cdot \beta_{\widehat{w_1 w_2}} \left(\tilde{\lambda}_1(f, w') \right). \end{aligned}$$

Applying our inductive hypothesis to the first two terms, we see that this equals

$$\tilde{\lambda}_1(f, w_1 w_2) \cdot \beta_{\widehat{w_1 w_2}} \left(\tilde{\lambda}_1(f, w') \right) = \tilde{\lambda}_1(f, w) \cdot \beta_{\hat{w}} \left(\tilde{\lambda}_1(f, w') \right). \quad \square$$

Claim 2. For $w, w' \in (S_Z^{\pm 1})^*$ with $\hat{w} = \hat{w}' \in \mathbb{Z}^n$, we have $\tilde{\lambda}_1(f, w) = \tilde{\lambda}_1(f, w')$ for $f \in S_A^{\pm 1}$.

Proof of claim. Recall that $\mathbb{Z}^n = \langle S_Z \mid R_Z \rangle$. Define $R'_Z = R_Z \cup \{ss^{-1} \mid s \in S_Z^{\pm 1}\} \subset (S_Z^{\pm 1})^*$. Since any two elements of $(S_Z^{\pm 1})^*$ that map to the same element of \mathbb{Z}^n must differ by a sequence of insertions and deletions of elements of R'_Z , we can assume without loss of generality that $w = uv$ and $w' = urv$ for some $u, v \in (S_Z^{\pm 1})^*$ and $r \in R'_Z$. A computer calculation described in Lemma 7.8 below shows that $\tilde{\lambda}_1(f, r) = 1$. We now apply Claim 1

several times to deduce that

$$\begin{aligned}\tilde{\lambda}_1(f, w') &= \tilde{\lambda}_1(f, urv) = \tilde{\lambda}_1(f, u) \cdot \beta_{\hat{f}\hat{u}} \left(\tilde{\lambda}_1(f, r) \right) \cdot \beta_{\hat{f}\hat{ur}} \left(\tilde{\lambda}_1(f, v) \right) \\ &= \tilde{\lambda}_1(f, u) \cdot \beta_{\hat{f}\hat{u}} \left(\tilde{\lambda}_1(f, v) \right) = \tilde{\lambda}_1(f, uv) = \tilde{\lambda}_1(f, w).\end{aligned}\quad \square$$

For $f \in S_A^{\pm 1}$, Claim 2 implies that the map $\tilde{\lambda}_1(f, \cdot)$ from $(S_Z^{\pm 1})^*$ to Γ_n descends to a map $\lambda_1(f, \cdot)$ from \mathbb{Z}^n to Γ_n . Claim 1 implies that $\lambda_1(f, \cdot)$ satisfies a version of condition TB1 from the definition of a twisted bilinear map, namely that $\lambda_1(f, z_1 z_2) = \lambda_1(f, z_1) \cdot \beta_{\hat{f}z_1}(\lambda_1(f, z_2))$ for all $z_1, z_2 \in \mathbb{Z}^n$. Our next claim is a version of condition TB3. We remark that the condition $f \in S_A$ in it is not a typo; we will extend it to $f \in S_A^{\pm 1}$ later.

Claim 3. $\lambda_1(f, z) \cdot \beta_{\hat{f}z} \left(\alpha_{\hat{f}}(k) \right) \cdot \lambda_1(f, z)^{-1} = \alpha_{\hat{f}}(\beta_z(k))$ for $f \in S_A$, $z \in \mathbb{Z}^n$, and $k \in \Gamma_n$.

Proof of claim. Let $w \in (S_Z^{\pm 1})^*$ satisfy $\hat{w} = z$. The proof is by induction on the length of w . For w of length 0, the claim is trivial. For w of length 1, there are two cases. For $w \in S_Z$, the claim follows from a computer calculation described below in Lemma 7.9. For $w = v^{-1}$ with $v \in S_Z$, Claim 1 implies that

$$1 = \tilde{\lambda}_1(f, v^{-1}v) = \tilde{\lambda}_1(f, v^{-1}) \cdot \beta_{\hat{f}\hat{v}^{-1}} \left(\tilde{\lambda}_1(f, v) \right),$$

so $\tilde{\lambda}_1(f, v^{-1}) = \beta_{\hat{f}\hat{v}^{-1}} \left(\tilde{\lambda}_1(f, v)^{-1} \right)$. Our goal is to show that

$$\tilde{\lambda}_1(f, v^{-1}) \cdot \beta_{\hat{f}\hat{v}^{-1}} \left(\alpha_{\hat{f}}(k) \right) \cdot \tilde{\lambda}_1(f, v^{-1})^{-1} = \alpha_{\hat{f}}(\beta_{\hat{v}^{-1}}(k)).$$

Plugging in our formula for $\tilde{\lambda}_1(f, v^{-1})$, we see that this is equivalent to showing that

$$\beta_{\hat{f}\hat{v}^{-1}} \left(\tilde{\lambda}_1(f, v)^{-1} \cdot \alpha_{\hat{f}}(k) \cdot \tilde{\lambda}_1(f, v) \right) = \alpha_{\hat{f}}(\beta_{\hat{v}^{-1}}(k)).$$

Manipulating this a bit, we see that it is equivalent to showing that

$$\alpha_{\hat{f}}(k) = \tilde{\lambda}_1(f, v) \cdot \beta_{\hat{f}\hat{v}} \left(\alpha_{\hat{f}}(\beta_{\hat{v}^{-1}}(k)) \right) \cdot \tilde{\lambda}_1(f, v)^{-1}.$$

Using the already proven case $w = v$ of the claim, the right hand side equals

$$\alpha_{\hat{f}}(\beta_{\hat{v}}(\beta_{\hat{v}^{-1}}(k))) = \alpha_{\hat{f}}(k),$$

as desired.

Now assume that w has length at least 2 and that the claim is true for all shorter words. Write $w = w_1 w_2$, where w_1 and w_2 are shorter words than w . Applying Claim 1, we see that $\tilde{\lambda}_1(f, w_1 w_2) \cdot \beta_{\hat{f}\widehat{w_1 w_2}} \left(\alpha_{\hat{f}}(k) \right) \cdot \tilde{\lambda}_1(f, w_1 w_2)^{-1}$ equals

$$\tilde{\lambda}_1(f, w_1) \cdot \beta_{\hat{f}\hat{w}_1} \left(\tilde{\lambda}_1(f, w_2) \cdot \beta_{\hat{f}\hat{w}_2} \left(\alpha_{\hat{f}}(k) \right) \cdot \tilde{\lambda}_1(f, w_2)^{-1} \right) \cdot \tilde{\lambda}_1(f, w_1)^{-1}. \quad (13)$$

Our inductive hypothesis implies that

$$\tilde{\lambda}_1(f, w_2) \cdot \beta_{\hat{f}_{\hat{w}_2}} \left(\alpha_{\hat{f}}(k) \right) \cdot \tilde{\lambda}_1(f, w_2)^{-1} = \alpha_{\hat{f}}(\beta_{\hat{w}_2}(k)).$$

Thus (13) equals

$$\tilde{\lambda}_1(f, w_1) \cdot \beta_{\hat{f}_{\hat{w}_1}} \left(\alpha_{\hat{f}}(\beta_{\hat{w}_2}(k)) \right) \cdot \tilde{\lambda}_1(f, w_1)^{-1}.$$

Another application of our inductive hypothesis shows that this equals

$$\alpha_{\hat{f}}(\beta_{\hat{w}_1}(\beta_{\hat{w}_2}(k))) = \alpha_{\hat{f}}(\beta_{\widehat{w_1 w_2}}(k)). \quad \square$$

We now construct $\tilde{\lambda}_2$. For $w \in (S_A^{\pm 1})^*$ and $z \in \mathbb{Z}^n$, we define $\tilde{\lambda}_2(w, z) \in \Gamma_n$ by induction on the length of w . If $w = 1$ (i.e. w has length 0), then we define $\tilde{\lambda}_2(w, z) = 1$. If $w \in S_A^{\pm 1}$ (i.e. w has length 1), then we define $\tilde{\lambda}_2(w, z) = \lambda_1(w, z)$. Finally, if w has length at least 2 and $\tilde{\lambda}_2(\cdot, z)$ has been defined for all shorter words, then write $w = sw'$ with $s \in S_A^{\pm 1}$ and define

$$\tilde{\lambda}_2(w, z) = \alpha_{\hat{s}} \left(\tilde{\lambda}_2(w', z) \right) \cdot \tilde{\lambda}_2(a_1, \hat{w}'z).$$

This formula should remind the reader of property TB2 from the definition of a twisted bilinear map, as should the following claim.

Claim 4. $\tilde{\lambda}_2(ww', z) = \alpha_{\hat{w}} \left(\tilde{\lambda}_2(w', z) \right) \cdot \tilde{\lambda}_2(w, \hat{w}'z)$ for $w, w' \in (S_A^{\pm 1})^*$ and $z \in \mathbb{Z}^n$.

Proof of claim. This can be proved by induction on the length of w just like Claim 1. The details are left to the reader. \square

The reader might expect at this point that we would prove an analogue of Claim 2 and thus show that $\tilde{\lambda}_2$ descends to a map $\lambda_2: \text{Aut}(F_n) \times \mathbb{Z}^n \rightarrow \Gamma_n$. However, before we can do this we must prove two preliminary results. The first extends Claim 3 to show that $\tilde{\lambda}_2$ satisfies a version of condition TB3.

Claim 5. $\tilde{\lambda}_2(w, z) \cdot \beta_{\hat{w}_z}(\alpha_{\hat{w}}(k)) \cdot \tilde{\lambda}_2(w, z)^{-1} = \alpha_{\hat{w}}(\beta_z(k))$ for $w \in (S_A^{\pm 1})^*$, $z \in \mathbb{Z}^n$ and $k \in \Gamma_n$.

Proof of claim. This can be proved by induction on the length of w just like Claim 3. The details are left to the reader. \square

The next claim extends Claim 1 to show that $\tilde{\lambda}_2$ satisfies a version of condition TB1.

Claim 6. $\tilde{\lambda}_2(w, zz') = \tilde{\lambda}_2(w, z) \cdot \beta_{\hat{w}_z} \left(\tilde{\lambda}_2(w, z') \right)$ for $w \in (S_A^{\pm 1})^*$ and $z, z' \in \mathbb{Z}^n$.

Proof of claim. The proof is by induction on the length of w . For w of length 0, this is trivial, and for w of length 1, it holds by Claim 1. Now assume that w has length at least 2 and that the desired formula holds whenever w has smaller length. Write $w = w_1 w_2$, where

w_1 and w_2 are shorter words than w . Applying Claim 4 and our inductive hypothesis, we see that

$$\begin{aligned}\tilde{\lambda}_2(w, zz') &= \alpha_{\hat{w}_1} \left(\tilde{\lambda}_2(w_2, zz') \right) \cdot \tilde{\lambda}_2(w_1, \hat{w}_2 z \hat{w}_2 z') \\ &= \alpha_{\hat{w}_1} \left(\tilde{\lambda}_2(w_2, z) \cdot \beta_{\hat{w}_2 z} \left(\tilde{\lambda}_2(w_2, z') \right) \right) \cdot \tilde{\lambda}_2(w_1, \hat{w}_2 z) \cdot \beta_{\hat{w}_1 z} \left(\tilde{\lambda}_2(w_1, \hat{w}_2 z') \right).\end{aligned}\quad (14)$$

Also, Claim 4 implies that $\tilde{\lambda}_2(w, z) \cdot \beta_{\hat{w} z} \left(\tilde{\lambda}_2(w, z') \right)$ equals

$$\alpha_{\hat{w}_1} \left(\tilde{\lambda}_2(w_2, z) \right) \cdot \tilde{\lambda}_2(w_1, \hat{w}_2 z) \cdot \beta_{\hat{w}_1 \hat{w}_2 z} \left(\alpha_{\hat{w}_1} \left(\tilde{\lambda}_2(w_2, z') \right) \cdot \tilde{\lambda}_2(w_1, \hat{w}_2 z') \right).\quad (15)$$

Our goal is to prove that (14) equals (15). Manipulating this, we see that our goal is equivalent to showing that

$$\tilde{\lambda}_2(w_1, \hat{w}_2 z) \cdot \beta_{\hat{w}_1 \hat{w}_2 z} \left(\alpha_{\hat{w}_1} \left(\tilde{\lambda}_2(w_2, z') \right) \right) \cdot \tilde{\lambda}_2(w_1, \hat{w}_2 z)^{-1} = \alpha_{\hat{w}_1} \left(\beta_{\hat{w}_2 z} \left(\tilde{\lambda}_2(w_2, z') \right) \right).$$

This is an immediate consequence of Claim 5. \square

We finally prove the promised analogue of Claim 2.

Claim 7. For $w, w' \in (S_A^{\pm 1})^*$ with $\hat{w} = \hat{w}' \in \text{Aut}(F_n)$, we have $\tilde{\lambda}_2(w, z) = \tilde{\lambda}_2(w', z)$ for $z \in \mathbb{Z}^n$.

Proof of claim. Recall that $\text{Aut}(F_n) = \langle S_A \mid R_A \rangle$. Define $R'_A = R_A \cup \{ss^{-1} \mid s \in S_A^{\pm 1}\} \subset (S_A^{\pm 1})^*$. A computer calculation described below in Lemma 7.10 shows that $\tilde{\lambda}_2(r, \hat{s}) = 1$ for $r \in R'_A$ and $s \in S_Z^{\pm 1}$. Writing z as a product of elements of $S_Z^{\pm 1}$, we can use Claim 6 to show that $\tilde{\lambda}_2(r, z) = 1$ for $r \in R'_A$. The proof now is identical to the proof of Claim 2; the details are left to the reader. \square

Claim 7 implies that $\tilde{\lambda}_2$ descends to a map $\lambda_2: \text{Aut}(F_n) \times \mathbb{Z}^n \rightarrow \Gamma_n$. This map is a twisted bilinear map: Claim 6 implies that it satisfies condition TB1, Claim 4 implies that it satisfies condition TB2, and Claim 5 implies that it satisfies condition TB3. As discussed at the beginning of the proof, $\lambda = \lambda_2$ is the twisted bilinear map whose existence we are trying to prove. \square

6.3 Proof of L-presentation

We now prove Theorem 6.2.

Proof of Theorem 6.2. Let $\Gamma_n = \langle S_K \mid R_K^0 \mid \phi(S_Q^{\pm 1}) \rangle$ be the group with the presentation described in Theorem 6.2. We map each generator of Γ_n to the generator of $\mathcal{K}_{n,1}^{\text{IA}}$ with the same name. Lemma 7.11 below checks that the basic relations R_K^0 are true in $\mathcal{K}_{n,1}^{\text{IA}}$; it then follows from the naturality from Lemmas 6.3 and 6.4 that the extended relations of Γ_n are also true in $\mathcal{K}_{n,1}^{\text{IA}}$. Therefore we have defined a homomorphism $\Psi: \Gamma_n \rightarrow \mathcal{K}_{n,1}^{\text{IA}}$. Since our

generating set from Theorem A is in the image of Ψ , we know Ψ is a surjection; our goal is to show that Ψ is an isomorphism.

For $f \in \text{Aut}(F_n)$, let $\alpha_f: \Gamma_n \rightarrow \Gamma_n$ be the homomorphism given by Lemma 6.3. Also, for $z \in \mathbb{Z}^n$, let $\beta_z: \Gamma_n \rightarrow \Gamma_n$ be the homomorphism given by Lemma 6.4. Finally, let $\lambda: \text{Aut}(F_n) \times \mathbb{Z}^n \rightarrow \Gamma_n$ be the twisted bilinear map given by Lemma 6.5. Plugging this data into Theorem 5.5, we obtain a short exact sequence

$$1 \longrightarrow \Gamma_n \longrightarrow \Delta_n \xrightarrow{\rho} \mathbb{Z}^n \rtimes_r \text{Aut}(F_n) \longrightarrow 1$$

together with homomorphisms $\iota_1: \text{Aut}(F_n) \rightarrow \Delta_n$ and $\iota_2: \mathbb{Z}^n \rightarrow \Delta_n$ such that $\rho \circ \iota_1 = \text{id}$ and $\rho \circ \iota_2 = \text{id}$. The naturality properties of the data in Lemmas 6.3, 6.4, and 6.5 imply that this short exact sequence fits into a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Gamma_n & \longrightarrow & \Delta_n & \longrightarrow & \mathbb{Z}^n \rtimes \text{Aut}(F_n) \longrightarrow 1 \\ & & \downarrow \Psi & & \downarrow \Phi & & \downarrow = \\ 1 & \longrightarrow & \mathcal{K}_{n,1}^{\text{IA}} & \longrightarrow & \mathcal{A}_{n,1} & \longrightarrow & \mathbb{Z}^n \rtimes \text{Aut}(F_n) \longrightarrow 1. \end{array} \quad (16)$$

By the five lemma, we see that to prove that Ψ is an isomorphism, it is enough to prove that Φ is an isomorphism. We will do this by constructing an explicit inverse homomorphism $\Phi^{-1}: \mathcal{A}_{n,1} \rightarrow \Delta_n$.

To do this, we first need some explicit elements of Δ_n and some relations between those elements. The needed elements are as follows.

- We will identify the generating set

$$S_K = \{\mathfrak{C}_{y,x_a}, \mathfrak{C}_{x_a,y} \mid x_a \in X\} \cup \{\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]} \mid x_a, x_b \in X, x_a \neq x_b, \alpha, \beta, \epsilon \in \{1, -1\}\}$$

for Γ_n with its image in Δ_n .

- For $\alpha \in \{1, -1\}$ and distinct $x_a, x_b \in X$, we define $\mathfrak{M}_{x_a^\alpha, x_b}$ to equal $\iota_A(M_{x_a^\alpha, x_b})$.
- For distinct $x_a, x_b \in X$, we define $\mathfrak{P}_{a,b}$ to equal $\iota_A(P_{a,b})$.
- For $x_a \in X$, we define \mathfrak{I}_a to equal $\iota_A(I_a)$.
- As in the proof of Lemma 6.4, we will regard \mathbb{Z}^n as being generated by the set $\{M_{x_a, y} \mid x_a \in X\}$, and for $x_a \in X$ we define $\mathfrak{M}_{x_a, y}$ to equal $\iota_B(M_{x_a, y})$.

The needed relations are as follows. That they hold is immediate from the construction of Δ_n in the proof of Theorem 5.5.

- The relations R_K^0 from the L-presentation for Γ_n .
- By construction, the group Δ_n contains subgroups $\Gamma_n \rtimes \text{Aut}(F_n)$ and $\Gamma_n \rtimes \mathbb{Z}^n$. Any relation which holds in $\Gamma_n \rtimes \text{Aut}(F_n)$ or $\Gamma_n \rtimes \mathbb{Z}^n$ (which are generated by the evident elements) also holds in Δ_n .
- For $f \in \text{Aut}(F_n)$ and $z \in \mathbb{Z}^n$, we have $\lambda(f, z) = fz f^{-1} z^{-1}$. Here fz comes from the action of $\text{Aut}(F_n)$ on \mathbb{Z}^n in the semidirect product $\mathbb{Z}^n \rtimes \text{Aut}(F_n)$. Also, $f \in \text{Aut}(F_n)$ and $z \in \mathbb{Z}^n$ and $fz \in \mathbb{Z}^n$ should be identified with their images in Δ_n .

Let $\mathcal{A}_{n,1} = \langle S_C \mid R_C \rangle$ be the presentation given by Theorem 5.7, so

$$\begin{aligned} S_C = & \{M_{x_a^\alpha, x_b} \mid 1 \leq a, b \leq n \text{ distinct}, \alpha \in \{1, -1\}\} \cup \{P_{a,b} \mid 1 \leq a < b \leq n\} \\ & \cup \{I_a \mid 1 \leq a \leq n\} \cup \{M_{x_a^\alpha, y}, C_{y, x_a} \mid 1 \leq a \leq n, \alpha \in \{1, -1\}\}. \end{aligned}$$

We define a set map $\tilde{\Phi}^{-1}: S_C \rightarrow \Delta_n$ as follows. First, most of the elements in S_C have evident analogues in Δ_n , so we define

$$\begin{aligned} \tilde{\Phi}^{-1}(M_{x_a^\alpha, x_b}) &= \mathfrak{M}_{x_a^\alpha, x_b} & \text{and} & & \tilde{\Phi}^{-1}(P_{a,b}) &= \mathfrak{P}_{a,b} & \text{and} & & \tilde{\Phi}^{-1}(I_a) &= \mathfrak{I}_a \\ \text{and } \tilde{\Phi}^{-1}(M_{x_a, y}) &= \mathfrak{M}_{x_a, y} & \text{and} & & \tilde{\Phi}^{-1}(C_{y, x_a}) &= \mathfrak{C}_{y, x_a}. \end{aligned}$$

The only remaining element of S_C is $M_{x_a^{-1}, y}$, and we define

$$\tilde{\Phi}^{-1}(M_{x_a^{-1}, y}) = \mathfrak{C}_{x_a, y} \mathfrak{M}_{x_a, y}^{-1}.$$

The map $\tilde{\Phi}^{-1}$ extends to a homomorphism $\tilde{\Phi}^{-1}: F(S_C) \rightarrow \Delta_n$. Computer calculations described in Lemma 7.12 below show that $\tilde{\Phi}^{-1}(r) = 1$ for $r \in R_C$, so $\tilde{\Phi}^{-1}$ descends to a homomorphism $\Phi^{-1}: \mathcal{A}_{n,1} \rightarrow \Delta_n$. Examining its effect on generators, we see that Φ^{-1} is the desired inverse to Φ , and the proof is complete. \square

7 Computer calculations

This section discusses the computer calculations used in the previous section. The preliminary section §7.1 discusses the basic framework we use. The actual computations are in §7.2.

7.1 Framework for calculations

We model $\text{Aut}(F_{n+1})$ using GAP, a software algebra system available for free at <http://www.gap-system.org/>. We encourage our readers to experiment with the included functions, and to look at the code that performs the verifications below. We use the same framework that the authors used in [6], so we quote part of our explanation of the framework from there. From [6]:

We use GAP's built-in functionality to model F_n as a free group on the eight generators \mathbf{xa} , \mathbf{xb} , \mathbf{xc} , \mathbf{xd} , \mathbf{xe} , \mathbf{xf} , \mathbf{xg} , and \mathbf{y} . Since our computations never involve more than 8 variables, computations in this group suffice to show that our computations hold in general.

We found it more convenient to model the free groups $F(S_A)$, $F(S_Q)$, and $F(S_K)$ without using the built-in free group functionality. Instead we model the generators using lists and program the basic free group operations directly. Continuing from [6]:

For example, we model the generator M_{x_a, x_b} as the list `["M", xa, xb]`, C_{y, x_a} as `["C", y, xa]`, and $M_{x_a^{-1}, [y, x_c]}$ as `["Mc", xa^-1, y, xc]`. We model $P_{a,b}$ as `["P", xa, xb]` and I_a as `["I", xa]`. The examples should make clear: the first entry in the list is a string key "M", "C", "Mc", "P", or "I", indicating whether the list represents a transvection, conjugation move, commutator transvection, swap or inversion. The parameters given as subscripts in the generator are then the remaining elements of the list, in the same order.

We model words in any of the free groups $F(S_A)$, $F(S_Q)$, and $F(S_K)$ as lists of generators. Continuing from [6]:

We model inverses of generators as follows: the inverse of ["M",xa,xb] is ["M",xa,xb^-1] and the inverse of ["C",xa,xb] is ["C",xa,xb^-1], but the inverse of ["Mc",xa,xb,xc] is ["Mc",xa,xc,xb]. Swaps and inversions are their own inverses. Technically, this means that ... we model structures where the order relations for swaps and inversions and the relation ... for inverting commutator transvections are built in. This is not a problem because our verifications always show that certain formulas are trivial modulo our relations

....

In particular, the inverse of ["Mc",xa,y,xb] is modeled as ["Mc",xa,xb,y]. Continuing from [6]:

The empty word [] represents the trivial element. We wrote several functions ... that perform common tasks on words. The function `pw` takes any number of words (reduced or not) as arguments and returns the freely reduced product of those words in the given order, as a single word. The function `iw` inverts its input word and the function `cyw` cyclically permutes its input word.

... The function `applyrels` is particularly useful, because it inserts multiple relations into a word. It takes in two inputs: a starting word and a list of words with placement indicators. The function recursively inserts the first word from the list in the starting word at the given position, reduces the word, and then calls itself with the new word as the starting word and with the same list of insertions, with the first dropped.

Most of the verifications amount to showing that some formula can be expressed as a product of conjugates of images of relations under the substitution rules. We model the substitution rule function ϕ using a function named `phi`. This takes a word in $(S_Q^{\pm 1})^*$ as its first input and a word from $F(S_K)$ as its second input and applies to the second word the composition of substitution rules given by the first one. We use a function `krel` to encode the basic relations R_K^0 from Theorem 6.2. Given a number n and a list of basis elements (or inverse basis elements) from F_{n+1} , `krel` returns the n th relation from Table 5, with the supplied basis elements as subscripts on the S_K -generators. If the parameters are inconsistent, it returns the empty word. We define a function `psi` that encodes the action of $\text{Aut}(F_n)$ on \mathbb{Z}^n from Section 5.1. We also defines a function `lambda` that computes the definition of $\tilde{\lambda}_2$ above; in the special case that its first input is in $S_A^{\pm 1}$, it computes $\tilde{\lambda}_1$ as well (this is also true of $\tilde{\lambda}_2$ by definition).

The functions described here and the checklists for the computations described below are all given in the file `BirmanIA.g`, which we make available with this paper. Each of the following lemmas refers to lists of outputs in `BirmanIA.g`. To check the validity of a given lemma, one needs to read the code that generates the list, evaluate the code, and make sure the output is correct (usually the desired output is a list of copies of the trivial word). We

have provided a list `BirmanIAChecklist` that gives the output of all the verifications in the paper.

7.2 The actual calculations

In addition to the relations from Table 5, we use some derived relations for convenience. These are output by a function `exkrel` and we do not list them here (they can be found by inspecting the code and the outputs from that function). What matters is that these relations always follow from the relations in the presentation.

Lemma 7.1. *All the relations output by `exkrel` are true in Γ_n .*

Proof. The source for the list `exkrellist` contains a reduction of one instance of each output of `exkrel` to the trivial word using only outputs from `krel`, images of outputs from `krel` under the action of `phi`, and previously verified relations from `exkrel`. Each of the entries in the list evaluates to the trivial word, so the reductions are correct. \square

There are a few places where we verify identities that are homomorphisms on both sides. To verify these most efficiently, we use generating sets for Γ_n that are smaller than S_K .

Lemma 7.2. *Suppose S is a set containing all the $\{\mathfrak{C}_{x_a, y}\}_a$ and $\{\mathfrak{C}_{y, x_a}\}_a$, and for each choice of a, b , suppose S contains at least one of the eight elements $\{\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]}\}_{\alpha, \beta, \epsilon}$. Then S is a generating set for Γ_n .*

Proof. Suppose S contains $\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^\beta]}$ and all the conjugation moves above. Then: we can use R4 to express $\mathfrak{M}_{x_a^\alpha, [y^\epsilon, x_b^{-\beta}]}$ in terms of elements of S , we can use R5 to express $\mathfrak{M}_{x_a^\alpha, [y^{-\epsilon}, x_b^\beta]}$ in terms of elements of S , and we can use R7 to express $\mathfrak{M}_{x_a^{-\alpha}, [y^\epsilon, x_b^\beta]}$ in terms of elements of S . Applying these relations repeatedly allows us to get all of the eight commutator transvections involving x_a and x_b from one of them. \square

Lemma 7.3. *Let $\Psi: F(S_K) \rightarrow \mathcal{K}_{n,1}^{IA}$ be the natural surjection. Then regarding S_Q as a subset of $\mathcal{A}_{n,1}$, we have*

$$\Psi(\phi(s)(t)) = s\Psi(t)s^{-1} \quad (s \in S_Q, t \in S_K).$$

Proof. First we note that this is clearly true, by definition, if s is a swap or inversion. Further, if we verify this for $s = \mathfrak{M}_{x_a, x_b}$, then it follows for $s = \mathfrak{M}_{x_a^{-1}, x_b}$ by conjugating the entire expression by an inversion. So it is enough to verify it for s of the form \mathfrak{M}_{x_a, x_b} or $\mathfrak{M}_{x_a, y}$. In the code generating the list `phiconjugationlist`, we check this equation for both such choices of s , and for all possible configurations of generator t with respect to the choice of s . \square

Lemma 7.4. *The map $\zeta: (S_A^{\pm 1})^* \rightarrow \text{End}(\Gamma_n)$ induced by $\phi: S_Q^{\pm 1} \rightarrow \text{End}(F(S_K))$ satisfies*

$$\zeta(s)(\zeta(s^{-1})(t)) = t$$

for all $s \in S_A^{\pm 1}$ and all $t \in S_K$.

Proof. In fact, it is enough to show this for t in a generating set for Γ_n , since then $\zeta(s)\circ\zeta(s^{-1})$ is the identity endomorphism of Γ_n . If s is a swap or inversion, then the lemma follows immediately from the definition. So we verify that $\phi(s)(\phi(s^{-1})(t)) = t$ (up to relations of Γ_n) for s of the form \mathfrak{M}_{x_a, x_b} , and for enough choices of t to give a generating set for Γ_n (using Lemma 7.2). This computation appears in the code generating the list `phiAinverseList`. For s of the form $\mathfrak{M}_{x_a^{-1}, x_b}$, our computations for $s = \mathfrak{M}_{x_a, x_b}$ suffice, after substituting x_a for x_a^{-1} in each computation. \square

Lemma 7.5. *The map $\eta: F(S_A) \rightarrow \text{Aut}(\Gamma_n)$ induced by $\phi: S_Q^{\pm 1} \rightarrow \text{End}(F(S_K))$ satisfies*

$$\eta(r)(t) = t$$

for all $t \in S_K$ and for every relation $r \in R_A$ from Nielsen's presentation for $\text{Aut}(F_n) = \langle S_A | R_A \rangle$.

Proof. It is enough to check that the equation

$$\phi(r)(t) = t \tag{17}$$

holds in Γ_n for every relation r from Nielsen's presentation, for choices of t ranging through a generating set for Γ_n .

Equation (17) works automatically for r a relation of type N1, since the action ϕ is defined for swaps and inversions using the natural action on $\{x_1^{\pm 1}, \dots, x_n^{\pm 1}\}$, and these relations hold for that action.

Equation (17) also works automatically for r a relation of type N2. In this case, the equation says that $s \mapsto \phi(s)$, for s a transvection, is equivariant with respect to the action of swaps and inversions. This is apparent from the definition of ϕ : the definition does not refer to specific elements x_i , but instead treats configurations the same way based on coincidences between them.

For the other cases, we use computations given in the source code for the lists `phiN3list`, `phiN4list`, and `phiN5list`. In each list we select a relation r and reduce $\phi(r)(t)t^{-1}$ to 1 in Γ_n , for choices of t constituting a generating set by Lemma 7.2. In each list we exploit natural symmetries of the equation to reduce the number of cases considered. \square

Lemma 7.6. *The map $\zeta: (S_Z^{\pm 1})^* \rightarrow \text{End}(\Gamma_n)$ induced by $\phi: S_Q^{\pm 1} \rightarrow \text{End}(F(S_K))$ satisfies*

$$\zeta(s)(\zeta(s^{-1})(t)) = t$$

for all $s \in S_Z^{\pm 1}$ and $t \in S_K$.

Proof. This is like the proof of Lemma 7.4, but simpler. We verify that $\phi(s)(\phi(s^{-1})(t)) = t$ in Γ_n for s of the form $\mathfrak{M}_{x_a, y}$, for t ranging over a generating set for Γ_n . This computation is in the code generating the list `phiZinverseList`. \square

Lemma 7.7. *The map $\eta: F(S_Z) \rightarrow \text{Aut}(\Gamma_n)$ induced by $\phi: S_Q^{\pm 1} \rightarrow \text{End}(F(S_K))$ satisfies*

$$\eta(r)(t) = t,$$

whenever r is a basic commutator of generators from S_Z and $t \in S_K$.

Proof. The computations showing this appear in `phiznlist`. □

Lemma 7.8. *For r of the form ss^{-1} for $s \in S_Z$, or $[s, t]$ for $s, t \in S_Z$, we have*

$$\tilde{\lambda}_1(f, r) = 1$$

in Γ_n for any $f \in S_A^{\pm 1}$.

Proof. The meaning here is that we must expand $\tilde{\lambda}_1(f, r)$ according the definition without simplifying r (not even cancelling inverse pairs), and then verify that the expression we get is a relation in Γ_n . To check the lemma, it is enough to verify that the function `lambda` returns relations in Γ_n when the first input is a generator or inverse generator and the second input is commutator or the product of a generator and its inverse. The computations for this lemma this appear in `lambda2ndinverselist` and `lambda2ndrellist`. □

Lemma 7.9. *We have*

$$\lambda_1(f, z) \cdot \beta_{\hat{f}_z} \left(\alpha_{\hat{f}}(t) \right) \cdot \lambda_1(f, z)^{-1} = \alpha_{\hat{f}}(\beta_z(t))$$

for $f \in S_A$, $z \in S_Z$, and $t \in \Gamma_n$.

Proof. The function `psi` takes as input a word a in S_A and a word b in S_Z and returns a word in S_Z representing $^a b$. Since the actions α and β are given by ϕ , we may rewrite the expression we are trying to prove as

$$\lambda_1(f, z) \cdot \phi(\psi(f)(z)) \circ \phi(f)(t) \cdot \lambda_1(f, z)^{-1} = \phi(f) \circ \phi(z)(t).$$

We note that this equation is an automorphism of Γ_n on both sides, so it is enough to verify it for t in a generating set. Computations checking this identity for all choices of f , z and t appear in the code generating `tb3list`. □

Lemma 7.10. *Suppose $r \in (S_A)^*$ is one of Nielsen's relations for $\text{Aut}(F_n)$ or is a product ff^{-1} for some $f \in S_A^{\pm 1}$. Then for any $z \in \mathbb{Z}^n$, we have*

$$\lambda_2(r, z) = 1$$

in Γ_n .

Proof. It is enough to show this for generators of \mathbb{Z}^n . We show this using the function `lambda` that encodes the definition of λ_2 . The code checking these identities generates the lists `lambda1stinversecheck`, `lambdaN1list`, `lambdaN2list`, `lambdaN3list`, `lambdaN4list`, and `lambdaN5list`. □

Lemma 7.11. *The basic relations R_K^0 of the L -presentation for Γ_n are true when interpreted as identities in $\mathcal{K}_{n,1}^{IA}$.*

Proof. This is verified in the list `verifyGammarellist`, which uses the function `krel` to generate the relations. All generic and non-generic forms of the relations are checked separately. \square

Lemma 7.12. *The relations from Jensen–Wahl’s presentation for $\mathcal{A}_{n,1}$ map to relations of Δ_n under the map $\tilde{\Phi}^{-1}$, so the map as defined on generators extends to a well defined homomorphism $\Phi^{-1}: \mathcal{A}_{n,1} \rightarrow \Delta_n$.*

Proof. This is verified in the list `JWfromDeltalist`. \square

References

- [1] L. Bartholdi, Endomorphic presentations of branch groups, *J. Algebra* 268 (2003), no. 2, 419–443.
- [2] J. S. Birman, Mapping class groups and their relationship to braid groups, *Comm. Pure Appl. Math.* **22** (1969), 213–238.
- [3] K. S. Brown, *Cohomology of groups*, corrected reprint of the 1982 original, Graduate Texts in Mathematics, 87, Springer, New York, 1994.
- [4] F. Cohen and J. Pakianathan, On automorphism groups of free groups and their nilpotent quotients, in preparation.
- [5] M. Day and A. Putman, A Birman exact sequence for $\text{Aut}(F_n)$, *Adv. Math.* 231 (2012), no. 1, 243–275.
- [6] M. Day and A. Putman, On the second homology group of the Torelli subgroup of $\text{Aut}(F_n)$, preprint 2014.
- [7] S. Eilenberg and S. MacLane, Cohomology Theory in Abstract Groups. II: Group extensions with a non-Abelian Kernel, *Ann. of Math.*, **48** (1947), no. 2, 326–341.
- [8] B. Farb, Automorphisms of F_n which act trivially on homology, in preparation.
- [9] B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, 49, Princeton Univ. Press, Princeton, NJ, 2012.
- [10] C. A. Jensen and N. Wahl, Automorphisms of free groups with boundaries, *Algebr. Geom. Topol.* **4** (2004), 543–569.
- [11] N. Kawazumi, Cohomological aspects of Magnus expansions, preprint 2005.
- [12] T. Satoh, A survey of the Johnson homomorphisms of the automorphism groups of free groups and related topics, preprint 2013, [arXiv:1204.0876](https://arxiv.org/abs/1204.0876).
- [13] J. Nielsen, Die Isomorphismengruppe der freien Gruppen. (German) *Math. Ann.* **91** (1924), no. 3-4, 169–209.

Matthew Day
 Department of Mathematical Sciences, 309 SCEN
 University of Arkansas
 Fayetteville, AR 72701
 E-mail: matthewd@uark.edu

Andrew Putman
 Department of Mathematics
 Rice University, MS 136
 6100 Main St.
 Houston, TX 77005
 E-mail: andyp@math.rice.edu