

Name: _____

Instructor: _____

Math 10550, EXAM II
October 13, 3016

- The Honor Code is in effect for this examination. All work is to be your own.
- No calculators.
- The exam lasts for 1 hr. and 15 min.
- Be sure that your name is on every page in case pages become detached.
- Be sure that you have all 18 pages of the test.
- Each multiple choice question is worth 7 points. Your score will be the sum of the best 10 scores on the multiple choice questions plus your score on questions 13-15.

PLEASE MARK YOUR ANSWERS WITH AN X, not a circle!					
1.	(a)	(b)	(c)	(d)	(e)
2.	(a)	(b)	(c)	(d)	(e)
.....					
3.	(a)	(b)	(c)	(d)	(e)
4.	(a)	(b)	(c)	(d)	(e)
.....					
5.	(a)	(b)	(c)	(d)	(e)
6.	(a)	(b)	(c)	(d)	(e)
.....					
7.	(a)	(b)	(c)	(d)	(e)
8.	(a)	(b)	(c)	(d)	(e)
.....					
9.	(a)	(b)	(c)	(d)	(e)
10.	(a)	(b)	(c)	(d)	(e)
.....					
11.	(a)	(b)	(c)	(d)	(e)
12.	(a)	(b)	(c)	(d)	(e)

Please do NOT write in this box.

Multiple Choice _____

13. _____

14. _____

15. _____

Total _____

Name: _____

Instructor: _____

Multiple Choice

1.(6 pts.) Find y' , if

$$x^4 + x^3y + 5xy^2 = 8.$$

(a) $\frac{-(4x^3 + 3x^2y + 5y^2)}{10xy}$

(b) $\frac{-(4x^3 + 3x^2y + 5y^2)}{x^3}$

(c) $\frac{-(4x^3 + 3x^2y)}{x^3 + 10xy}$

(d) The derivative does not exist.

(e) $\frac{-(4x^3 + 3x^2y + 5y^2)}{x^3 + 10xy}$

Solution: We differentiate both sides with respect to x

$$\frac{d(x^4 + x^3y + 5xy^2)}{dx} = \frac{d(8)}{dx}$$

$$4x^3 + 3x^2y + x^3\frac{dy}{dx} + 5y^2 + 10xy\frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{-4x^3 - 3x^2y - 5y^2}{x^3 + 10xy}$$

Name: _____

Instructor: _____

2.(6 pts.) If $\sin(\pi xy) = \pi(x + y)$ find $\frac{dy}{dx}$ at $(1, -1)$ by implicit differentiation.

Solution: Differentiating both sides with respect to x ,

$$\cos(\pi xy) \left(\pi y + \pi x \frac{dy}{dx} \right) = \pi \left(1 + \frac{dy}{dx} \right).$$

Now, letting $x = 1$, $y = -1$ we get

$$\pi \cos(-\pi) \left(-1 + \frac{dy}{dx} \right) = \pi \left(1 + \frac{dy}{dx} \right).$$

Isolating dy/dx we get

$$\frac{dy}{dx} = 0.$$

- (a) $\frac{\pi}{2}$ (b) 0 (c) 1 (d) -1 (e) π

Name: _____

Instructor: _____

3.(6 pts.) A particle is moving in a straight line along a horizontal axis with a position function given by

$$s(t) = t^2 - 4t + 4,$$

where distance is measured in feet and time is measured in seconds. What is the distance travelled by the particle in the time period $1 \leq t \leq 4$ seconds?

SOLUTION:

Recall that the distance is given by how much has the particle travelled, thus we need to look for any point where our particle changed direction. To do so we consider where (if any) does the derivative of s change signs. We compute

$$s'(t) = 2t - 4$$

Which is zero at $t = 2$ and since it is an increasing line we know $s'(t) < 0$ for $t < 2$ and $s'(t) > 0$ for $t > 2$. Thus it changes direction on $[1, 4]$ only at $t = 2$. Hence the total travelled distance (since the particle is moving in a straight line along a horizontal axis), in feet, is given by

$$D = |s(4) - s(2)| + |s(2) - s(1)| = |4 - 0| + |0 - 1| = 5.$$

- (a) 5 feet (b) 8 feet (c) 3 feet
(d) 0 feet (e) 2 feet

Name: _____

Instructor: _____

4.(6 pts.) A beetle travels in straight line with position $s(t)$ (measured in feet) at time t (measured in seconds), $t \geq 0$. $s(t)$ is given by

$$s(t) = 3t^4 - 20t^3 + 36t^2.$$

At what time, after the motion gets started, does the beetle first come to rest?

Solution: We need to compute the critical points:

$$\begin{aligned} s'(t) = 12t^3 - 60t^2 + 72t &= 0 \\ 12t(t^2 - 5t + 6) &= 0 \end{aligned}$$

The critical points are $t = 0, 2, 3$, so after the motion starts, the beetle come to rest at $t = 2$.

- (a) $t = 1$ (b) $t = 3$ (c) $t = 2$
(d) This beetle never stops(e) $t = 4$

Name: _____

Instructor: _____

5.(6 pts.) A right triangle has base x feet and height y feet. If the base increases at 2 ft/second, and the height increases at 1 ft/second, find the rate of change in the area of the right triangle when $x = 8$ and $y = 5$.

SOLUTION:

The formula for the area A of a triangle with base x in feet and height y in feet is given by $A = \frac{1}{2}xy$. Differentiating with respect to t , we get that

$$\frac{dA}{dt} = \frac{1}{2} \frac{dx}{dt} y + \frac{1}{2} x \frac{dy}{dt}$$

Plugging in the values given in the problem, we see that

$$\begin{aligned} \frac{dA}{dt} &= \frac{1}{2}(2)(5) + \frac{1}{2}(8)(1) \\ &= 9 \end{aligned}$$

- (a) 2 ft²/second (b) 10.5 ft²/second (c) 18 ft²/second
(d) -1 ft²/second (e) 9 ft²/second

Name: _____

Instructor: _____

6.(6 pts.) Suppose f is differentiable and $-2 \leq f'(x) \leq 1$ for all x and $f(2) = 3$. What are the minimum and maximum possible values for $f(5)$?

SOLUTION:

Since f is differentiable we can apply the Mean Value Theorem on $[2, 5]$ to get the existence of an $x \in (2, 5)$ such that

$$f'(x) = \frac{f(5) - f(2)}{5 - 2}$$

And by assumption (on f') we have,

$$-2 \leq \frac{f(5) - 3}{3} \leq 1$$

$$\implies -3 \leq f(5) \leq 6$$

(a) $-3 \leq f(5) \leq 0$

(b) $-3 \leq f(5) \leq 6$

(c) $3 \leq f(5) \leq 6$

(d) $-5 \leq f(5) \leq 4$

(e) $-10 \leq f(5) \leq 10$

Name: _____

Instructor: _____

7.(6 pts.) Use linear approximation of $f(x) = \sqrt[3]{x}$ at $a = -8$ to estimate $\sqrt[3]{-8.12}$.

SOLUTION:

Using the linear approximation formula, we see that

$$L(x) = f(a) + f'(a)(x - a)$$

The derivative of f is $f'(x) = \frac{1}{3x^{2/3}}$. We see that $f(-8) = -2$ and

$$\begin{aligned} f'(-8) &= \frac{1}{3} \frac{1}{(-2)^2} \\ &= \frac{1}{12} \end{aligned}$$

So

$$L(x) = -2 + \frac{1}{12}(x + 8)$$

Thus, the approximation is

$$\begin{aligned} L(-8.12) &= -2 + \frac{1}{12}(-8.12 + 8.12) \\ &= -2 + -.01 \\ &= -2.01 \end{aligned}$$

- (a) -1.99 (b) -2.04 (c) **-2.01** (d) -1.8 (e) -2.2

Name: _____

Instructor: _____

8.(6 pts.) Find the linearization of the function $f(x) = \sin^2(x)$ at $a = \frac{\pi}{4}$.

SOLUTION:

Recall that the "Linearization" of a function at a certain point a is just its tangent line at the given point; and it is given by

$$L(x) = f'(a)(x - a) + f(a).$$

Since $f'(x) = 2(\sin(x)) \cos(x)$ we get $f'(a) = f'(\frac{\pi}{4}) = 2(\sin(\frac{\pi}{4})) \cos(\frac{\pi}{4}) = 2 \cdot \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{2}}{2} =$

1. Also $f(a) = f(\frac{\pi}{4}) = \left(\sin(\frac{\pi}{4})\right)^2 = \left(\frac{\sqrt{2}}{2}\right)^2 = \frac{1}{2}$. Hence,

$$L(x) = x - \frac{\pi}{4} + \frac{1}{2}.$$

(a) $x + \frac{1}{\sqrt{2}} - \frac{\pi}{4}$

(b) $\frac{x}{2} + \frac{1}{2} - \frac{\pi}{4}$

(c) $-\frac{10}{4}x - \frac{1}{4}$

(d) $\frac{1}{2}x + \frac{3}{2}$

(e) $x + \frac{1}{2} - \frac{\pi}{4}$

Name: _____

Instructor: _____

9.(6 pts.) Which of the following gives a complete list of the critical numbers/points of the function

$$f(x) = 3x^{2/3} \cdot (x + 1)^3$$

SOLUTION:

To be a critical number, you must either be a number a in the domain of f such that $f(a) = 0$ OR a number a in the domain of f such that f' does not exist at a . We observe that f is defined for every real number. Its derivative is given by

$$\begin{aligned} f'(x) &= \frac{2(x+1)^3}{x^{1/3}} + 9x^{2/3}(x+1)^2 \\ &= \frac{2(x+1)^3 + 9x(x+1)^2}{x^{1/3}} \end{aligned}$$

We see that f' is not defined at $a = 0$ and 0 is in the domain of f . So 0 is a critical number. Now, to find the rest of the critical values, we set $f' = 0$. We see this reduces to the equation

$$2(x+1)^3 + 9x(x+1)^2 = 0$$

This gives

$$(x+1)^2(2(x+1) + 9x) = 0$$

So $x = -1$ is a solution and solving

$$2(x+1) + 9x = 11x + 2 = 0$$

tells us that $x = -2/11$ is another solution. Thus, the full set of critical numbers is $x = 0, -2/11, -1$.

(a) $x = \frac{-2}{11}, -1$

(b) $x = \frac{2}{11}, -1$

(c) $x = 0, \frac{-2}{11}, -1$


(d) $x = \frac{2}{11}, 1$

(e) $x = -1, 0$

Name: _____

Instructor: _____

10.(6 pts.) A girl flies a kite at a height of 4 meters, the wind is carrying the kite horizontally away from her at a rate of 2 m/sec (there is no vertical movement in the kite). How fast must she let out the string when the kite is 5 meters away from her?



Solution: Let $l(t)$ be the hypotenuse of the triangle above. Our goal is to find $\frac{dl}{dt}$ when

$l(t) = 5$. Let $h(t)$ be the horizontal distance. Then implicitly differentiate

$$h(t)^2 + 16 = l(t)^2$$

to get

$$2h \frac{dh}{dt} = 2l \frac{dl}{dt}.$$

Then $\frac{dl}{dt} = \frac{2h}{l}$ since $\frac{dh}{dt} = 2$. If $l = 5$, then $16^2 + h^2 = 25$, so $h = 3$. Hence, when $l = 5$, $\frac{dl}{dt} = \frac{6}{5}$, so the answer is (a).

(a) $\frac{6}{5}$ m/sec

(b) $\frac{10}{3}$ m/sec

(c) $10\sqrt{3}$ m/sec

(d) 3 m/sec

(e) 5 m/sec

Name: _____

Instructor: _____

11.(6 pts.) Let $f(x) = 4x^2 - 4x + 4$. Find the absolute maximum and absolute minimum of f on the interval $[0, 2]$. (That is find the maximum and minimum value of $f(x)$ on the given interval).

SOLUTION:

First we compute the derivative, $f'(x) = 8x - 4$. Setting $f'(x) = 0 = 8x - 4$, we see that $x = 1/2$ is the only critical number (note that $f'(x)$ is defined for every real number). The absolute maximum and absolute minimum must occur at a critical number or at an endpoint of the interval. So we simply evaluate f at each of these points. We see that

$$f(0) = 4$$

$$f(2) = 4 \cdot 4 - 4 \cdot 2 + 4 = 12$$

and

$$f(1/2) = 4(1/4) - 4(1/2) + 4 = 3$$

From our computations, we see that f has max value 12 and min value 3.

- (a) Max value = 6, Min value = 4
- (b) Max value = 12, Min value = 3
- (c) Max value = 4, Min value = 3
- (d) Max value = 12, No Minimum value exists
- (e) Max value = 12, Min value = 4

Name: _____

Instructor: _____

12.(6 pts.) Consider the function $f(x) = x^4 - 8x^3 + 5$. Which of the following statements is true?

SOLUTION:

We begin by first computing the derivative and second derivative. We compute

$$\begin{aligned}f'(x) &= 4x^3 - 24x^2 \\ &= 4x^2(x - 6)\end{aligned}$$

and

$$\begin{aligned}f''(x) &= 12x^2 - 48x \\ &= 12x(x - 4)\end{aligned}$$

Now, we see that the zeros of $f''(x)$ are 0, 4. When $x < 0$, we see that $f''(x)$ is positive by choosing any test point such as -1 . We see that

$$\begin{aligned}f''(-1) &= 12(-1)(-1 - 4) \\ &= 60 \\ &> 0\end{aligned}$$

When $0 < x < 4$, we see that $f''(x)$ is negative by choosing a test point such as 1. We see that

$$\begin{aligned}f''(1) &= 12(-3) \\ &= -36 \\ &< 0\end{aligned}$$

When $x > 4$, we see that $f''(x)$ is positive by choosing a test point such as 5. We see that

$$\begin{aligned}f''(5) &= 5(12 \cdot 5 - 48) \\ &= 5 \cdot 12 \\ &= 60 \\ &> 0\end{aligned}$$

Our computations show that f'' changes signs at $x = 0$ and $x = 4$. This tells us that 0 and 4 are both points of inflection. Now, the zeros of the derivative (i.e, the critical numbers of f) are 0 and 6. We see that $f''(6) > 0$ and so the second derivative test tells us that at $x = 6$, there is a local minimum. By choosing test points such as -1 and 1, we see that $f'(-1) = -20 < 0$ and $f'(1) = -28 < 0$. Thus, at the point $x = 0$, there is neither a local minimum nor a local maximum. From all that we have shown, we see that a must be the correct solution. Note that this solution is given for completeness but we did not have to do all this work to arrive at the correct answer. For example, once we

Name: _____

Instructor: _____

saw that the only points of inflection occurred at $x = 0$ and $x = 4$, we could eliminate three of the answer choices. Thus, we were left to only decide what happened at $x = 0$.

- (a) f has a local minimum at $x = 6$, a local maximum at $x = 0$, and points of inflection at $x = 0$ and 4 .
- (b) f has a local minimum at $x = 6$, no local maximum, and points of inflection at $x = 0, 4$ and -4 .
- (c) f has local minima at $x = 0$ and 6 , no local maximum, and a point of inflection at $x = 4$.
- (d) f has a local minimum at $x = 6$, no local maximum, and points of inflection at $x = 0$ and 4 .
- (e) f has a local maximum at $x = 0$, no local minimum, and a point of inflection at $x = 4$.


Name: _____

Instructor: _____

Partial Credit

You must show your work on the partial credit problems to receive credit!

13.(11 pts.) Pedestrian A is walking towards the intersection C of two streets intersecting at a right angle. Pedestrian B is walking away from intersection C . Pedestrian A is going North at 2 mph, and Pedestrian B is going East at 3 mph. How fast is the distance from Pedestrian A to Pedestrian B changing when Pedestrian A is 4 miles South of intersection C , and Pedestrian B is 3 miles East of intersection C .



SOLUTION:

Let us denote by $z(t)$ the distance at any time between the pedestrians, by $y(t)$ the distance between A and the intersection and by $x(t)$ the distance between B and the intersection. We are given,

$$\frac{dy}{dt} = -2 \quad ; \quad \frac{dx}{dt} = 3.$$

And we want to know

$$\left. \frac{dz}{dt} \right|_{y=4, x=3}.$$

We know by the Pythagorean Theorem that $z^2(t) = y^2(t) + x^2(t)$ and differentiating with respect to time in both sides gives

$$\begin{aligned} 2z(t) \frac{dz}{dt} &= 2y(t) \frac{dy}{dt} + 2x(t) \frac{dx}{dt} \\ \implies \frac{dz}{dt} &= \frac{1}{z(t)} \left[y(t) \frac{dy}{dt} + x(t) \frac{dx}{dt} \right]. \end{aligned}$$

We also know (again by the Pythagorean theorem) that if $y = 4$ and $x = 3$, then $z = 5$, so that in mph

$$\left. \frac{dz}{dt} \right|_{y=4, x=3} = \frac{1}{5} [4(-2) + 3(3)] = \frac{1}{5}$$

Name: _____

Instructor: _____

14.(13 pts.) Consider the function

$$f(x) = x^3 + x - \frac{1}{x},$$

with domain $(0, \infty)$. With this restriction on the domain, show that the equation $f(x) = 0$ has one and exactly one real solution for $x \in (0, \infty)$. Identify the theorem(s) you are using and show the validity of the required conditions to apply the theorems are true to gain full credit.

SOLUTION:

We first note that f is continuous on $(0, \infty)$. Thus, we may attempt to use the Intermediate Value Theorem. Now, we see that $f(1/2) = 1/8 + 1/2 - 2 = 5/8 - 2 < 0$. Also, we see that $f(2) = 8 + 2 - 1/2 = 10 - 1/2 > 0$. Thus, f changes signs between $1/2$ and 2 . Since f is also continuous on $(1/2, 2)$, the Intermediate Value Theorem tells us that there is a root of f somewhere in $(1/2, 2)$.

To justify that there is at most one solution, we invoke the Mean Value Theorem or Rolle's Theorem. If there were more than one root of f , then there must be at least two. Call them a and b and say $b < a$. As f is continuous and differentiable on all of $(0, \infty)$, we have that f must be continuous on $[b, a]$ and differentiable on (b, a) . The Mean Value Theorem or Rolle's Theorem both imply that there is some c in (b, a) such that

$$\begin{aligned} f'(c) &= \frac{f(b) - f(a)}{b - a} \\ &= 0, \end{aligned}$$

Now $f'(x) = 3x^2 + 1 + \frac{1}{x^2}$. We see that $f'(x) > 0$ for all values of x in $(0, \infty)$, since $f'(x) = 1 + (\text{sum of squares}) \geq 1$, since a sum of squares is always non-negative. Since the derivative is always strictly greater than 0 on $(0, \infty)$ and so in particular on (b, a) . So we have a contradiction.

Thus the assumption that there is more than one solution must be false.

Name: _____

Instructor: _____

15.(16 pts.) Let $f(x) = x - \sin(2x)$, with domain $(-\frac{\pi}{2}, \frac{\pi}{2})$.

(a) Find the critical numbers/points for f .

SOLUTION:

Since f' is continuous we only look for when does $f'(x) = 1 - 2\cos(2x) = 0$. This happens when $\cos(2x) = \frac{1}{2}$, which implies that $2x = \pm\frac{\pi}{3} \implies x = \pm\frac{\pi}{6}$.

(b) Find the intervals where f is increasing and decreasing. (justify your answer)

SOLUTION:

Knowing that f has critical points at $\pm\frac{\pi}{6}$ we select points from the three new formed intervals $(-\frac{\pi}{2}, -\frac{\pi}{6})$, $(-\frac{\pi}{6}, \frac{\pi}{6})$ and $(\frac{\pi}{6}, \frac{\pi}{2})$, say $-\frac{\pi}{4}$, 0 and $\frac{\pi}{4}$ respectively. We see that $f'(\pm\frac{\pi}{4}) = 1 > 0$ and that $f'(0) = 1 - 2 = -1 < 0$. So that by the first derivative test f is increasing on $(-\frac{\pi}{2}, -\frac{\pi}{6}) \cup (\frac{\pi}{6}, \frac{\pi}{2})$ and decreasing on $(-\frac{\pi}{6}, \frac{\pi}{6})$.

(c) Classify the critical points as local maxima or local minima or neither and justify your conclusions in each case.

SOLUTION:

By what we just found in part (b) and the first derivative test, we know that f has a local maximum at $-\frac{\pi}{6}$ and a local minimum at $\frac{\pi}{6}$. Moreover the second derivative test gives $f''(-\frac{\pi}{6}) = 4\sin(-\frac{\pi}{3}) < 0$ implies it is a local maximum while $f''(\frac{\pi}{6}) = 4\sin(\frac{\pi}{3}) > 0$ implies it is a local minimum.

Name: _____

Instructor: _____

ROUGH WORK

Name: _____

Instructor: ANSWERS

Math 10550, EXAM II
October 13, 3016

- The Honor Code is in effect for this examination. All work is to be your own.
- No calculators.
- The exam lasts for 1 hr. and 15 min.
- Be sure that your name is on every page in case pages become detached.
- Be sure that you have all 18 pages of the test.
- Each multiple choice question is worth 7 points. Your score will be the sum of the best 10 scores on the multiple choice questions plus your score on questions 13-15.

PLEASE MARK YOUR ANSWERS WITH AN X, not a circle!				
1.	(a)	(b)	(c)	(d) (●)
2.	(a)	(●)	(c)	(d) (e)
.....				
3.	(●)	(b)	(c)	(d) (e)
4.	(a)	(b)	(●)	(d) (e)
.....				
5.	(a)	(b)	(c)	(d) (●)
6.	(a)	(●)	(c)	(d) (e)
.....				
7.	(a)	(b)	(●)	(d) (e)
8.	(a)	(b)	(c)	(d) (●)
.....				
9.	(a)	(b)	(●)	(d) (e)
10.	(●)	(b)	(c)	(d) (e)
.....				
11.	(a)	(●)	(c)	(d) (e)
12.	(a)	(b)	(c)	(●) (e)

Please do NOT write in this box.

Multiple Choice _____

13. _____

14. _____

15. _____

Total _____