

## Lecture 2 : The Natural Logarithm.

Recall


$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad n \neq -1.$$

What happens if  $n = -1$ ?

**Definition** We can define a function which is an **anti-derivative for  $x^{-1}$**  using the Fundamental Theorem of Calculus: We let

$$\ln x = \int_1^x \frac{1}{t} dt, \quad x > 0.$$


This function is called the natural logarithm.



**Note** that  $\ln(x)$  is the area under the continuous curve  $y = \frac{1}{t}$  between 1 and  $x$  if  $x > 1$  and minus the area under the continuous curve  $y = \frac{1}{t}$  between 1 and  $x$  if  $x < 1$ .


We have  $\ln(2)$  is the area of the region shown in the picture on the left above and  $\ln(1/2)$  is minus the area of the region shown in the picture on the right above.

I do not have a formula for  $\ln(x)$  in terms of functions studied before, however I could estimate the value of  $\ln(2)$  using a Riemann sum. The approximating rectangles for a left Riemann sum with 10 approximating rectangles is shown below. Their area adds to 0.718771 ( to 6 decimal places). If we took the limit of such sums as the number of approximating rectangles tends to infinity, we would get the actual value of  $\ln(2)$ , which is 0.693147 ( to 6 decimal places). The natural logarithm function is a built in function on most scientific calculators.



With very little work, using a right Riemann sum with 1 approximating rectangle, we can get a lower bound for  $\ln(2)$ . The picture below demonstrates that

The picture below demonstrates that  $\ln 2 = \int_1^2 \frac{1}{t} dt > 1/2$ .



### Properties of the Natural Logarithm:

We can use our tools from Calculus I to derive a lot of information about the natural logarithm.

1. Domain =  $(0, \infty)$  (by definition)
2. Range =  $(-\infty, \infty)$  (see later)
3.  $\ln x > 0$  if  $x > 1$ ,  $\ln x = 0$  if  $x = 1$ ,  $\ln x < 0$  if  $x < 1$ .

This follows from our comments above after the definition about how  $\ln(x)$  relates to the area under the curve  $y = 1/x$  between 1 and  $x$ .

4.  $\frac{d(\ln x)}{dx} = \frac{1}{x}$

This follows from the definition and the Fundamental Theorem of Calculus.

5. The graph of  $y = \ln x$  is increasing, continuous and concave down on the interval  $(0, \infty)$ .

Let  $f(x) = \ln(x)$ ,  $f'(x) = 1/x$  which is always positive for  $x > 0$  (the domain of  $f$ ), Therefore the graph of  $f(x)$  is increasing on its domain. We have  $f''(x) = \frac{-1}{x^2}$  which is always negative, showing that the graph of  $f(x)$  is concave down. The function  $f$  is continuous since it is differentiable.



6. The function  $f(x) = \ln x$  is a one-to-one function.

Since  $f'(x) = 1/x$  which is positive on the domain of  $f$ , we can conclude that  $f$  is a one-to-one function.

7. Since  $f(x) = \ln x$  is a one-to-one function, there is a unique number,  $e$ , with the property that

$$\boxed{\ln e = 1.}$$

We have  $\ln(1) = 0$  since  $\int_1^1 1/t dt = 0$ . Using a Riemann sum with 3 approximating rectangles, we see that  $\ln(4) > 1/1 + 1/2 + 1/3 > 1$ . Therefore by the intermediate value theorem, since  $f(x) = \ln(x)$  is continuous, there must be some number  $e$  with  $1 < e < 4$  for which  $\ln(e) = 1$ . This number is unique since the function  $f(x) = \ln(x)$  is one-to-one.



We will be able to estimate the value of  $e$  in the next section with a limit.  $e \approx 2.7182818284590$ .

The following properties are very useful when calculating with the natural logarithm:

(i)  $\ln 1 = 0$

(ii)  $\ln(ab) = \ln a + \ln b$

(iii)  $\ln\left(\frac{a}{b}\right) = \ln a - \ln b$

(iv)  $\ln a^r = r \ln a$

where  $a$  and  $b$  are positive numbers and  $r$  is a rational number.

**Proof** (ii) We show that  $\ln(ax) = \ln a + \ln x$  for a constant  $a > 0$  and any value of  $x > 0$ . The rule follows with  $x = b$ . Let  $f(x) = \ln x$ ,  $x > 0$  and  $g(x) = \ln(ax)$ ,  $x > 0$ . We have  $f'(x) = \frac{1}{x}$  and  $g'(x) = \frac{1}{ax} \cdot a = \frac{1}{x}$ .

Since both functions have equal derivatives,  $f(x) + C = g(x)$  for some constant  $C$ . Substituting  $x = 1$  in this equation, we get  $\ln 1 + C = \ln a$ , giving us  $C = \ln a$  and  $\ln ax = \ln a + \ln x$ .

(iii) Note that  $0 = \ln 1 = \ln \frac{a}{a} = \ln a + \ln \frac{1}{a}$ , giving us that  $\ln \frac{1}{a} = -\ln a$ .

Thus we get  $\ln \frac{a}{b} = \ln a + \ln \frac{1}{b} = \ln a - \ln b$ .

(iv) Comparing derivatives, we see that

$$\frac{d(\ln x^r)}{dx} = \frac{rx^{r-1}}{x^r} = \frac{r}{x} = \frac{d(r \ln x)}{dx}.$$

Hence  $\ln x^r = r \ln x + C$  for any  $x > 0$  and any rational number  $r$ . Letting  $x = 1$  we get  $C = 0$  and the result holds.

**Example** Expand

$$\ln \frac{x^2 \sqrt{x^2 + 1}}{x^3}$$

using the rules of logarithms.

**Example** Express as a single logarithm:

$$\ln x + 3 \ln(x + 1) - \frac{1}{2} \ln(x + 1).$$

**Example** Evaluate  $\int_1^{e^2} \frac{1}{t} dt$

---


We can use the rules of logarithms given above to derive the following information about limits.

$$\boxed{\lim_{x \rightarrow \infty} \ln x = \infty, \quad \lim_{x \rightarrow 0} \ln x = -\infty.}$$

**Proof** We saw above that  $\ln 2 > 1/2$ . If  $x > 2^n$ , then  $\ln x > \ln 2^n$  (Why ?). So  $\ln x > n \ln 2 > n/2$ . Hence as  $x \rightarrow \infty$ , the values of  $\ln x$  also approach  $\infty$ .

Also  $\ln \frac{1}{2^n} = -n \ln 2 < -n/2$ . Thus as  $x$  approaches 0 the values of  $\ln x$  approach  $-\infty$ .

Note that we can now draw a reasonable sketch of the graph of  $y = \ln(x)$ , using all of the information derived above.




**Example** Find the limit  $\lim_{x \rightarrow \infty} \ln\left(\frac{1}{x^2+1}\right)$ .

---

We can extend the applications of the natural logarithm function by composing it with the absolute value function. We have :

$$\ln |x| = \begin{cases} \ln x & x > 0 \\ \ln(-x) & x < 0 \end{cases}$$

This is an even function with graph



We have  $\ln|x|$  is also an antiderivative of  $1/x$  with a larger domain than  $\ln(x)$ .

$$\boxed{\frac{d}{dx}(\ln|x|) = \frac{1}{x}} \quad \text{and} \quad \boxed{\int \frac{1}{x} dx = \ln|x| + C}$$

We can use the chain rule and integration by substitution to get

$$\boxed{\frac{d}{dx}(\ln|g(x)|) = \frac{g'(x)}{g(x)}} \quad \text{and} \quad \boxed{\int \frac{g'(x)}{g(x)} dx = \ln|g(x)| + C}$$

**Example** Differentiate  $\ln|\sqrt[3]{x-1}|$ .

**Example** Find the integral

$$\int \frac{x}{3-x^2} dx.$$

### Logarithmic Differentiation

To differentiate  $y = f(x)$ , it is often easier to use logarithmic differentiation :

1. Take the natural logarithm of both sides to get  $\ln y = \ln(f(x))$ .
2. Differentiate with respect to  $x$  to get  $\frac{1}{y} \frac{dy}{dx} = \frac{d}{dx} \ln(f(x))$
3. We get  $\frac{dy}{dx} = y \frac{d}{dx} \ln(f(x)) = f(x) \frac{d}{dx} \ln(f(x))$ .

**Example** Find the derivative of  $y = \sqrt[4]{\frac{x^2+1}{x^2-1}}$ .

## Extra Examples

Please try to work through these questions before looking at the solutions.

**Example** Expand  $\ln\left(\frac{e^2\sqrt{a^2+1}}{b^3}\right)$

**Example** Differentiate  $\ln|\sqrt[3]{x-1}|$ .

**Example** Find  $d/dx \ln(|\cos x|)$ .

**Example** Find the integral

$$\int \cot x dx$$

**Example** Find the integral

$$\int_e^{e^2} \frac{1}{x \ln x} dx.$$

**Example** Find the derivative of  $y = \frac{\sin^2 x \tan^4 x}{(x^2-1)^2}$ .

**Old Exam Question** Differentiate the function

$$f(x) = \frac{(x^2-1)^4}{\sqrt{x^2+1}}.$$

## Solutions

**Example** Expand  $\ln\left(\frac{e^2\sqrt{a^2+1}}{b^3}\right)$

$$\begin{aligned}\ln\left(\frac{e^2\sqrt{a^2+1}}{b^3}\right) &= \ln(e^2\sqrt{a^2+1}) - \ln(b^3) = \ln(e^2) + \ln(\sqrt{a^2+1}) - 3\ln b \\ &= 2\ln e + \frac{1}{2}\ln(a^2+1) - 3\ln b = 2 + \frac{1}{2}\ln(a^2+1) - 3\ln b.\end{aligned}$$

**Example** Differentiate  $\ln|\sqrt[3]{x-1}|$ .

We use the chain rule here

$$\frac{d}{dx} \ln|\sqrt[3]{x-1}| = \frac{1}{\sqrt[3]{x-1}} \cdot \frac{1}{3}(x-1)^{-2/3} = \frac{1}{3(x-1)}.$$

**Example** Find  $d/dx \ln(|\cos x|)$ .

Again, we use the chain rule

$$\frac{d}{dx} \ln|\cos x| = \frac{1}{\cos x} \cdot (-\sin x) = -\tan x.$$

**Example** Find the integral

$$\int \cot x dx$$

$$\int \cot x dx = \int \frac{\cos x}{\sin x} dx.$$

We use substitution. Let  $u = \sin x$ ,  $du = \cos x dx$ .

$$\int \frac{\cos x}{\sin x} dx = \int \frac{du}{u} = \ln|u| + C = \ln|\sin x| + C.$$

**Example** Find the integral

$$\int_e^{e^2} \frac{1}{x \ln x} dx.$$

We use substitution. Let  $u = \ln x$ ,  $du = \frac{1}{x} dx$ .  $u(e) = \ln e = 1$ ,  $u(e^2) = \ln e^2 = 2$ .

$$\int_e^{e^2} \frac{1}{x \ln x} dx = \int_1^2 \frac{du}{u} = \ln|u| \Big|_1^2 = \ln 2 - \ln 1 = \ln 2.$$

**Example** Find the derivative of  $y = \frac{\sin^2 x \tan^4 x}{(x^2-1)^2}$ .

We use Logarithmic differentiation. If  $y = \frac{\sin^2 x \tan^4 x}{(x^2-1)^2}$ , then

$$\ln y = \ln(\sin^2 x) + \ln(\tan^4 x) - \ln((x^2 - 1)^2) = 2 \ln(\sin x) + 4 \ln(\tan x) - 2 \ln(x^2 - 1).$$

Differentiating both sides with respect to  $x$ , we get

$$\frac{1}{y} \frac{dy}{dx} = \frac{2 \cos x}{\sin x} + \frac{4 \sec^2 x}{\tan x} - \frac{2(2x)}{x^2 - 1}.$$

Multiplying both sides by  $y$  and converting to a function of  $x$ , we get

$$\frac{dy}{dx} = y \left[ \frac{2 \cos x}{\sin x} + \frac{4 \sec^2 x}{\tan x} - \frac{4x}{x^2 - 1} \right] = \left( \frac{\sin^2 x \tan^4 x}{(x^2 - 1)^2} \right) \left[ \frac{2 \cos x}{\sin x} + \frac{4 \sec^2 x}{\tan x} - \frac{4x}{x^2 - 1} \right].$$

**Old Exam Question** Differentiate the function

$$f(x) = \frac{(x^2 - 1)^4}{\sqrt{x^2 + 1}}.$$

We use Logarithmic differentiation. If  $y = \frac{(x^2-1)^4}{\sqrt{x^2+1}}$ , then

$$\ln y = 4 \ln(x^2 - 1) - \frac{1}{2} \ln(x^2 + 1).$$

Differentiating both sides with respect to  $x$ , we get

$$\frac{1}{y} \frac{dy}{dx} = \frac{4(2x)}{x^2 - 1} - \frac{2x}{2(x^2 + 1)}.$$

Multiplying both sides by  $y$  and converting to a function of  $x$ , we get

$$\frac{dy}{dx} = y \left[ \frac{8x}{x^2 - 1} - \frac{x}{(x^2 + 1)} \right] = \left( \frac{(x^2 - 1)^4}{\sqrt{x^2 + 1}} \right) \left[ \frac{8x}{x^2 - 1} - \frac{x}{(x^2 + 1)} \right].$$