

# Trigonometric Substitution

To solve integrals containing the following expressions;

$$\sqrt{a^2 - x^2} \quad \sqrt{x^2 + a^2} \quad \sqrt{x^2 - a^2},$$

it is sometimes useful to make the following substitutions:

Expression	Substitution	Identity
$\sqrt{a^2 - x^2}$	$x = a \sin \theta, \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ or $\theta = \sin^{-1} \frac{x}{a}$	$1 - \sin^2 \theta = \cos^2 \theta$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta, \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ or $\theta = \tan^{-1} \frac{x}{a}$	$1 + \tan^2 \theta = \sec^2 \theta$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta, \quad 0 \leq \theta < \frac{\pi}{2}$ or $\pi \leq \theta < \frac{3\pi}{2}$ or $\theta = \sec^{-1} \frac{x}{a}$	$\sec^2 \theta - 1 = \tan^2 \theta$

**Note** The calculations here are much easier if you use the substitution in reverse:  $x = a \sin \theta$  as opposed to  $\theta = \sin^{-1} \frac{x}{a}$ .

# Integrals involving $\sqrt{a^2 - x^2}$

We make the substitution  $x = a \sin \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \cos \theta d\theta$ ,  
 $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = a |\cos \theta| = a \cos \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice. )

## Example

$$\int \frac{x^3}{\sqrt{4 - x^2}} dx$$

# Integrals involving $\sqrt{a^2 - x^2}$

We make the substitution  $x = a \sin \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \cos \theta d\theta$ ,  
 $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = a |\cos \theta| = a \cos \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice. )

## Example

$$\int \frac{x^3}{\sqrt{4 - x^2}} dx$$

- ▶ Let  $x = 2 \sin \theta$ ,  $dx = 2 \cos \theta d\theta$ ,  $\sqrt{4 - x^2} = \sqrt{4 - 4 \sin^2 \theta} = 2 \cos \theta$ .

# Integrals involving $\sqrt{a^2 - x^2}$

We make the substitution  $x = a \sin \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \cos \theta d\theta$ ,  
 $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = a |\cos \theta| = a \cos \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice.)

## Example

$$\int \frac{x^3}{\sqrt{4 - x^2}} dx$$

- ▶ Let  $x = 2 \sin \theta$ ,  $dx = 2 \cos \theta d\theta$ ,  $\sqrt{4 - x^2} = \sqrt{4 - 4 \sin^2 \theta} = 2 \cos \theta$ .
- ▶  $\int \frac{x^3 dx}{\sqrt{4 - x^2}} = \int \frac{8 \sin^3 \theta (2 \cos \theta d\theta)}{2 \cos \theta} = \int 8 \sin^3 \theta d\theta = \int 8 \sin^2 \theta \sin \theta d\theta = 8 \int (1 - \cos^2 \theta) \sin \theta d\theta$ .

# Integrals involving $\sqrt{a^2 - x^2}$

We make the substitution  $x = a \sin \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \cos \theta d\theta$ ,  
 $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = a |\cos \theta| = a \cos \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice.)

## Example

$$\int \frac{x^3}{\sqrt{4-x^2}} dx$$

- ▶ Let  $x = 2 \sin \theta$ ,  $dx = 2 \cos \theta d\theta$ ,  $\sqrt{4-x^2} = \sqrt{4-4\sin^2 \theta} = 2 \cos \theta$ .
- ▶  $\int \frac{x^3 dx}{\sqrt{4-x^2}} = \int \frac{8 \sin^3 \theta (2 \cos \theta d\theta)}{2 \cos \theta} = \int 8 \sin^3 \theta d\theta = \int 8 \sin^2 \theta \sin \theta d\theta = 8 \int (1 - \cos^2 \theta) \sin \theta d\theta$ .
- ▶ Let  $w = \cos \theta$ ,  $dw = -\sin \theta d\theta$ ,

$$\begin{aligned} 8 \int (1 - \cos^2 \theta) \sin \theta d\theta &= -8 \int (1 - w^2) dw = 8 \int (w^2 - 1) dw = \frac{8w^3}{3} - 8w + C \\ &= \frac{8(\cos \theta)^3}{3} - 8 \cos \theta + C = \frac{8(\cos(\sin^{-1} \frac{x}{2}))^3}{3} - 8 \cos(\sin^{-1} \frac{x}{2}) + C. \end{aligned}$$


Integrals involving  $\sqrt{a^2 - x^2}$ 

$$\int \frac{x^3}{\sqrt{4-x^2}} dx = \frac{8(\cos(\sin^{-1} \frac{x}{2}))^3}{3} - 8 \cos(\sin^{-1} \frac{x}{2}) + C.$$

# Integrals involving $\sqrt{a^2 - x^2}$

$$\int \frac{x^3}{\sqrt{4-x^2}} dx = \frac{8(\cos(\sin^{-1} \frac{x}{2}))^3}{3} - 8 \cos(\sin^{-1} \frac{x}{2}) + C.$$

- To get an expression for  $\cos(\sin^{-1} \frac{x}{2})$ , we use an appropriate triangle




From the triangle, we get

$$\cos(\sin^{-1} \frac{x}{2}) = \frac{\sqrt{4-x^2}}{2}$$

# Integrals involving $\sqrt{a^2 - x^2}$

$$\int \frac{x^3}{\sqrt{4-x^2}} dx = \frac{8(\cos(\sin^{-1} \frac{x}{2}))^3}{3} - 8 \cos(\sin^{-1} \frac{x}{2}) + C.$$

- To get an expression for  $\cos(\sin^{-1} \frac{x}{2})$ , we use an appropriate triangle



From the triangle, we get  
 $\cos(\sin^{-1} \frac{x}{2}) = \frac{\sqrt{4-x^2}}{2}$

- hence  $\int \frac{x^3 dx}{\sqrt{4-x^2}} = \frac{8\left(\frac{\sqrt{4-x^2}}{2}\right)^3}{3} - 8\frac{\sqrt{4-x^2}}{2} + C = \frac{(4-x^2)^{3/2}}{3} - 4\sqrt{4-x^2} + C$



# Integrals involving $\sqrt{a^2 - x^2}$

**Example**  $\int \frac{dx}{x^2\sqrt{9-x^2}}$

# Integrals involving $\sqrt{a^2 - x^2}$

**Example**  $\int \frac{dx}{x^2\sqrt{9-x^2}}$

▶ Let  $x = 3 \sin \theta$ ,  $dx = 3 \cos \theta d\theta$ ,  $\sqrt{9 - x^2} = \sqrt{9 - 9 \sin^2 \theta} = 3 \cos \theta$ .

# Integrals involving $\sqrt{a^2 - x^2}$

**Example**  $\int \frac{dx}{x^2\sqrt{9-x^2}}$


▶ Let  $x = 3 \sin \theta$ ,  $dx = 3 \cos \theta d\theta$ ,  $\sqrt{9 - x^2} = \sqrt{9 - 9 \sin^2 \theta} = 3 \cos \theta$ .

▶  $\int \frac{dx}{x^2\sqrt{9-x^2}} = \int \frac{3 \cos \theta d\theta}{(9 \sin^2 \theta)3 \cos \theta} = \int \frac{1}{9 \sin^2 \theta} d\theta = \frac{-\cot \theta}{9} + C = \frac{-\cot(\sin^{-1} \frac{x}{3})}{9} + C$

# Integrals involving $\sqrt{a^2 - x^2}$

**Example**  $\int \frac{dx}{x^2 \sqrt{9-x^2}}$

- ▶ Let  $x = 3 \sin \theta$ ,  $dx = 3 \cos \theta d\theta$ ,  $\sqrt{9-x^2} = \sqrt{9-9\sin^2 \theta} = 3 \cos \theta$ .
- ▶  $\int \frac{dx}{x^2 \sqrt{9-x^2}} = \int \frac{3 \cos \theta d\theta}{(9 \sin^2 \theta) 3 \cos \theta} = \int \frac{1}{9 \sin^2 \theta} d\theta = \frac{-\cot \theta}{9} + C = \frac{-\cot(\sin^{-1} \frac{x}{3})}{9} + C$
- ▶ To get an expression for  $\cot(\sin^{-1} \frac{x}{3})$ , we use an appropriate triangle




From the triangle, we get  
 $\cot(\sin^{-1} \frac{x}{3}) = \frac{\sqrt{9-x^2}}{x}$  and hence

$$\int \frac{dx}{x^2 \sqrt{9-x^2}} = \frac{-\sqrt{9-x^2}}{9x} + C$$

# Integrals involving $\sqrt{a^2 - x^2}$

**Example**  $\int \frac{dx}{x^2 \sqrt{9-x^2}}$

- ▶ Let  $x = 3 \sin \theta$ ,  $dx = 3 \cos \theta d\theta$ ,  $\sqrt{9-x^2} = \sqrt{9-9\sin^2 \theta} = 3 \cos \theta$ .
- ▶  $\int \frac{dx}{x^2 \sqrt{9-x^2}} = \int \frac{3 \cos \theta d\theta}{(9 \sin^2 \theta) 3 \cos \theta} = \int \frac{1}{9 \sin^2 \theta} d\theta = \frac{-\cot \theta}{9} + C = \frac{-\cot(\sin^{-1} \frac{x}{3})}{9} + C$
- ▶ To get an expression for  $\cot(\sin^{-1} \frac{x}{3})$ , we use an appropriate triangle



From the triangle, we get  
 $\cot(\sin^{-1} \frac{x}{3}) = \frac{\sqrt{9-x^2}}{x}$  and hence

$$\int \frac{dx}{x^2 \sqrt{9-x^2}} = \frac{-\sqrt{9-x^2}}{9x} + C$$

- ▶ **Note** You can also use this method to derive what you already know

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} + C$$

# Integrals involving $\sqrt{x^2 + a^2}$

We make the substitution  $x = a \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + a^2} = \sqrt{a^2 \tan^2 \theta + a^2} = a |\sec \theta| = a \sec \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice.)

## Example

$$\int \frac{dx}{\sqrt{x^2 + 4}}$$

# Integrals involving $\sqrt{x^2 + a^2}$

We make the substitution  $x = a \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + a^2} = \sqrt{a^2 \tan^2 \theta + a^2} = a |\sec \theta| = a \sec \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice.)

## Example

$$\int \frac{dx}{\sqrt{x^2 + 4}}$$

- Let  $x = 2 \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = 2 \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + 4} = \sqrt{4 \tan^2 \theta + 4} = 2 \sqrt{\sec^2 \theta} = 2 \sec \theta$ .

# Integrals involving $\sqrt{x^2 + a^2}$

We make the substitution  $x = a \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + a^2} = \sqrt{a^2 \tan^2 \theta + a^2} = a |\sec \theta| = a \sec \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice.)

## Example

$$\int \frac{dx}{\sqrt{x^2 + 4}}$$

- ▶ Let  $x = 2 \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = 2 \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + 4} = \sqrt{4 \tan^2 \theta + 4} = 2 \sqrt{\sec^2 \theta} = 2 \sec \theta$ .
- ▶  $\int \frac{dx}{\sqrt{x^2 + 4}} = \int \frac{2 \sec^2 \theta d\theta}{2 \sec \theta} = \int \sec \theta d\theta = \ln |\sec \theta + \tan \theta| + C =$   
 $\ln \left| \sec\left(\tan^{-1} \frac{x}{2}\right) + \tan\left(\tan^{-1} \frac{x}{2}\right) \right| + C = \ln \left| \sec\left(\tan^{-1} \frac{x}{2}\right) + \frac{x}{2} \right| + C$




# Integrals involving $\sqrt{x^2 + a^2}$

We make the substitution  $x = a \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + a^2} = \sqrt{a^2 \tan^2 \theta + a^2} = a |\sec \theta| = a \sec \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice.)

## Example

$$\int \frac{dx}{\sqrt{x^2 + 4}}$$

- ▶ Let  $x = 2 \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = 2 \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + 4} = \sqrt{4 \tan^2 \theta + 4} = 2\sqrt{\sec^2 \theta} = 2 \sec \theta$ .
- ▶  $\int \frac{dx}{\sqrt{x^2 + 4}} = \int \frac{2 \sec^2 \theta d\theta}{2 \sec \theta} = \int \sec \theta d\theta = \ln |\sec \theta + \tan \theta| + C =$   
 $\ln \left| \sec\left(\tan^{-1} \frac{x}{2}\right) + \tan\left(\tan^{-1} \frac{x}{2}\right) \right| + C = \ln \left| \sec\left(\tan^{-1} \frac{x}{2}\right) + \frac{x}{2} \right| + C$
- ▶ To get an expression for  $\sec\left(\tan^{-1} \frac{x}{2}\right)$ , we use an appropriate triangle



From the triangle, we get  
 $\sec\left(\tan^{-1} \frac{x}{2}\right) = \frac{\sqrt{x^2 + 4}}{2}$  and hence

$$\int \frac{dx}{\sqrt{x^2 + 4}} = \ln \left| \frac{\sqrt{x^2 + 4}}{2} + \frac{x}{2} \right| + C.$$

# Integrals involving $\sqrt{x^2 + a^2}$

We make the substitution  $x = a \tan \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $dx = a \sec^2 \theta d\theta$ ,  
 $\sqrt{x^2 + a^2} = \sqrt{a^2 \tan^2 \theta + a^2} = a |\sec \theta| = a \sec \theta$  (since  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  by choice. )

**Note** You can also use this substitution to get the familiar

$$\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + C.$$

# Integrals involving $\sqrt{x^2 + a^2}$ , Completing the square.

Sometimes we can convert an integral to a form where trigonometric substitution can be applied by completing the square.

**Example** Evaluate

$$\int \frac{dx}{\sqrt{x^2 - 4x + 13}}.$$

# Integrals involving $\sqrt{x^2 + a^2}$ , Completing the square.

Sometimes we can convert an integral to a form where trigonometric substitution can be applied by completing the square.

**Example** Evaluate

$$\int \frac{dx}{\sqrt{x^2 - 4x + 13}}.$$

▶  $x^2 - 4x + 13 = x^2 - 2(2)x + 2^2 - 2^2 + 13 = (x - 2)^2 + 9.$

# Integrals involving $\sqrt{x^2 + a^2}$ , Completing the square.

Sometimes we can convert an integral to a form where trigonometric substitution can be applied by completing the square.

**Example** Evaluate

$$\int \frac{dx}{\sqrt{x^2 - 4x + 13}}.$$

- ▶  $x^2 - 4x + 13 = x^2 - 2(2)x + 2^2 - 2^2 + 13 = (x - 2)^2 + 9.$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 4x + 13}} = \int \frac{dx}{\sqrt{(x-2)^2 + 9}} = \int \frac{du}{\sqrt{(u)^2 + 9}},$  where  $u = x - 2.$

# Integrals involving $\sqrt{x^2 + a^2}$ , Completing the square.

Sometimes we can convert an integral to a form where trigonometric substitution can be applied by completing the square.

**Example** Evaluate

$$\int \frac{dx}{\sqrt{x^2 - 4x + 13}}.$$

- ▶  $x^2 - 4x + 13 = x^2 - 2(2)x + 2^2 - 2^2 + 13 = (x - 2)^2 + 9.$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 4x + 13}} = \int \frac{dx}{\sqrt{(x-2)^2 + 9}} = \int \frac{du}{\sqrt{(u)^2 + 9}},$  where  $u = x - 2.$
- ▶ *Now we apply the substitution  $u = 3 \tan \theta,$   $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2},$   
 $du = 3 \sec^2 \theta d\theta,$   $\sqrt{u^2 + 9} = \sqrt{9 \tan^2 \theta + 9} = 3\sqrt{\sec^2 \theta} = 3 \sec \theta.$*

# Integrals involving $\sqrt{x^2 + a^2}$ , Completing the square.

Sometimes we can convert an integral to a form where trigonometric substitution can be applied by completing the square.

**Example** Evaluate

$$\int \frac{dx}{\sqrt{x^2 - 4x + 13}}.$$

- ▶  $x^2 - 4x + 13 = x^2 - 2(2)x + 2^2 - 2^2 + 13 = (x - 2)^2 + 9.$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 4x + 13}} = \int \frac{dx}{\sqrt{(x-2)^2 + 9}} = \int \frac{du}{\sqrt{(u)^2 + 9}},$  where  $u = x - 2.$
- ▶ *Now we apply the substitution  $u = 3 \tan \theta,$   $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2},$   
 $du = 3 \sec^2 \theta d\theta,$   $\sqrt{u^2 + 9} = \sqrt{9 \tan^2 \theta + 9} = 3\sqrt{\sec^2 \theta} = 3 \sec \theta.$*
- ▶  $\int \frac{du}{\sqrt{(u)^2 + 9}} = \int \frac{3 \sec^2 \theta d\theta}{3 \sec \theta} = \int \sec \theta d\theta = \ln |\sec \theta + \tan \theta| + C =$   
 $\ln \left| \sec(\tan^{-1} \frac{u}{3}) + \tan(\tan^{-1} \frac{u}{3}) \right| + C.$


# Integrals involving $\sqrt{x^2 + a^2}$ , Completing the square.

►  $\int \frac{dx}{\sqrt{x^2 - 4x + 13}} = \ln \left| \sec\left(\tan^{-1} \frac{u}{3}\right) + \tan\left(\tan^{-1} \frac{u}{3}\right) \right| + C, \text{ where } u = x - 2.$



# Integrals involving $\sqrt{x^2 + a^2}$ , Completing the square.

- ▶  $\int \frac{dx}{\sqrt{x^2 - 4x + 13}} = \ln \left| \sec(\tan^{-1} \frac{u}{3}) + \tan(\tan^{-1} \frac{u}{3}) \right| + C$ , where  $u = x - 2$ .
- ▶ To get an expression for  $\sec(\tan^{-1} \frac{u}{3})$ , we use an appropriate triangle




From the triangle, we get

$$\sec(\tan^{-1} \frac{u}{3}) = \frac{\sqrt{u^2 + 9}}{3}$$

Integrals involving  $\sqrt{x^2 + a^2}$ , Completing the square.

- ▶  $\int \frac{dx}{\sqrt{x^2 - 4x + 13}} = \ln \left| \sec(\tan^{-1} \frac{u}{3}) + \tan(\tan^{-1} \frac{u}{3}) \right| + C$ , where  $u = x - 2$ .
- ▶ To get an expression for  $\sec(\tan^{-1} \frac{u}{3})$ , we use an appropriate triangle



From the triangle, we get

$$\sec(\tan^{-1} \frac{u}{3}) = \frac{\sqrt{u^2 + 9}}{3}$$

- ▶  $\int \frac{dx}{\sqrt{x^2 - 4x + 13}} = \ln \left| \frac{\sqrt{u^2 + 9}}{3} + \frac{u}{3} \right| + C = \ln \left| \frac{\sqrt{(x-2)^2 + 9}}{3} + \frac{(x-2)}{3} \right| + C$ .

# Integrals involving $\sqrt{x^2 - a^2}$

We make the substitution  $x = a \sec \theta$ ,  $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ , ( This amounts to saying  $\theta = \sec^{-1} \frac{x}{a}$  ),  $dx = a \sec \theta \tan \theta d\theta$ ,  
 $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = a |\tan \theta| = a \tan \theta$  (since  $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$  by choice)

**Example** Evaluate

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx$$

# Integrals involving $\sqrt{x^2 - a^2}$

We make the substitution  $x = a \sec \theta$ ,  $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ , ( This amounts to saying  $\theta = \sec^{-1} \frac{x}{a}$  ),  $dx = a \sec \theta \tan \theta d\theta$ ,  
 $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = a |\tan \theta| = a \tan \theta$  (since  $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$  by choice)

**Example** Evaluate

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx$$

- Let  $x = 5 \sec \theta$   $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ , then  $dx = 5 \sec \theta \tan \theta d\theta$ ,  
 $\sqrt{x^2 - 25} = \sqrt{25 \sec^2 \theta - 25} = 5 \sqrt{\tan^2 \theta} = 5 |\tan \theta| = 5 \tan \theta$ .

# Integrals involving $\sqrt{x^2 - a^2}$

We make the substitution  $x = a \sec \theta$ ,  $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ , ( This amounts to saying  $\theta = \sec^{-1} \frac{x}{a}$  ),  $dx = a \sec \theta \tan \theta d\theta$ ,  
 $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = a |\tan \theta| = a \tan \theta$  (since  $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$  by choice)

**Example** Evaluate

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx$$

▶ Let  $x = 5 \sec \theta$   $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ , then  $dx = 5 \sec \theta \tan \theta d\theta$ ,  
 $\sqrt{x^2 - 25} = \sqrt{25 \sec^2 \theta - 25} = 5 \sqrt{\tan^2 \theta} = 5 |\tan \theta| = 5 \tan \theta$ .

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx = \int \frac{5 \sec \theta \tan \theta}{25 \sec^2 \theta (5 \tan \theta)} d\theta = \int \frac{1}{25 \sec \theta} d\theta$$

$$= \frac{1}{25} \int \cos \theta d\theta = \frac{1}{25} \sin \theta + C = \frac{1}{25} \sin(\sec^{-1} \frac{x}{5}) + C$$


# Integrals involving $\sqrt{x^2 - a^2}$

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx = \frac{1}{25} \sin(\sec^{-1} \frac{x}{5}) + C$$

# Integrals involving $\sqrt{x^2 - a^2}$

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx = \frac{1}{25} \sin(\sec^{-1} \frac{x}{5}) + C$$

- To get an expression for  $\sin(\sec^{-1} \frac{x}{5})$ , we use an appropriate triangle




From the triangle, we get

$$\sin(\sec^{-1} \frac{x}{5}) = \frac{\sqrt{x^2 - 25}}{x}$$

# Integrals involving $\sqrt{x^2 - a^2}$

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx = \frac{1}{25} \sin(\sec^{-1} \frac{x}{5}) + C$$

- To get an expression for  $\sin(\sec^{-1} \frac{x}{5})$ , we use an appropriate triangle



From the triangle, we get

$$\sin(\sec^{-1} \frac{x}{5}) = \frac{\sqrt{x^2 - 25}}{x}$$


- Hence  $\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx = \frac{1}{25} \cdot \frac{\sqrt{x^2 - 25}}{x} + C = \frac{\sqrt{x^2 - 25}}{25x} + C$ .



# Integrals involving $\sqrt{x^2 - a^2}$

$$\int \frac{1}{x^2\sqrt{x^2 - 25}} dx = \frac{1}{25} \sin(\sec^{-1} \frac{x}{5}) + C$$

- To get an expression for  $\sin(\sec^{-1} \frac{x}{5})$ , we use an appropriate triangle



From the triangle, we get

$$\sin(\sec^{-1} \frac{x}{5}) = \frac{\sqrt{x^2 - 25}}{x}$$

- Hence  $\int \frac{1}{x^2\sqrt{x^2 - 25}} dx = \frac{1}{25} \cdot \frac{\sqrt{x^2 - 25}}{x} + C = \frac{\sqrt{x^2 - 25}}{25x} + C$ .
- **Note** You can also use this substitution to get

$$\int \frac{1}{x\sqrt{x^2 - a^2}} dx = \frac{1}{a} \sec^{-1} \frac{x}{a} + C.$$

# Integrals involving $\sqrt{x^2 - a^2}$ , Completing the square

**Example** Evaluate  $\int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}}$ .

# Integrals involving $\sqrt{x^2 - a^2}$ , Completing the square

**Example** Evaluate  $\int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}}$ .

- ▶ *Completing the square*  $x^2 - 6x + 8 = x^2 - 2(3)x + 9 - 9 + 8 = (x - 3)^2 - 1$

# Integrals involving $\sqrt{x^2 - a^2}$ , Completing the square

**Example** Evaluate  $\int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}}$ .

- ▶ *Completing the square*  $x^2 - 6x + 8 = x^2 - 2(3)x + 9 - 9 + 8 = (x - 3)^2 - 1$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 6x + 8}} dx = \int \frac{dx}{\sqrt{(x-3)^2 - 1}} dx = \int \frac{du}{\sqrt{u^2 - 1}}$ , where  $u = x - 3$ .

# Integrals involving $\sqrt{x^2 - a^2}$ , Completing the square

**Example** Evaluate  $\int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}}$ .

- ▶ *Completing the square*  $x^2 - 6x + 8 = x^2 - 2(3)x + 9 - 9 + 8 = (x - 3)^2 - 1$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 6x + 8}} = \int \frac{dx}{\sqrt{(x-3)^2 - 1}} = \int \frac{du}{\sqrt{u^2 - 1}}$ , where  $u = x - 3$ .
- ▶ Let  $u = \sec \theta$   $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ ,  $du = \sec \theta \tan \theta d\theta$ .

# Integrals involving $\sqrt{x^2 - a^2}$ , Completing the square

**Example** Evaluate  $\int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}}$ .

- ▶ *Completing the square*  $x^2 - 6x + 8 = x^2 - 2(3)x + 9 - 9 + 8 = (x - 3)^2 - 1$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 6x + 8}} = \int \frac{dx}{\sqrt{(x-3)^2 - 1}} = \int \frac{du}{\sqrt{u^2 - 1}}$ , where  $u = x - 3$ .
- ▶ Let  $u = \sec \theta$   $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ ,  $du = \sec \theta \tan \theta d\theta$ .
- ▶  $= \int \frac{\sec \theta \tan \theta d\theta}{\sqrt{\sec^2 \theta - 1}} = \int \frac{\sec \theta \tan \theta d\theta}{\sqrt{\sec^2 \theta - 1}} = \int \sec \theta d\theta$

# Integrals involving $\sqrt{x^2 - a^2}$ , Completing the square


**Example** Evaluate  $\int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}}$ .

- ▶ *Completing the square*  $x^2 - 6x + 8 = x^2 - 2(3)x + 9 - 9 + 8 = (x - 3)^2 - 1$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 6x + 8}} = \int \frac{dx}{\sqrt{(x-3)^2 - 1}} = \int \frac{du}{\sqrt{u^2 - 1}}$ , where  $u = x - 3$ .
- ▶ Let  $u = \sec \theta$   $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ ,  $du = \sec \theta \tan \theta d\theta$ .
- ▶  $= \int \frac{\sec \theta \tan \theta d\theta}{\sqrt{\sec^2 \theta - 1}} = \int \frac{\sec \theta \tan \theta d\theta}{\sqrt{\sec^2 \theta - 1}} = \int \sec \theta d\theta$
- ▶  $= \ln |\sec \theta + \tan \theta| + C = \ln |\sec(\sec^{-1} u) + \tan(\sec^{-1} u)|$

# Integrals involving $\sqrt{x^2 - a^2}$ , Completing the square

**Example** Evaluate  $\int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}}$ .

- ▶ *Completing the square*  $x^2 - 6x + 8 = x^2 - 2(3)x + 9 - 9 + 8 = (x - 3)^2 - 1$
- ▶  $\int \frac{dx}{\sqrt{x^2 - 6x + 8}} = \int \frac{dx}{\sqrt{(x-3)^2 - 1}} = \int \frac{du}{\sqrt{u^2 - 1}}$ , where  $u = x - 3$ .
- ▶ Let  $u = \sec \theta$   $0 \leq \theta < \frac{\pi}{2}$  or  $\pi \leq \theta < \frac{3\pi}{2}$ ,  $du = \sec \theta \tan \theta d\theta$ .
- ▶  $= \int \frac{\sec \theta \tan \theta d\theta}{\sqrt{\sec^2 \theta - 1}} = \int \frac{\sec \theta \tan \theta d\theta}{\sqrt{\sec^2 \theta - 1}} = \int \sec \theta d\theta$
- ▶  $= \ln |\sec \theta + \tan \theta| + C = \ln |\sec(\sec^{-1} u) + \tan(\sec^{-1} u)|$
- ▶ *To get an expression for  $\tan(\sec^{-1} u)$ , we use an appropriate triangle*



*From the triangle, we get*

$\tan(\sec^{-1} u) = \sqrt{u^2 - 1}$  Hence

$$\begin{aligned} & \int \frac{dx}{\sqrt{x^2 - 6x + 8}} \\ &= \ln |u + \sqrt{u^2 - 1}| + C \\ &= \ln |(x - 3) + \sqrt{(x - 3)^2 - 1}| + C. \\ & \int_4^6 \frac{dx}{\sqrt{x^2 - 6x + 8}} \\ &= \ln |3 + \sqrt{8}| - \ln |1| = \ln |3 + \sqrt{8}| \end{aligned}$$