

The Picard-Vessiot theory, constrained cohomology, and linear differential algebraic groups

Anand Pillay*
University of Notre Dame

November 6, 2016

Abstract

We prove that a differential field K is algebraically closed and Picard-Vessiot closed if and only if the differential Galois cohomology group, $H_{\partial}^1(K, G)$, is trivial for any linear differential algebraic group G over K . We give applications to the parameterized Picard-Vessiot theory and pose several problems.

1 Introduction and preliminaries

In this paper we are mainly concerned with differential fields (K, ∂) of characteristic 0 and we try to relate two “theories”, the Galois theory of linear differential equations over K (i.e. the Picard-Vessiot theory), and Kolchin’s “constrained cohomology” (which we prefer to call “differential Galois cohomology”) of linear differential algebraic groups over K . We work in the category of differential algebraic varieties in the sense of Kolchin [12]. Because we are concerned with differential algebraic groups and their torsors, we may equally well (and in fact do) work in the category of definable sets in the first order theory DCF_0 of differentially closed fields (with respect to

*Research supported by NSF grant DMS-1360702, as well as NSF DMS-1606334 and NSA number H98230-15-1-0245

a single derivation ∂) of characteristic 0. The main result is the following, where we explain the notions later.

Theorem 1.1. *Let (K, ∂) be a differential field. Then K is algebraically closed and Picard-Vessiot closed if and only if $H_{\partial}^1(K, G) = \{1\}$ for any linear differential algebraic group G over K , namely if every differential algebraic principal homogeneous space X for G , defined over K , has a K -point.*

Remark 1.2. (i) *A Picard-Vessiot (or PV) extension L of K is by definition a differential field extension L of K generated over K by a fundamental system of solutions Y_1, \dots, Y_n of a linear differential system $\partial Y = AY$ over K , such that $C_L = C_K$, namely L has “no new constants”. So formally, to be PV-closed means that K has no proper PV extension L .*

(ii) *When C_K is algebraically closed (in particular when K is algebraically closed), then there always exists a PV extension of K for any linear differential system $\partial Y = AY$, hence to be Picard-Vessiot closed is equivalent to K containing a fundamental system of solutions for any linear system $\partial Y = AY$ over K .*

(iii) *When C_K is algebraically closed then K being PV-closed implies K algebraically closed.*

(iv) *For G a differential algebraic group over K , $H_{\partial}^1(K, G)$ is defined in [12], Chapter VII, in terms of certain cocycles from $\mathcal{G} = \text{Aut}(K^{\text{diff}}/K)$ to $G(K^{\text{diff}})$, where K^{diff} is the differential closure of K inside some “universal” differentially closed field \mathcal{U} . $H_{\partial}^1(K, G)$ parametrizes the set of differential algebraic principal homogeneous spaces X for G over K up to isomorphism over K (in the appropriate category). So triviality of $H_{\partial}^1(K, G)$ means that for every such PHS X for G over K , $X(K) \neq \emptyset$.*

We will also point out how our proof of Theorem 1.1 extends to prove the following:

Theorem 1.3. *Let K be a differential field. Then K is algebraically closed and closed under generalized strongly normal extensions if and only if $H_{\partial}^1(K, G) = \{1\}$ for any (not necessarily linear) differential algebraic group G over K .*

Theorem 1.3 can be seen as evidence for the naturalness of the generalized strongly normal theory.

Theorem 1.1 has the following consequence for the *parameterized Picard-Vessiot theory* [7], where again the additional notions will be explained subsequently.

Corollary 1.4. *Let K be a $\{\partial_x, \partial_t\}$ -differential field where ∂_x and ∂_t are commuting derivations. Let K^{∂_x} denote the field of constants of K with respect to ∂_x . Let $\partial_x Y = AY$ be a parametrized linear differential equation over K . Assume that K^{∂_x} is both algebraically closed and PV-closed as a ∂_t -field. Then K has a unique parameterized Picard Vessiot extension for the equation.*

Although our results are stated in purely differential algebraic terms, we will freely use a mixture of differential algebraic and model-theoretic notations and methods in the proofs. In fact, differential algebraic geometry in the sense of Kolchin is synonymous with the model theory of differentially closed fields, in terms of content and subject matter, although model theory brings with it a certain generality in point of view.

This paper is concerned mainly with ordinary differential fields of characteristic 0, namely fields K of characteristic 0 equipped with a single derivation, although in Corollary 1.4 and section 3 we touch on the case of a set Δ of commuting derivations. For the majority of the introduction we will focus on $\Delta = \{\partial\}$. For differential algebraic geometry, namely the theory (and category) of differential algebraic varieties defined over a differential field K , the reader could consult [12], although some of the axiomatic presentation is a bit obscure. But Section 1.1 of [8] can also serve as an introduction to this material. The reader is also referred to Buium's book [3] for an approach somewhat more informed by modern algebraic geometry. Key objects of this paper are differential algebraic groups G and their principal homogeneous spaces (or torsors) X . Namely G is a differential algebraic group defined over K , X is a differential algebraic variety defined over K , and we have a regular (strictly transitive) group action of G on X (on the left) which is a morphism $G \times X \rightarrow X$ over K defined over K in the sense of differential algebraic varieties.

DCF_0 is the first order theory of differentially closed fields of characteristic 0 in the language $L = \{+, -, 0, 1, \times, \partial\}$. It is complete, ω -stable, with quantifier elimination, and elimination of imaginaries. We take \mathcal{U} to be a "saturated" model of DCF_0 in which all differential fields we are concerned with are assumed to embed. \mathcal{U} is precisely a "universal" differential field in the sense of Kolchin. K will usually denote a "small" differential subfield of \mathcal{U} . \mathcal{C} denotes the field of constants of \mathcal{U} , and C_K the field of constants of K . But when we work with partial fields (several commuting derivations)

notation will be a bit different. The reader is referred to [14] for the basic model theory of DCF_0 . By a K -definable set we mean a subset of \mathcal{U}^n which is definable with parameters from K in the L -structure \mathcal{U} . Our paper [19] gives an exhaustive account of Kolchin's theory of differential algebraic varieties and groups, and its relation to the category of definable sets and definable groups, and the interested reader is referred there. We summarize:

Fact 1.5. (i) any group G definable over K has a unique structure of a differential algebraic group over K .

(ii) If G is a definable group over K and X is a principal homogeneous space for G over K , in the category of definable sets over K , then X can be given (uniquely) the structure of a differential algebraic variety over K such that X is a PHS for G over K in the category of differential algebraic varieties over K ,

(iii) If (G, X) is a definable (differential algebraic) PHS over K then there is an embedding (over K) in an algebraic PHS (G_1, X_1) over K .

Also connectedness of groups has the same meaning in the definable and differential algebraic categories.

An important notion in this paper is the *differential closure* K^{diff} of a differential field K . It is the analogue in differential algebra of the algebraic closure K^{alg} of a field K , and from the point of view of model theory is the “prime model over K ”. Now $Aut(K^{alg}/K)$ is a profinite group, but $Aut(K^{diff}/K)$ does not have such a nice “geometric” structure. As in the case of usual Galois cohomology, the set of PHS's over K for a given differential algebraic group G over K can be described in terms of suitable cocycles from $Aut(K^{diff}/K)$ to $G(K^{diff})$, modulo the equivalence relation of being cohomologous. This is put into a somewhat more general model-theoretic environment in [20], where it is pointed out that *definable* cocycles suffice. For the purposes of the current paper, these descriptions are not needed. In any case $H^1_{\partial}(K, G)$ denotes the set of (differential algebraic) PHS's for G , over K up to isomorphism over K , where the isomorphism should respect the G -action. We will make use of Kolchin's theorem (Theorem 4, Chapter VII, [12]) which says:

Fact 1.6. Suppose G is an algebraic group over K . Then $H^1_{\partial}(K, G) = H^1(K, G)$. In particular if K is algebraically closed then $H^1(K, G) = \{1\}$.

Definition 1.7. (i) A linear differential algebraic group (LDAG) over K is a subgroup of $GL_n(\mathcal{U})$ which is differential algebraic over K , equivalently

definable over K in \mathcal{U} .

(ii) An Abelian differential algebraic group over K is a differential algebraic (equivalently definable) over K subgroup of $A(\mathcal{U})$ for some Abelian variety A over K .

Due to Fact 1.5 (iii) and the Chevalley decomposition for algebraic groups we have:

Fact 1.8. *Let G be a connected differential algebraic group over K . Then we have an exact sequence $1 \rightarrow C \rightarrow G \rightarrow B \rightarrow 1$ of connected differential algebraic groups over K where C is linear and B is Abelian.*

We will make use of the following result from [4]:

Fact 1.9. *If G is a LDAG over K and N is a normal definable (over K) subgroup, then G/N is an LDAG over K .*

The analogous result for Abelian DAG's fails.

Model theory provides various ordinal-valued dimensions attached to definable sets and types in DCF_0 , the most important for our purposes being Morley rank RM and U -rank. In DCF_0 every complete type $p(x)$ (in finitely many variables) has U -rank $< \omega^2$. The U -rank of a definable set is the supremum of the U -ranks of complete types extending that definable set. For definable groups G in DCF_0 , $RM(G) = U(G)$ and moreover the supremum above is attained by the so-called "generic types" of G . (See [22]). We will call G finite-dimensional if $RM(G) < \omega$. This is Buium's nomenclature [3] and corresponds to G having "differential type 0" in the notation of [8] for example. In general the Morley rank and U -rank of a definable group G will have the form $\omega \cdot m + k$, for m, k nonnegative integers. The "leading coefficient" (which is k when $m = 0$ and m otherwise) is related to but distinct from the "typical differential dimension" in the Kolchin theory.

We will be making use of a number of results about U -rank and definable groups from [23] (some of which come from the seminal paper [1]) with full references.

The model theoretic environment for partial differential algebraic geometry is the theory $DCF_{0,m}$ of differentially closed fields with respect to m commuting derivations $\partial_1, \dots, \partial_m$. Essentially everything we have said above

about the ordinary case passes over to the partial case, except now the U -ranks (and Morley ranks) of types are bounded by ω^{m+1} , namely every complete type has the form $\omega^m k_m + \omega^{m-1} k_{m-1} + \dots + \omega k_1 + k_0$, where k_0, \dots, k_m are nonnegative integers.

Finally we discuss the Galois theoretic notions which are central to this paper. We are mainly concerned with the Picard-Vessiot theory, but we mention “generalized strongly normal extensions” in Theorem 1.3, and the parameterized Picard-Vessiot (*PPV*) theory, with respect to $DCF_{0,2}$ in Corollary 1.4.

We repeat part of Remark 1.2. A linear differential equation over K , in vector form, is something of the form

$$(*) \partial Y = AY$$

where Y is a $n \times 1$ (column) vector and A is an $n \times n$ matrix over K . The set V of solutions in \mathcal{U} is a subset of \mathcal{U}^n which is an n -dimensional \mathcal{C} -vector space. What we call a fundamental system of solutions is precisely a \mathcal{C} -basis of V , and is the same thing as a matrix $Z \in GL_n(\mathcal{U})$ whose columns are solutions of (*). A *PV* extension of K for (*) is a differential field extension L of the form $K\langle Z \rangle$ for some fundamental system Z of solutions, such that $C_K = C_L$. L is always contained in some differential closure K^{diff} of K , and when C_K is algebraically closed, such a *PV* extension L exists and is unique.

A first generalization of the notion of Picard-Vessiot extension is that of a *strongly normal extension* and is due to Kolchin [11]. L is a strongly normal extension of K if for some α such that L is generated over K by α , we have (i) for any realization β of $tp(\alpha/K)$, β is in the differential field generated by K , α and \mathcal{C} , and (ii) $C_K = C_L$.

A further generalization is the notion of a “generalized strongly normal extension of K ” which was introduced in [16] and elaborated on in [17], [15] and [18]. Let X be a K -definable set (such as the field of constants). $L = K\langle \alpha \rangle$ is an X -strongly normal extension of K if (i) for any realization β of $tp(\alpha/K)$, β is in the differential field generated by α , K , and (the coordinates of elements of) X , and (ii) $X(K) = X(K^{diff})$. (We should mention that condition (ii) has been weakened in [13], although this does not affect the current paper.)

When K is algebraically closed, generalized strongly normal extensions (as well as strongly normal extensions) can be explicitly described in terms of solving certain kinds of differential equations, as follows: Given a connected

algebraic group over K , a ∂ -structure on G is an extension of the derivation ∂ on K to a derivation D of the structure sheaf (coordinate ring when G is affine) of G . Such a derivation D is equivalent to a homomorphic regular section s over K of a certain “shifted tangent bundle” $T_\partial G$ of G . We call (G, s) an algebraic ∂ -group over K , which is an object belonging to algebraic geometry (and was introduced and studied in [3]). It gives rise to a map (in fact crossed homomorphism) $dlog_{(G,s)}$ belonging to differential algebraic geometry, from G to its Lie algebra LG , namely the map taking $y \in G(\mathcal{U})$ to $\partial(y)s(y)^{-1}$ where the latter is computed in the algebraic group $T_\partial(G)$. We let $(G, s)^\partial$ be the kernel of $dlog_{(G,s)}$, a finite-dimensional differential algebraic group over K , and if $a \in LG(K)$, we call the equation $dlog_{(G,s)}(y) = a$ a logarithmic differential equation over K on (G, s) . In any case, it is shown in [18] that (assuming K to be algebraically closed), L is a generalized strongly normal extension of K (i.e. an X -strongly normal extension of K for some X) iff there is a connected algebraic ∂ -group (G, s) over K , a logarithmic differential equation $dlog_{(G,s)}(-) = a \in LG(K)$ over K and a solution $y \in G(K^{diff})$ of the equation such that (i) $L = K(y)$ and (ii) $(G, s)^\partial(K) = (G, s)^\partial(K^{diff})$. The case when G is over C_K and s is the “trivial” ∂ -structure on G recovers Kolchin’s strongly normal theory.

2 Proofs of Main results

We use notation and conventions from the previous section. We use repeatedly the fact that $C_{K^{diff}} = C_K^{alg}$, in particular if C_K is algebraically closed then it coincides with the constants of K^{diff} .

Proof of Theorem 1.1.

We first point out the (trivial) right to left direction. If K is not algebraically closed, then a proper Galois extension of K gives a finite principal homogeneous space X over K for a finite group G , such that $X(K) = \emptyset$. So we may assume K is algebraically closed. If K is not PV -closed K has a proper PV -extension for a system $\partial Y = AY$ over K . L is generated by a fundamental system $b = (b_1, \dots, b_n)$ of solutions for the equation. The map taking $\sigma \in Aut_\partial(L/K)$ to $\sigma(b)b^{-1}$ (multiplication in GL_n) does not depend on the choice of b , and establishes an isomorphism between $Aut_\partial(L/K)$ and a differential algebraic subgroup G of GL_n defined over K . The orbit of b under G (where G acts on the left) is a differential algebraic left PHS for G , defined

over K , and with no K -point, contradicting our assumptions. (The above construction of G should be considered well-known but details are also given in Section 3 of [10].)

We now prove the left to right direction. We are assuming that K is algebraically closed and PV closed. We let G be an arbitrary linear differential algebraic group over K and we aim to prove that $H_{\partial}^1(K, G) = \{1\}$.

We will be using the following:

Lemma 2.1 (INDUCTIVE PRINCIPLE). *Suppose that we have a short exact sequence $1 \rightarrow N \rightarrow G \rightarrow H \rightarrow 1$ of differential algebraic groups over K . Suppose that $H_{\partial}^1(K, H) = \{1\}$ and $H_{\partial}^1(K, N) = \{1\}$. Then $H_{\partial}^1(K, G) = \{1\}$.*

Proof. This follows from the usual exact sequence in cohomology, but we can be more explicit: Suppose X is a differential algebraic PHS for G over K . Then the set Y of N -orbits in X is a differential algebraic PHS for H over K . If Y has a K -point, then this yields an N -orbit $Z \subseteq X$, defined over K , and then a K -point of Z yields a K -point of X . \square

There are several ways to proceed, modulo Lemma 2.1 above, depending on what and how much to quote of the differential algebraic and model-theoretic literature. Cassidy and Singer [8], working in the context of several commuting derivations, prove that an arbitrary “strongly connected” differential algebraic group over K has a subnormal series (of differential algebraic groups over K) where the successive quotients are “almost simple”. Moreover in the ordinary case (one derivation) they describe the “almost simple” linear differential algebraic groups, so it would be enough to check Theorem 1.1 for each of these. In fact we will take a slightly different route, with more model-theoretic input, notation, and information about the structure of linear differential algebraic groups, but ending up checking Theorem 1.1 in essentially the same special cases (and more or less recovering the decomposition theorem of [8]).

Case (1) G is “finite-dimensional” (so finite Morley rank, or finite U -rank). By the inductive principle we may assume G is connected. Let N be the “definable solvable radical of G ”, namely the maximal normal solvable connected definable subgroup. It exists and is unique as G has finite Morley rank, in particular N is defined over K . Then clearly $H = G/N$ is defined over K , linear (by Fact 1.9), and *semisimple* in the sense that H has no

connected normal abelian definable subgroup. This notion of semisimplicity agrees with the notion in [5].

By Lemma 2.1, we can separate into subcases:

Case (1)(a). $G = N$, namely G is solvable.

As G is a differential algebraic subgroup of $GL_n(\mathcal{U})$ defined over K , we can consider \bar{G} , the Zariski closure of G , a connected solvable linear algebraic group over K (in the algebraic geometric sense). By the results in Section III.10 of [2], \bar{G} is filtered by a chain of normal algebraic subgroups (over K) each successive quotient being isomorphic over K to the additive group \mathbb{G}_a or multiplicative group \mathbb{G}_m . Intersecting this chain with G , we see that G is filtered by a chain of normal definable (over K) subgroups, where we may assume that each successive quotient is a subgroup of $(\mathcal{U}, +)$ or (\mathcal{U}^*, \times) . By Lemma 2.1 we may assume that either G is a (finite-dimensional) differential algebraic subgroup of $(\mathcal{U}, +)$, or a (finite-dimensional) differential algebraic subgroup of (\mathcal{U}^*, \times) .

Suppose first $G \leq (\mathcal{U}, +)$. By Cassidy [6], G (being finite dimensional) is defined by a homogeneous linear differential equation over K (in a single indeterminate y but of arbitrary finite order). So $G(K^{diff})$ is a finite-dimensional vector space over C_K (the latter being algebraically closed) and adjoining to K a basis, yields a Picard-Vessiot extension of K . By our assumptions such a basis must already lie in K whereby $G(K^{diff})$ is isomorphic, over K to $(C_K)^d$ for some d . Suppose now that X is a *PHS* for G over K , so $X(K^{diff})$ is a *PHS* for $(C_K)^d$ over K . Let $\alpha \in X(K^{diff})$, so $L = K\langle\alpha\rangle$ is clearly a strongly normal extension of K . For $\sigma \in Aut_\partial(L/K)$ there is unique $g_\sigma \in (C_K)^d$ such that $\sigma(\alpha) = g_\sigma\alpha$. As $(C_K)^d$ is commutative we see that the map taking σ to g_σ yields an isomorphism between $Aut_\partial(L/K)$ and some algebraic subgroup of C_K^d , in particular with a linear algebraic group (in the constants), whereby, by Corollary 2 in Section 9, Chapter VI of [11], L is a *PV* extension of K . So by our assumptions, $\alpha \in X(K)$. We have shown that $H_\partial^1(K, G) = \{1\}$. (Alexey Ovchinnikov mentioned that we can also use Corollary 1 on p. 193 of [12] in this case.)

Now suppose $G \leq (\mathcal{U}^*, \times)$. We can use the logarithmic derivative map from (\mathcal{U}^*, \times) to $(\mathcal{U}, +)$ which takes y to $\partial(y)/y$ to map G onto a finite-dimensional definable (over K) subgroup of $(\mathcal{U}, +)$ with kernel a definable (over C_K) subgroup of $\mathbb{G}_m(C)$. We can apply the previous paragraph to the image of G . The kernel is handled in a similar fashion (any *PHS* for it over K will yield a strongly normal, hence *PV* extension of K , so has to be trivial). So Lemma 2.1 tells that $H_\partial^1(K, G)$ is again trivial.

We have completed Case (1)(a).

Case (1)(b). $G = H$, namely G is “semisimple”. So G is a linear (connected) ∂ -semisimple group over K in the sense of [5]. Putting together Theorems 14 and 15 of [5], G is isogenous over K to a direct product of simple (up to finite centres) noncommutative finite-dimensional linear differential algebraic groups G_1, \dots, G_k , each over K . By Lemma 2.1, we may assume that G is equal to some G_i and is moreover centreless. By Theorem 1.5 of [21] $G(K^{diff})$ is isomorphic over K^{diff} to a group of the form $H(C_{K^{diff}}) = H(C_K)$ where H is a simple linear algebraic group over C_K . By Proposition 4.1 of [18] and our assumptions, such an isomorphism can be found defined over K . So we may assume $G = H(\mathcal{C})$. Now suppose X is a (left) PHS for $H(\mathcal{C})$, over K . Let again $\alpha \in X(K^{diff})$ and $L = K\langle\alpha\rangle$. The map taking $\sigma \in Aut_{\partial}(L/K)$ to g_{σ} in $H(C_K)$ establishes an isomorphism between $Aut_{\partial}(L/K)$ and B^{opp} for some algebraic subgroup B of $H(C_K)$. As B^{opp} is also a linear algebraic group in C_K , we see again by the result of Kolchin mentioned earlier, that L is a PV extension of K . So by our assumptions $\alpha \in X(K)$. So X has a K -point, and we have completed Case (1)(b) and also Case (1).

Case (2). G is arbitrary.

We will again use model theoretic notions although purely differential algebraic notions would also suffice. In general G has U -rank $\omega.m + k$ for some nonnegative integers, m, k and G is “finite dimensional” iff $m = 0$. We call G *1-connected* if G has no definable quotient G/H (a homogeneous space) with $U(G/H)$ finite (or G/H finite-dimensional). By the descending chain condition on definable subgroups (given by ω -stability) G has a unique 1-connected component, namely a smallest subgroup $G^{1,0}$ such that $G/G^{1,0}$ is finite-dimensional. Moreover if G is defined over K so is $G^{1,0}$. By Corollary 6.3 of [23] (U -rank inequalities for groups) and Theorem 6.7 of [23] (the Berline-Lascar decomposition), G is 1-connected iff it is connected and has U -rank of the form $\omega.m$. In any case by Case (1) and Lemma 2.1,

We may assume that G is 1-connected.

We let N be the maximal normal solvable 1-connected definable subgroup of G , clearly defined over K , which exists and is unique for the same reasons as in Case (1). Then $H = G/N$ is also 1-connected, and is “almost semisimple” in the sense that H has no proper nontrivial normal 1-connected abelian definable subgroup. By Lemma 2.1 it is enough to treat the cases

where $G = N$ and where $G = H$. So:

Case (2)(a). $G = N$. Exactly as in Case (1)(a), by looking at the Zariski closure \bar{G} of G , a connected solvable linear algebraic group, we find a sequence G_i of normal K -definable subgroups of G with the quotients (without loss of generality) subgroups of $(\mathcal{U}, +)$ or (\mathcal{U}, \times) . By Case (1) we may assume that each quotient G_i/G_{i-1} has infinite Morley rank (or U -rank) and thus has Morley rank (or U -rank) ω . But both $(\mathcal{U}, +)$ and (\mathcal{U}^*, \times) are connected with U -rank ω whereby G_i/G_{i-1} is equal to $(\mathcal{U}, +)$ or (\mathcal{U}^*, \times) . In particular each quotient is an algebraic group over K . We can now use Fact 1.6 to deduce that $H_{\partial}^1(G)$ is trivial.

Case (2)(b). $G = H$. Let Z be the centre of G . By ‘‘almost semisimplicity’’ of G , Z is finite-dimensional. On the other hand using the 1-connectedness of G , $G/Z(G)$ is simple (and centreless). So by the inductive argument and Case (1), we may assume that G is centreless (and still linear). We may assume that the Zariski closure of G (in the ambient general linear group) is semisimple and even centreless (for example by Theorem 14 of [5]). By Theorem 15 of [5] we may assume that G is simple and its Zariski closure \bar{G} is a simple algebraic group over K . As G has infinite Morley rank, we can conclude from Theorem 17 of [5], or Proposition 5.1 of [21], that $G = \bar{G}$, namely G is an algebraic group. Again by Fact 1.6, $H_{\partial}^1(K, G)$ is trivial. The proof is complete.

Proof of Theorem 1.3. The nontrivial (left to right) direction of Theorem 1.3 is actually implicitly claimed in [17]: Corollary 3.6 in that paper proves that if the first order theory of K is superstable, G is a finite Morley rank group definable over K , and X is a K -definable PHS for G , then $X(K) = X(K^{diff})$. And the sentence following the proof of that result says that it is not hard to prove (assuming $Th(K)$ superstable) that $X(K) \neq \emptyset$ whenever X is a PHS over K for *any* definable group G over K , although no actual proof is given. As is readily seen the only use of superstability in the proof of Corollary 3.6 is Theorem 1.2 in that paper which says that superstability of $Th(K)$ implies that K is closed under generalized strongly normal extensions. As the proof of 3.6 in [17] makes heavy use of the full trichotomy theorem for U -rank 1-types in DCF_0 , we take the opportunity to sketch how to extend the rather direct proof of Theorem 1.1 above to a proof of Theorem 1.3.

First the right to left direction: If K is not algebraically closed then a Galois extension of K gives rise to a PHS over K for a finite group,

as before. So we assume K to be algebraically closed. By [18], described in the last paragraph of the introduction to the current paper, a generalized strongly normal extension L of K is given by adjoining a solution $\alpha \in G(K^{diff})$ of a logarithmic differential equation $dlog_{G,s}(-) = a$ on an algebraic D -group (G, s) over K with $a \in LG(K)$, where (G, s) is K -large, namely $(G, s)^\partial(K^{diff}) = (G, s)^\partial(K)$. As in the proof of Theorem 1.1 the map taking $\sigma \in Aut_\partial(L/K)$ to $\sigma(\alpha)\alpha^{-1}$ (multiplication etc. in G) established an isomorphism with a differential algebraic subgroup H^+ of $G(K^{diff})$, defined over K . Moreover the orbit of α under left multiplication by H^+ is a left PHS for H^+ with no K -point, contradicting the assumptions.

For the left to right direction we first use Fact 1.8 to obtain an exact sequence $1 \rightarrow N \rightarrow G \rightarrow H \rightarrow 1$ of differential algebraic groups over K , where N is linear (namely embeds over K into some GL_n) and H is “Abelian”, namely embeds over K in an abelian variety. Hence the proof proceeds exactly as in the proof of Theorem 1.1 (left to right), using Lemma 2.1, but with an additional case, namely when G is a definable (over K) subgroup of a simple abelian variety A (over K). We will now refer to Fact 1.7(ii) of [15] where the following is stated, with explanations and references. The so-called Manin map μ is a definable (over K) homomorphism from A onto $(U, +)^d$ where $d = dim(A)$ and where $ker(\mu)$ is the unique smallest connected infinite definable subgroup of A , which we usually write as A^\sharp . In the case when A descends to the constants (so to C_K) then μ coincides with Kolchin’s logarithmic derivative, and A^\sharp is simply $A(\mathcal{C})$, and as C_K is algebraically closed we see that $A^\sharp(K) = A^\sharp(K^{diff})$. In the case when A does not descend to the constants, then A^\sharp is *strongly minimal*. Strong minimality plus the fact that A^\sharp contains the torsion subgroup of A implies that $A^\sharp(K) = A^\sharp(K^{diff})$ (see Lemma 2.2 of [15]).

Going back to our K -definable subgroup G of A , let $B = \mu(A)$, and $C = G \cap A^\sharp$. So B is a linear DAG over K and by Theorem 1.1, $H_\partial^1(K, B)$ is trivial. So using Lemma 2.1, it remains to show that $H_\partial^1(K, C)$ is trivial. If C is finite, we are fine as K is algebraically closed. Otherwise by strong minimality of A^\sharp , $C = A^\sharp$. Let X be a left PHS for C defined over K . Let $\alpha \in X(K^{diff})$. Then clearly $L = K\langle\alpha\rangle$ is a generalized strongly normal extension of K (with respect to C), as $C(K) = C(K^{diff})$, so by our assumption $\alpha \in X(K)$, and we finish.

Proof of Corollary 1.4.

We are in the context of fields K equipped with two commuting derivations

∂_x and ∂_t . The model-theoretic environment is $DCF_{0,2}$, mentioned earlier. Our notation for fields of constants is now a bit different. Given a differential field K , K^{∂_x} denotes the field of constants of K with respect to ∂_x , which note is a ∂_t field. If K is a model of $DCF_{0,2}$ then $(K^{\partial_x}, \partial_t)$ is a model of DCF_0 .

As K^{∂_x} is algebraically closed, by [25] there exists a *PPV* extension L of K for the equation $\partial_x Y = AY$. Let Z be a fundamental matrix of solutions which generates L over K (as a $\{\partial_x, \partial_t\}$ -field). Let K^{diff} be a differential closure of K (with respect to both derivations). Let L_1 be the differential field generated by L and $(K^{diff})^{\partial_x}$, and let K_1 be the differential field generated by K and $(K^{diff})^{\partial_x}$, and let $Aut(L_1/K_1)$ be group of automorphisms of L_1 over K_1 (as a $\{\partial_x, \partial_t\}$ -field). If $\sigma \in Aut(L_1/K_1)$ then $\sigma(Z) = Zc_\sigma$ for some matrix $c_\sigma \in GL_n((K^{diff})^{\partial_x})$. The map $\sigma \rightarrow c_\sigma$ gives an isomorphism between $Aut(L_1/K_1)$ and some linear differential algebraic (with respect to ∂_t) subgroup G of $GL_n((K^{diff})^{\partial_x})$ defined over K^{∂_x} . Now Propositions 4.25, 4.28, and Theorem 5.5 of [9] (which use a differential Tannakian formalism) yield a 1 – 1 map from the set of *PPV* extensions of K for the equation $\partial_x Y = AY$, up to isomorphism over K (as $\{\partial_x, \partial_t\}$ -fields) to $H_\partial^1(K^{\partial_x}, G)$ (with respect to the ambient differentially closed field $(K^{diff})^{\partial_x}, \partial_t$) which actually coincides with the differential closure of $(K^{\partial_x}, \partial_t)$. The assumption that $(K^{\partial_x}, \partial_t)$ is both algebraically closed and *PV*-closed, together with Theorem 1.1, implies that $H_\partial^1(K^{\partial_x}, G)$ is trivial, whereby we obtain uniqueness of the *PPV* extension L of K .

3 Additional questions and remarks.

There are several natural environments to which one would like to extend Theorem 1.1. The first is the positive characteristic case, where the field is equipped with an iterative Hasse derivation. The model theoretic context is the theory of separably closed fields of characteristic $p > 0$ with Ershov invariant 1. Even an appropriate formulation of the problem would be interesting as from the model theoretic point of view we are working with type-definable linear groups.

The second is difference Galois theory, with respect to a single automorphism. From the model-theoretic point of view, the first order theory would be *ACFA*. An appropriate formulation of the problem would be interesting.

The third is that of several commuting derivations. The right hand side

condition would be that the constrained cohomology of any linear differential algebraic group over K is trivial. There are various possibilities for the left hand side condition, namely the generalization of “ PV -closed” to this context.

We expect that in the second and third cases, the existence of “exotic” linear difference (differential) algebraic groups implies that the analogues of Theorem 1.1 fail. A paper on the topic, joint with Zoé Chatzidakis, is in preparation.

Finally: We can view Theorem 1.1 as a strong differential analogue of the trivial fact that a field K is algebraically closed iff $H^1(K, G)$ is trivial for every linear algebraic group over K . What then would be the differential analogue of Serre’s theorem [24] that a field K has finitely many extensions of degree n for all n iff $H^1(K, G)$ is finite for every linear algebraic group G over K ?

Acknowledgements. This paper is a write-up of talks given by the author at the Differential Algebra workshop in CUNY, April 2016, and the Differential Algebra and Related Topics meeting in CUNY, October 2016. Thanks to the organizers for the hospitality and stimulating environment. Relevant grants are mentioned earlier.

References

- [1] Ch. Berline and D. Lascar, Superstable groups, *Annals of Pure and Applied Logic* 30 (1986), 1-43.
- [2] A. Borel, *Linear Algebraic Groups*, Springer 1991.
- [3] A. Buium, *Differential algebraic groups of finite dimension*, Springer Lecture Notes 1509, Springer, 1992.
- [4] Ph. J. Cassidy, Differential algebraic groups, *American J. Math.* 94 (1972), 891-954.
- [5] Ph. J. Cassidy, The classification of the semisimple differential algebraic groups and linear semisimple differential algebraic Lie algebras, *Journal of Algebra*, 121 (1989), 169-238.

- [6] Ph. J. Cassidy, Unipotent differential algebraic groups, in Contributions to Algebra, Academic Press, New York 1977, 83-115.
- [7] Ph. J. Cassidy and M. Singer, Galois theory of parameterized differential equations and linear differential algebraic groups, in *Differential equations and quantum groups*, vol. 9, IRMA Lect. Math. Theor. Phys., 113-155, European Math. Soc., 2007.
- [8] Ph. J. Cassidy and M. Singer, A Jordan-Hölder Theorem for differential algebraic groups, *Journal of Algebra*, 328 (2011), 190-217.
- [9] H. Gillet, S. Gorchinskiy, and A. Ovchinnikov, Parameterized Picard-Vessiot extensions and Atiyah extensions, *Advances in Mathematics* 238 (2013), 322-411
- [10] M. Kamensky and A. Pillay, Interpretations and differential Galois extensions, *International Math. Research Notices*, 2016.
- [11] E. R. Kolchin, *Differential Algebra and Algebraic Groups*, Academic Press, 1973.
- [12] E. R. Kolchin, *Differential Algebraic Groups*, Academic Press, 1985.
- [13] O. Léon Sanchez and A. Pillay, Some definable Galois theory and examples, submitted to *Bulletin Symbolic Logic*.
- [14] D. Marker, Differential fields, in *Model theory of Fields*, second edition, Lecture Notes in Logic, ASL-AK Peters, 2006.
- [15] D. Marker and A. Pillay, Differential Galois Theory III; some inverse problems, *Illinois J. Math.*, 41 (1997), 453-461.
- [16] A. Pillay, Differential Galois Theory I, *Illinois J. Math.*, 42 (1998), 678-698.
- [17] A. Pillay, Differential Galois Theory II, *Annals of Pure and Applied Logic* 88 (1997), 181-191.
- [18] A. Pillay, Algebraic D -groups and differential Galois theory, *Pacific J. Math.*, 216 (2004), 343-360.

- [19] A. Pillay, Some foundational questions concerning differential algebraic groups, *Pacific J. Math.*, 179 (1997), 179-200.
- [20] A. Pillay, Remarks on Galois cohomology and definability, *Journal of Symbolic Logic*, 62 (1997), 487-492.
- [21] A. Pillay, Differential algebraic groups and the number of countable models, in *Model theory of Fields*, second edition, Lecture Notes in Logic, ASL – A.K. Peters, 2006.
- [22] A. Pillay and W.Y. Pong, On Lascar Rank and Morley Rank of Definable Groups in Differentially Closed Fields, *JSL*, 67 (2002), 1189-1196.
- [23] B. Poizat, *Groupes Stables*, Nur al-mantiq wal-ma'rifah, Villeurbanne, 1987.
- [24] J.P Serre, *Cohomologie Galoisienne*, Springer Lecture Notes in Mathematics, 1994.
- [25] M. Wibmer, Existence of ∂ -parameterized Picard-Vessiot extensions over fields with algebraically closed constants, *Journal of Algebra* 361 (2012), 163-171.