

STABLE GROUPS AND EXPANSIONS OF $(\mathbb{Z}, +, 0)$

GABRIEL CONANT AND ANAND PILLAY

ABSTRACT. We show that if G is a stable group of finite weight with no infinite, infinite-index, chains of definable subgroups, then G is superstable of finite U -rank. Combined with recent work of Palacín and Sklinos, we conclude that $(\mathbb{Z}, +, 0)$ has no proper stable expansions of finite weight. A corollary of this result is that if $P \subseteq \mathbb{Z}^n$ is definable in a finite dp-rank expansion of $(\mathbb{Z}, +, 0)$, and $(\mathbb{Z}, +, 0, P)$ is stable, then P is definable in $(\mathbb{Z}, +, 0)$. In particular, this answers a question of Marker on stable expansions of the group of integers by sets definable in Presburger arithmetic.

1. INTRODUCTION AND SUMMARY OF MAIN RESULTS

The work in this paper is motivated by questions surrounding first-order expansions of the group $(\mathbb{Z}, +, 0)$, which are well-behaved with respect to some notion of model theoretic tameness (e.g. stability or NIP). The group $(\mathbb{Z}, +, 0)$ is a well-known example of a stable group, and so this program is a natural analog of the very fruitful study of “tame” (e.g. o-minimal or NIP) expansions of the real ordered field $(\mathbb{R}, +, \cdot, <, 0)$. Expansions of $(\mathbb{Z}, +, 0)$ have emerged in the context of definable subgroups of finitely generated free groups, as well as the general growing industry of research on ordered abelian groups satisfying notions of tameness coming from dp-rank in NIP first-order theories (e.g. [7], [9], [24]). We will provide more detail on these contexts toward the end of the introduction. For now, we state an explicit question, originally asked by Marker in 2011.

Question 1.1 (Marker). Is there a set $P \subseteq \mathbb{Z}^n$, definable in Presburger arithmetic $(\mathbb{Z}, +, <, 0)$, such that $(\mathbb{Z}, +, 0, P)$ is a proper stable expansion of $(\mathbb{Z}, +, 0)$?

The focus on Presburger arithmetic in the previous question is not unnatural. Indeed, $(\mathbb{Z}, +, <, 0)$ is an ordered structure, and thus unstable, but is still well understood and very well behaved model theoretically (to be precise, its theory is NIP of dp-rank 1 [8]). Our first main result will show that, in fact, these model theoretic notions completely control the answer to Marker’s question.

Theorem 1.2. *If $P \subseteq \mathbb{Z}^n$ is definable in a finite dp-rank expansion of $(\mathbb{Z}, +, 0)$, and $(\mathbb{Z}, +, 0, P)$ is stable, then P is definable in $(\mathbb{Z}, +, 0)$.*

The notion of dp-rank in NIP theories has been an important tool in extending the work of stability theory to the unstable stable setting (see, e.g., [23]), and so Theorem 1.2 establishes a fundamental fact about the behavior of NIP expansions of $(\mathbb{Z}, +, 0)$. The proof of this theorem will be obtained from a more general result

Date: August 23, 2016.

2010 Mathematics Subject Classification. Primary: 03C45; Secondary: 03C64.

Key words and phrases. Stable groups, finite weight, finite dp-rank, Presburger arithmetic.

The second author was supported by NSF grant DMS-136072.

on stable groups (Theorem 1.4 below), combined with the following recent result of Palacín and Sklinos [15].

Fact 1.3. [15] $(\mathbb{Z}, +, 0)$ has no proper stable expansions of finite U -rank.

Precise definitions of these notions will be given in Section 2. For now, we emphasize that, *a priori*, Fact 1.3 alone is not sufficient to answer Marker's question, or obtain Theorem 1.2. In particular, while the dp-rank of complete stable theory is *bounded above* by its U -rank, there is no further general relationship between these two ranks. Indeed, there are stable groups of dp-rank 1 and infinite, even undefined, U -rank (see Example 2.9). Therefore, the work involved in proving Theorem 1.2 consists of showing that if a stable expansion of $(\mathbb{Z}, +, 0)$ has finite dp-rank, then it must have finite U -rank (we also show that these ranks are the same). We will in fact obtain this conclusion from a general characterization of superstable groups of finite U -rank, which exploits the notion of *weight* in stable theories. Before stating this result, we clarify the following terminology.

Given a group G , definable in (a model of) a complete theory T , we say G has *finite dp-rank* if there is a uniform finite bound on the dp-rank of any type containing the formula defining G . If G is a *stable* group (i.e. if T is stable), then the U -rank of G , denoted $U(G)$, is the supremum of the U -ranks of types containing a formula defining G . Replacing U -rank with *weight*, we similarly define the *weight* of G , denoted $\text{wt}(G)$. We let $<_\infty$ denote the partial order on groups given by: $H <_\infty K$ if $H \leq K$ and $[K : H] = \infty$. If G is superstable of finite U -rank then, by well-known facts, G necessarily has finite weight and no infinite $<_\infty$ -chains of definable subgroups. Our main result is that these conditions are also sufficient.

Theorem 1.4. *Let G be a stable group. The following are equivalent.*

- (i) G is superstable of finite U -rank.
- (ii) G has finite weight and no infinite $<_\infty$ -chains of definable subgroups.

The proof is given in Section 3, and involves a new application of Zilber indecomposability in the setting of weight. Theorem 1.4 quickly yields Theorem 1.2, modulo Fact 1.3 and general results on weight and dp-rank in stable theories.

Proof of Theorem 1.2. Let $G = (\mathbb{Z}, +, 0, P)$, where $P \subseteq \mathbb{Z}^n$ is definable in a finite dp-rank expansion of $(\mathbb{Z}, +, 0)$. Since dp-rank cannot increase after taking a reduct, G has finite dp-rank $n < \omega$. Assume G is stable. Then $\text{wt}(G) = n$ (see [1], [14]), and so $U(G)$ is finite by Theorem 1.4 (in fact, $U(G) = \text{wt}(G) = n$ by Corollary 1.5 below). By Fact 1.3, P is definable in $(\mathbb{Z}, +, 0)$. \square

Altogether, the real content of this paper is the proof of Theorem 1.4. From the proof of this result, we will also obtain the following corollary.

Corollary 1.5. *Suppose G is a stable group of finite weight, with no infinite $<_\infty$ -chains of definable subgroups. Then there is a uniform finite bound on the length of a $<_\infty$ -chain of definable subgroups of G . Moreover, if $n < \omega$ is the length of the longest $<_\infty$ -chain of definable subgroups of G , then $U(G) \leq n \text{wt}(G)$.*

In Section 2, we will also recall some classical examples showing that the upper bound in this result cannot be improved in general.

We end this section with a discussion of related work and open questions. The motivation for Question 1.1 partly arose from interest in the induced structure on proper definable subgroups of finitely-generated free groups, which are examples of

stable groups [21]. In particular, the maximal proper definable subgroups of such groups are exactly the centralizers of some nontrivial element (see [16]), and thus isomorphic *as groups* to $(\mathbb{Z}, +, 0)$. Therefore, studying stable expansions of $(\mathbb{Z}, +, 0)$ was seen as an alternate approach toward the unpublished result of Perin that the induced structure on centralizers in the free group is always a pure group. Another proof of this has been recently given by Byron and Sklinos [4].

Beyond this connection to the free group, there has been a recent flurry of interest in expansions of $(\mathbb{Z}, +, 0)$. On the stable side, we have the following ambitious question (which is similar to a question of Goodrick quoted in [15]).

Question 1.6. Characterize the sets $P \subseteq \mathbb{Z}^n$, for which $(\mathbb{Z}, +, 0, P)$ is stable.

On the unstable side, Dolich and Goodrick [7] have shown that $(\mathbb{Z}, +, <, 0)$ has no proper *strong* expansions (which includes expansions of finite dp-rank). Concerning *reducts* of Presburger arithmetic, a recent result of the second author [6] is that there are no intermediate structures strictly between $(\mathbb{Z}, +, 0)$ and $(\mathbb{Z}, +, <, 0)$. In a different direction, Kaplan and Shelah [11] show that if $P = \{z \in \mathbb{Z} : |z| \text{ is prime}\}$ then $(\mathbb{Z}, +, 0, P)$ is unstable and, assuming a fairly strong conjecture in number theory, $(\mathbb{Z}, +, 0, P)$ is supersimple of SU -rank 1 (see also Remark 1.8(3) below).

The investigation of stable expansions of $(\mathbb{Z}, +, 0)$ also fits naturally into the general question of when good properties of a structure are preserved after adding a new predicate. For example Pillay and Steinhorn [17] proved that there are no proper o-minimal expansions of $(\mathbb{N}, <)$, while Marker [12] exhibited proper strongly minimal expansions of $(\mathbb{N}, x \mapsto x + 1)$. Zilber [26] showed that there are proper ω -stable expansions of the complex field $(\mathbb{C}, +, \cdot, 0, 1)$ (in particular, adding a predicate for the roots of unity), while Marker [13] proved that there are no proper stable expansions of $(\mathbb{C}, +, \cdot, 0, 1)$ by a semialgebraic set.

Even more generally, Theorem 1.4 fits into the investigation of when stronger forms of stability can be proved for stable groups satisfying various assumptions on definable subgroups. For example, in [2], Baldwin and Pillay prove that if G is superstable of finite U -rank, and G has no proper connected type-definable normal subgroups, then G is ω -stable. In [10], Gagelman proves that if G is superstable of finite U -rank and satisfies the descending chain condition on definable subgroups, then G is ω -stable. It would be interesting to know if the finiteness conditions on weight and U -rank in Theorem 1.4 can be relaxed to obtain a characterization of superstable groups of a similar flavor. In particular, it is well known that if G is a superstable group, then every type in G has finite weight (i.e. G is *strongly stable*) and G has no infinite descending $<_\infty$ -chains of definable subgroups (i.e. G satisfies the *superstable descending chain condition*). Therefore, we ask the following question, which is an analog of Theorem 1.4 for superstable groups.

Question 1.7. Suppose G is a strongly stable group satisfying the superstable descending chain condition. Is G superstable?

We end with some important remarks.

Remark 1.8.

- (1) Many of the results above on $(\mathbb{Z}, +, 0)$ do not hold if one considers expansions of structures elementarily equivalent to $(\mathbb{Z}, +, 0)$. For example, there are models $(M, +, 0)$ of $\text{Th}(\mathbb{Z}, +, 0)$ with proper stable expansions of finite U -rank.

- (2) Theorem 1.2 also holds with inp-rank in place of dp-rank , since these ranks coincide in the stable case (see [1]). Therefore the theorem can be applied in the more general class of NTP_2 theories.
- (3) Fact 1.3 does not hold if stable is replaced by simple. For example, using results of Chatzidakis and Pillay [5] on “generic” predicates, one can find a set $P \subseteq \mathbb{Z}$ such that $(\mathbb{Z}, +, 0, P)$ is unstable, but supersimple of U -rank 1.

2. PRELIMINARIES

The purpose of this section is to collect the preliminary tools and facts that we will need in the proof of the main result (Theorem 1.4). Our intent is to include sufficient detail so as to make this paper accessible to a wider audience beyond those researchers well-versed in stability theory. For example, Lemma 2.3 and Proposition 2.11 are folkloric facts, which seem to be primarily used in the superstable context, and to not appear in the literature in more general settings. Therefore we have included proofs suitable for the general stable case.

Throughout this section, T is a stable first-order theory, and we assume $T = T^{\text{eq}}$. We work in a sufficiently saturated monster model \mathbb{M} of T , and use letters A, B, \dots for small parameter sets in \mathbb{M} , where a parameter set A is *small* (written $A \subset \mathbb{M}$) if \mathbb{M} is $|T(A)|^+$ -saturated. In general, a cardinal κ is *small* or *bounded* if \mathbb{M} is κ^+ -saturated. We use letters X, Y, \dots for definable or type-definable sets, and we always identify such a set X with its set of realizations $X(\mathbb{M})$ in the monster model. As usual, by a *type-definable* set we mean an intersection of a small collection of definable sets. Given a type p , and a type-definable set X , we write $p \models X$ if p extends a type defining X . We use \perp for nonforking independence in T . We assume familiarity with stability and U -rank.

Definition 2.1.

- (1) Given a sequence $(\bar{b}_i)_{i \in I}$ of tuples and $C \subset \mathbb{M}$, we say $(\bar{b}_i)_{i \in I}$ is C -**independent** if $\bar{b}_i \perp_C \{\bar{b}_j : j \neq i\}$ for all $i \in I$.
- (2) Given $C \subset \mathbb{M}$ and $p \in S(C)$, define the **weight of p** , denoted $\text{wt}(p)$, to be the supremum over cardinals κ for which there is $B \supseteq C$, a realization $\bar{a} \models p$, and a B -independent sequence $(\bar{b}_i)_{i < \kappa}$ such that $\bar{a} \perp_C B$ and $\bar{a} \not\perp_B \bar{b}_i$ for all $i < \kappa$.
- (3) Let rk denote either U -rank or weight.
 - (i) If $\bar{a} \in \mathbb{M}$ and $C \subset \mathbb{M}$ then $\text{rk}(\bar{a}/C)$ denotes $\text{rk}(\text{tp}(\bar{a}/C))$.
 - (ii) If X is type-definable, then $\text{rk}(X) = \sup\{\text{rk}(p) : p \models X\}$.

We will use the following basic properties of U -rank and weight.

Proposition 2.2.

Let rk denote either U -rank or weight.

- (a) *Given $\bar{a} \in \mathbb{M}$ and $C \subset \mathbb{M}$, $\text{rk}(\bar{a}/C) = 0$ if and only if $\bar{a} \in \text{acl}(C)$.*
- (b) *Fix $\bar{a}, \bar{b} \in \mathbb{M}$ and $C \subset \mathbb{M}$. If $\bar{a} \in \text{acl}(\bar{b}, C)$ and $\bar{b} \in \text{acl}(\bar{a}, C)$ then $\text{rk}(\bar{a}/C) = \text{rk}(\bar{b}/C)$.*
- (c) *Suppose X is type-definable and f is a definable function with domain containing X . Then $\text{rk}(f(X)) \leq \text{rk}(X)$.*

Proof. These are straightforward exercises. Parts (b) and (c) follow easily from part (a) together with Lascar’s inequality for U -rank (see [18, Theorem 19.4]), and a sufficiently similar inequality for weight (see [22, Lemma V.3.11(2)]). \square

In a superstable theory, the weight of a type p is bounded by the sum of the integer coefficients in the Cantor normal form of $U(p)$ (see [18, Theorem 19.9]). In particular, one has $\text{wt}(p) \leq U(p)$, which still holds for stable theories in general.

Lemma 2.3. *If $C \subset \mathbb{M}$ and $p \in S(C)$, then $\text{wt}(p) \leq U(p)$.*

Proof. Fix $p \in S(C)$. Suppose we have a set $B \supseteq C$, a realization $\bar{a} \models p$, and a B -independent sequence $(\bar{b}_i)_{i < \kappa}$, for some cardinal κ , such that $\bar{a} \downarrow_C B$ and $\bar{a} \not\downarrow_B \bar{b}_i$ for all $i < \kappa$. We prove $U(\bar{a}/B) \geq \kappa$, which implies $U(p) \geq \kappa$. Given $i \leq \kappa$, define $B_i = B \cup \{\bar{b}_j : i \leq j\}$ (so $B_\kappa = B$). We prove, by induction on $i \leq \kappa$, that $U(\bar{a}/B_i) \geq i$. Given this, we will then have $U(\bar{a}/B) = U(\bar{a}/B_\kappa) \geq \kappa$.

The base case is trivial; so suppose $\lambda \leq \kappa$ is a limit ordinal and $U(\bar{a}/B_i) \geq i$ for all $i < \lambda$. For any $i < \lambda$, we have $B_\lambda \subseteq B_i$, and so $U(\bar{a}/B_\lambda) \geq U(\bar{a}/B_i) \geq i$. Therefore $U(\bar{a}/B_\lambda) \geq \lambda$. Finally, fix $i < \kappa$ and suppose $U(\bar{a}/B_i) \geq i$. Since $B_{i+1} \downarrow_B \bar{b}_i$ and $\bar{a} \not\downarrow_B \bar{b}_i$, we have $\bar{a} \not\downarrow_{B_{i+1}} \bar{b}_i$ by transitivity. Therefore $U(\bar{a}/B_{i+1}) \geq i + 1$. \square

For general stable theories, Lemma 2.3 is the most one can say concerning the relationship between weight and U -rank for arbitrary types (see Example 2.9). However, when working “close” to types of U -rank 1, weight and U -rank coincide. This will be a key tool in the proof of our main result.

Proposition 2.4. *Fix $C \subset \mathbb{M}$, and suppose $X \subseteq \mathbb{M}$ is such that $U(a/C) \leq 1$ for all $a \in X$. If \bar{b} is a finite tuple in $\text{acl}(XC)$ then $U(\bar{b}/C) = \text{wt}(\bar{b}/C)$.*

Proof. We first consider the case that \bar{b} is a tuple of elements of X . The essential observation is the following:

(\dagger) If $a \in X$ and $C \subseteq B \subseteq D \subset \mathbb{M}$ then, since $U(a/C) \leq 1$, we have $a \not\downarrow_B D$ if and only if $a \in \text{acl}(D) \setminus \text{acl}(B)$.

In particular, by (\dagger) and symmetry of forking, we have the following exchange property: given $a, a' \in X$ and $C \subseteq B \subset \mathbb{M}$, if $a \in \text{acl}(B, a') \setminus \text{acl}(B)$ then $a' \in \text{acl}(B, a)$. Therefore, a tuple $\bar{a} \in X$ has a well-defined basis over C , and we show $U(\bar{a}/C) = \dim_{\text{acl}}(\bar{a}/C) = \text{wt}(\bar{a}/C)$. The first equality follows from Proposition 2.2(a) and repeated application of Lascar’s inequality. For the second equality, it follows from (\dagger) that a basis for \bar{a} over C is C -independent (with respect to nonforking) and, moreover, $\bar{a} \not\downarrow_C a_i$ for any $a_i \in \bar{a} \setminus \text{acl}(C)$. Therefore $\text{wt}(\bar{a}/C) \geq \dim_{\text{acl}}(\bar{a}/C)$, and so equality holds by Lemma 2.3.

Now consider the general case of $\bar{b} \in \text{acl}(XC)$. By Proposition 2.2(a), we may assume that some coordinate of \bar{b} is not in $\text{acl}(C)$. Fix $\bar{a} = (a_1, \dots, a_n) \in X$, algebraically independent over C , with $\bar{b} \in \text{acl}(\bar{a}, C)$. Let $k \leq n$ be maximal such that, for some $i_1 < \dots < i_k \leq n$, we have $\bar{b} \downarrow_C (a_{i_1}, \dots, a_{i_k})$ (it is possible here that $k = 0$). Without loss of generality, assume $\bar{b} \downarrow_C (a_1, \dots, a_k)$. Let $\bar{a}_1 = (a_1, \dots, a_k)$ and $\bar{a}_2 = (a_{k+1}, \dots, a_n)$. We show $\bar{a}_2 \in \text{acl}(\bar{b}, \bar{a}_1, C)$. If not, then there is some $k < i \leq n$ such that $a_i \notin \text{acl}(\bar{b}, \bar{a}_1, C)$, and so $a_i \downarrow_{\bar{a}_1, C} \bar{b}$ by (\dagger). Together with $\bar{a}_1 \downarrow_C \bar{b}$, we have $(\bar{a}_1, a_i) \downarrow_C \bar{b}$, contradicting the maximality of k . Altogether, $\bar{a}_2 \in \text{acl}(\bar{b}, \bar{a}_1, C)$ and $\bar{b} \in \text{acl}(\bar{a}_2, \bar{a}_1, C)$. By the first case above, and Proposition 2.2(b), $U(\bar{b}/C) = U(\bar{a}_2/C) = \text{wt}(\bar{a}_2/C) = \text{wt}(\bar{b}/C)$. \square

Remark 2.5. The notion of weight also behaves nicely in simple theories. For example, after replacing all occurrences of U -rank with SU -rank, the statements

of Proposition 2.2, Lemma 2.3, and Proposition 2.4 hold when T is simple (with identical proofs).

We now turn to stable groups. Once again, a *stable group* is a group G definable in (some model of) a stable theory T . We will continue to identify the definable set G with its realization $G(\mathbb{M})$ in the monster model. Given a definable group G , we let G^0 denote the *connected component of G* , which is the intersection of all definable subgroups of G of finite index. By stability (see Fact 2.6 below), G^0 is the intersection of at most $|T|$ many definable subgroups of G of finite index, and hence is type-definable (over the same parameters used to define G). We say G is *connected* if $G = G^0$.

We first recall the following classical results.

Fact 2.6. *Let G be a stable group.*

- (a) (Baldwin-Saxl [19, Proposition 1.4]) *Let $\{H_i : i \in I\}$ be a family of uniformly definable subgroups of G , and set $H = \bigcap_{i \in I} H_i$. Then $H = \bigcap_{i \in I_0} H_i$ for some finite $I_0 \subseteq I$. In particular, H is definable.*
- (b) (Poizat [19, Theorem 5.17]) *Any type-definable subgroup of G is the intersection of at most $|T|$ many definable subgroups of G .*

Remark 2.7. Unlike the previous preliminaries, these facts on stable groups do not immediately go through if T is only assumed to be simple. In fact, there are simple unstable groups where part (a) fails [25, Example 1]. On the other hand, whether part (b) holds for groups definable in simple theories is a well known open question.

For the sake of clarity, it is worth making a few remarks concerning weight and U -rank of stable groups. In particular, given a definable group G and $A \subset \mathbb{M}$, we let $S_G(A)$ denote the space of complete types, over parameters in A , which contain a formula defining G . Then, if rk denotes either U -rank or weight, we can express $\text{rk}(G)$ as

$$\text{rk}(G) = \sup\{\text{rk}(p) : p \in S_G(A) \text{ for some } A \subset \mathbb{M}\}.$$

We say G has *finite U -rank* (resp. *finite weight*) if $U(G) < \omega$ (resp. $\text{wt}(G) < \omega$).

If G is stable then $U(p) = U(G)$ for any generic type p in G (see [3, Lemma III.4.5(i)]). However, this can fail for weight. For example, it is possible that all types in G have finite weight, but $\text{wt}(G)$ is not finite (e.g. Example 2.9(1) below). Since our focus is on the case that $\text{wt}(G)$ is finite, we will not concern ourselves with this situation.

Remark 2.8. When considering examples of stable groups, it is often the case that the group G is the whole structure (i.e. defined by the formula $x = x$). Therefore, given a group $G = (G, \cdot, 1, \dots)$, when we speak of the U -rank or weight of G , we continue to mean as calculated in a monster model according to the definitions and conventions above.

The following examples illustrate some of the possible variety concerning weight, U -rank, and $<_\infty$ -chains in stable groups.

Example 2.9.

- (1) Let $\Pi_2 = \{2^n : n \in \mathbb{N}\}$ and let $G = (\mathbb{Z}, +, 0, \Pi_2)$. Then G is superstable of U -rank ω (see [15], [20]). Therefore, G is strongly stable, but does not have finite weight by Theorem 1.4.

- (2) Fix an integer $n > 0$ and let $G = (\mathbb{Q}^n, +, 0, (H_k)_{k < n})$ where, for each $k < n$, $H_k = \mathbb{Q}^k \times \{0\}^{n-k}$. We have a sequence $(E_k)_{k < n}$ of definable equivalence relations, given by $E_k(x, y) \leftrightarrow x - y \in H_k$. Let \mathbb{M} be a monster model. Given $a, b \in \mathbb{M}$, let $d(a, b) = \min\{k < n : E_k(a, b)\}$. Then d is an ultrametric on \mathbb{M} , taking values in $\{0, 1, \dots, n\}$; and nonforking independence is characterized by: $A \downarrow_C B$ if and only if, for all $a \in \text{acl}(AC)$, $d(a, \text{acl}(BC)) = d(a, \text{acl}(C))$ (where algebraic closure is the same as in $(\mathbb{Q}^n, +, 0)$). Using this, one may verify that G is superstable of U -rank n and weight 1.
- (3) Let $G = (\mathbb{Q}^\omega, +, 0, (H_n)_{n < \omega})$ where, for each $n < \omega$, $H_n = \mathbb{Q}^n \times \{0\}^\omega$. Using a similar argument as in part (2), one may show that G is superstable of U -rank ω and weight 1.
- (4) Let $G = (\mathbb{Q}^\omega, +, 0, (K_n)_{n < \omega})$ where, for each $n < \omega$, $H_n = \{0\}^n \times \mathbb{Q}^\omega$. Then G is strictly stable of weight 1.

Our final preliminary tools concern indecomposable sets in stable groups.

Definition 2.10. Let G be a stable group. A type-definable set $X \subseteq G$ is **indecomposable** if, for all type-definable subgroups $H \leq G$, either X/H is unbounded or $|X/H| = 1$ (where $X/H = \{xH : x \in X\}$).

Proposition 2.11. *Let G be a stable group. Fix $A \subset \mathbb{M}$ and a stationary type $p \in S_G(A)$. Let $X = p(\mathbb{M})$. Then $X \subseteq G$ is indecomposable.*

Proof. Let \mathcal{F} denote the family of type-definable subgroups $H \leq G$ such that X/H is bounded. Let H_0 be the intersection of the elements of \mathcal{F} . Using Fact 2.6, it is a standard exercise to show that H_0 is a type-definable subgroup of G and X/H_0 is bounded (i.e. $H_0 \in \mathcal{F}$). Note also that A -invariance of X implies A -invariance of H_0 , and so H_0 is type-definable over A .

Let $\tilde{p} \in S_G(\mathbb{M})$ be the unique global nonforking extension of p . Let $C \subset X$ be a bounded set such that $X/H_0 = \{cH_0 : c \in C\}$, and fix a realization $u \in G$ of $\tilde{p}|_{AC}$. Then $u \in X$, and so $u \in cH_0$ for some $c \in C$, which means $\tilde{p} \models cH_0$. If $f \in \text{Aut}(\mathbb{M}/A)$ then, by A -invariance of H_0 and \tilde{p} , we have $\tilde{p} \models f(c)H_0$, and so $f(cH_0) = f(c)H_0 = cH_0$. Consequently, cH_0 is type-definable over A , and so $p \models cH_0$. Therefore $X \subseteq cH_0$, which implies $X \subseteq cH$ for all $H \in \mathcal{F}$. \square

A well-known result of Berline and Lascar is the Indecomposability Theorem for superstable groups [3, Theorem V.3.1]. In order to use this result without the assumption of superstability, we state the following corollary of its proof.

Fact 2.12. *Suppose G is a stable group and $\{X_i : i \in I\}$ is a family of indecomposable type-definable subsets of G , each containing 1. Given $n > 0$ and $\sigma = (i_0, \dots, i_n) \in I^{<\omega}$, let $X_\sigma = X_{i_0} \cdot X_{i_1} \cdot \dots \cdot X_{i_n}$. Assume that there is a uniform finite bound on $U(X_\sigma)$, where σ ranges over $I^{<\omega}$. Then $\bigcup_{i \in I} X_i$ generates a connected type-definable subgroup H of G . In particular, there are i_0, \dots, i_n such that $H = X_{i_0} \cdot \dots \cdot X_{i_n} \cdot X_{i_n}^{-1} \cdot \dots \cdot X_{i_0}^{-1}$.*

3. PROOF OF THE MAIN RESULT

Toward the proof of Theorem 1.4, the first step is to prove that if G is a stable group of finite weight, then one may construct a type-definable subgroup of G with several nice properties.

Lemma 3.1. *If G is an infinite stable group of finite weight then there is an infinite connected type-definable normal subgroup $H \leq G$, with $U(H) = \text{wt}(H)$.*

Proof. Fix a stationary type $p \in S_G(A)$, for some $A \subset \mathbb{M}$, such that $U(p) = 1$. For example, choose p minimal in the fundamental order among non-algebraic types in $S_G(A)$ (with A varying over small parameter sets in \mathbb{M}), and then replace p by a nonforking extension to a model.

Let $Y = p(\mathbb{M})$. Then $Y \subseteq G$ is indecomposable by Proposition 2.11. Fix some $u \in Y$, and set $X = u^{-1}Y$. Given $g \in G$, let $X^g = gXg^{-1}$. Then $\{X^g : g \in G\}$ is a family of indecomposable type-definable subsets of G , each of which contains 1. By Proposition 2.2(c), $U(X^g) = 1$ for all $g \in G$. Given a sequence $\sigma = (g_0, \dots, g_n)$ of elements of G , we set $X_\sigma = X^{g_0} \cdot \dots \cdot X^{g_n}$.

We claim that, for any $\sigma \in G^{<\omega}$, $U(X_\sigma) = \text{wt}(X_\sigma)$. Suppose, for a contradiction, that $\text{wt}(X_\sigma) < U(X_\sigma)$, with $\sigma = (g_0, \dots, g_n)$. Then there is a type q such that $q \models X_\sigma$ and $U(q) > \text{wt}(X_\sigma)$. After taking a nonforking extension, we may assume $q \in S(C)$, where $C \subset \mathbb{M}$ is such that X^{g_i} is type-definable over C for all $0 \leq i \leq n$. Let $a \models q$. Then $a \in X_\sigma$, so we may write $a = b_0 \cdot \dots \cdot b_n$ with $b_i \in X^{g_i}$. For any $0 \leq i \leq n$, $\text{tp}(b_i/C) \models X^{g_i}$ by choice of C , and so $U(b_i/C) \leq 1$. Since $a \in \text{acl}(b_0, \dots, b_n)$, we have $U(a/C) = \text{wt}(a/C)$ by Proposition 2.4. But $\text{wt}(a/C) \leq \text{wt}(X_\sigma)$, which contradicts $U(a/C) = U(q) > \text{wt}(X_\sigma)$.

We have $U(X_\sigma) \leq \text{wt}(G)$ for all $\sigma \in G^{<\omega}$. Therefore, we may apply Fact 2.12 to conclude that $\bigcup_{g \in G} X^g$ generates an infinite connected type-definable subgroup H of G , which is normal by construction. Moreover, $H = X_\sigma$ for some $\sigma \in G^{<\omega}$, and so $U(H) = \text{wt}(H)$. \square

Given a group G , and a normal subgroup K , we let $\rho_{G/K}$ denote the pullback function on subgroups of G/K , i.e., given $H \leq G/K$, define $\rho_{G/K}(H) = \{g \in G : gK \in H\} \leq G$. The following are easy observations.

Proposition 3.2. *Let G be a group and K a normal subgroup of G .*

- (a) *If $H \leq G/K$ then K is a normal subgroup of $\rho_{G/K}(H)$ and $H = \rho_{G/K}(H)/K$.*
- (b) *If $H_1 \leq H_2 \leq G/K$ then $\rho_{G/K}(H_1) \leq \rho_{G/K}(H_2) \leq G$ and $[\rho_{G/K}(H_2) : \rho_{G/K}(H_1)] = [H_2 : H_1]$.*

We now prove the main result.

Proof of Theorem 1.4. As mentioned in the introduction, one direction of this result follows from standard facts. Specifically, we have already seen in Lemma 2.3 that finite U -rank implies finite weight. Moreover, if G is superstable of finite U -rank then it follows from Lascar's inequality for cosets [3, Corollary III.8.2] that, in fact, G has a *uniform* finite bound on the length of a $<_\infty$ -chain of definable subgroups.

For the other direction, fix a stable group G of finite weight, with no infinite $<_\infty$ -chains of definable subgroups. We want to show that G is superstable of finite U -rank. We may clearly assume G is infinite. Suppose, toward a contradiction, that $U(G)$ is infinite (or undefined). We inductively construct sequences $(G_i)_{i < \omega}$ and $(K_i)_{i < \omega}$ of definable groups as follows.

Let $K_0 = \{1\}$ and set $G_0 = G_1 = G$. Fix $n > 0$, and suppose we have constructed $(G_i)_{i \leq n}$ and $(K_i)_{i < n}$ satisfying the following properties.

- (1) For all $0 < i < n$, K_i is an infinite definable normal subgroup of G_i .
- (2) For all $0 < i \leq n$, G_i is infinite and equal to G_{i-1}/K_{i-1} .
- (3) $U(G) \leq (n-1) \text{wt}(G) \oplus U(G_n)$.

Note that G_n is an infinite stable group and, by Proposition 2.2(c), $\text{wt}(G_n) \leq \text{wt}(G) < \omega$. By Lemma 3.1, we may fix an infinite connected type-definable normal subgroup $H \leq G_n$, with $U(H) = \text{wt}(H)$.

From the assumption on $<_\infty$ -chains of definable subgroups of G , along with Proposition 3.2, it is easy to see that G_n also has no infinite $<_\infty$ -chains of definable subgroups. Since H is type-definable, it follows (with the help of Fact 2.6(b)) that there is a definable subgroup K_n of G_n such that $H \leq K_n$ and $[K_n : H]$ is bounded. (In fact, the superstable descending chain condition is equivalent to this assertion on type-definable subgroups.) By Fact 2.6(a) and normality of H , we may replace K_n by $\bigcap_{g \in G} gK_n g^{-1}$, and thus assume K_n is normal.

Since H is connected and type-definable, with bounded index in K_n , it follows that $H = K_n^0$, which implies $U(H) = U(K_n)$ (see, e.g., [3, Sections III.4, IV,3]). Since $U(H) = \text{wt}(H)$, we must have $U(K_n) = \text{wt}(K_n) \leq \text{wt}(G)$. Let $G_{n+1} = G_n/K_n$. Then, using (3) and Lascar's inequality for cosets [3, Corollary III.8.2],

$$\begin{aligned} U(G) &\leq (n-1)\text{wt}(G) \oplus U(G_n) \\ &\leq (n-1)\text{wt}(G) \oplus U(K_n) \oplus U(G_n/K_n) \leq n\text{wt}(G) \oplus U(G_{n+1}). \end{aligned}$$

It therefore follows from our assumption on $U(G)$ that G_{n+1} must be infinite. Altogether, $(G_i)_{i \leq n+1}$ and $(K_i)_{i \leq n}$ satisfy (1), (2), (3) above.

We have now constructed $(G_n)_{n < \omega}$ and $(K_n)_{n < \omega}$. Given $n < \omega$, let ρ_n denote the pullback function ρ_{G_n/K_n} . Let $L_0 = \{1\}$ and, given $n > 0$, set $L_n = \rho_0 \dots \rho_{n-1}(K_n)$. Using Proposition 3.2, it is easy to see that $(L_n)_{n < \omega}$ is an ascending chain of definable subgroups of G and, for any $n > 0$, since K_n is infinite we have $L_{n-1} <_\infty L_n$. Therefore $(L_n)_{n < \omega}$ is an $<_\infty$ -chain, which is a contradiction. \square

Remark 3.3. In the previous proof, note that we needed to assume that G had no infinite *ascending or descending* $<_\infty$ -chains of definable groups. Specifically, no infinite descending $<_\infty$ -chains allows us to construct the definable group K_n from the type-definable group H , which is necessary to complete the induction step of the argument. No infinite ascending $<_\infty$ -chains ensures us that the inductive construction halts at some finite stage. The stable group described in Example 2.9(3) also illustrates the necessity of these assumptions.

Proof of Corollary 1.5. Let G be a stable group of finite weight, with no infinite $<_\infty$ -chains of definable subgroups. Then G has finite U -rank by Theorem 1.4 and so, by Lascar's inequality for cosets, there is a uniform finite bound on the length of a $<_\infty$ -chain of definable subgroups of G . Let $n < \omega$ be the length of the longest $<_\infty$ -chain of definable subgroups of G . Repeating the construction in the previous proof, it follows that there must be some $m \leq n$ such that G_{m+1} is finite (since otherwise we would construct a $<_\infty$ -chain (L_0, \dots, L_{n+1}) of length $n+1$). \square

Acknowledgements. We are grateful to Rizos Sklinos and Erik Walsberg for their comments on an earlier draft.

REFERENCES

- [1] Hans Adler, *Strong theories, burden, and weight*, preprint, <http://www.logic.univie.ac.at/~adler/docs/strong.pdf>, 2007.
- [2] J. T. Baldwin and A. Pillay, *Semisimple stable and superstable groups*, Ann. Pure Appl. Logic **45** (1989), no. 2, 105–127, Stability in model theory, II (Trento, 1987). MR 1044119 (91f:03064)

- [3] Ch. Berline and D. Lascar, *Superstable groups*, Ann. Pure Appl. Logic **30** (1986), no. 1, 1–43, Stability in model theory (Trento, 1984). MR 831435 (87k:03028)
- [4] Ayala Byron and Rizos Sklinos, *Fields definable in the free group*, arXiv:1512.07922 [math.LO], 2015.
- [5] Z. Chatzidakis and A. Pillay, *Generic structures and simple theories*, Ann. Pure Appl. Logic **95** (1998), no. 1-3, 71–92. MR 1650667 (2000c:03028)
- [6] Gabriel Conant, *There are no intermediate structures between the group of integers and Presburger arithmetic*, arXiv:1603.00454 [math.LO], 2016.
- [7] Alfred Dolich and John Goodrick, *Strong theories of ordered abelian groups*, arXiv:1511.08274 [math.LO], 2015.
- [8] Alfred Dolich, John Goodrick, and David Lippel, *Dp-minimality: basic facts and examples*, Notre Dame J. Form. Log. **52** (2011), no. 3, 267–288. MR 2822489 (2012h:03102)
- [9] Joseph Flenner and Vincent Guingona, *Convexly orderable groups and valued fields*, J. Symb. Log. **79** (2014), no. 1, 154–170. MR 3226016
- [10] Jerry Gagelman, *A note on superstable groups*, J. Symbolic Logic **70** (2005), no. 2, 661–663. MR 2140052 (2006a:03046)
- [11] Itay Kaplan and Saharon Shelah, *Decidability and classification of the theory of integers with primes*, arXiv:1601.07099 [math.LO], 2016.
- [12] David Marker, *A strongly minimal expansion of (ω, s)* , J. Symbolic Logic **52** (1987), no. 1, 205–207. MR 877867 (88h:03048)
- [13] ———, *Semialgebraic expansions of \mathbf{C}* , Trans. Amer. Math. Soc. **320** (1990), no. 2, 581–592. MR 964900 (90k:03034)
- [14] Alf Onshuus and Alexander Usvyatsov, *On dp-minimality, strong dependence and weight*, J. Symbolic Logic **76** (2011), no. 3, 737–758. MR 2849244 (2012m:03082)
- [15] Daniel Palacín and Rizos Sklinos, *Superstable expansions of free abelian groups*, Notre Dame J. Form. Log., to appear, available: arXiv 1405.0568.
- [16] Anand Pillay, *Forking in the free group*, J. Inst. Math. Jussieu **7** (2008), no. 2, 375–389. MR 2400726 (2009f:20033)
- [17] Anand Pillay and Charles Steinhorn, *Discrete o-minimal structures*, Ann. Pure Appl. Logic **34** (1987), no. 3, 275–289, Stability in model theory (Trento, 1984). MR 899083 (88j:03023)
- [18] Bruno Poizat, *A course in model theory*, Universitext, Springer-Verlag, New York, 2000, An introduction to contemporary mathematical logic, Translated from the French by Moses Klein and revised by the author. MR 1757487 (2001a:03072)
- [19] ———, *Stable groups*, Mathematical Surveys and Monographs, vol. 87, American Mathematical Society, Providence, RI, 2001, Translated from the 1987 French original by Moses Gabriel Klein. MR 1827833 (2002a:03067)
- [20] ———, *Supergénéria*, J. Algebra **404** (2014), 240–270, À la mémoire d’Éric Jaligot. [In memoriam Éric Jaligot]. MR 3177894
- [21] Z. Sela, *Diophantine geometry over groups VIII: Stability*, Ann. of Math. (2) **177** (2013), no. 3, 787–868. MR 3034289
- [22] Saharon Shelah, *Classification theory and the number of nonisomorphic models*, second ed., Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam, 1990. MR 1083551 (91k:03085)
- [23] Pierre Simon, *A guide to NIP theories*, vol. 44, Cambridge University Press, 2015.
- [24] ———, *On dp-minimal ordered structures*, J. Symbolic Logic **76** (2011), no. 2, 448–460. MR 2830411 (2012e:03071)
- [25] Frank Wagner, *Groups in simple theories*, Logic Colloquium ’01, Lect. Notes Log., vol. 20, Assoc. Symbol. Logic, Urbana, IL, 2005, pp. 440–467. MR 2143908
- [26] Boris Zilber, *A note on the model theory of the complex field with roots of unity*, unpublished note, <https://people.maths.ox.ac.uk/zilber/publ.html>, 1990.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN, 46656, USA