

# *Relative Manin-Mumford for semi-abelian surfaces*

D. Bertrand, D. Masser, A. Pillay, U. Zannier \*

July 2013 (ArXiv version) †

*Abstract.* - We show that Ribet sections are the only obstruction to the validity of the relative Manin-Mumford conjecture for one dimensional families of semi-abelian surfaces. Applications include special cases of the Zilber-Pink conjecture for curves in a mixed Shimura variety of dimension four, as well as the study of polynomial Pell equations with non-separable discriminants.

## **Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	The viewpoint of group schemes : relative Manin-Mumford . . .	1
1.2	The viewpoint of mixed Shimura varieties : Pink's conjecture . . .	4
1.3	Plan of the paper . . . . .	7
<b>2</b>	<b>Restatement of the Main Theorem</b>	<b>8</b>
2.1	Introducing $q$ and $p$ . . . . .	8
2.2	Isoconstant issues . . . . .	10
2.3	Antisymmetric relations and restatement . . . . .	11

---

\* *Authors' addresses* : bertrand@math.jussieu.fr, david.masser@unibas.ch, A.Pillay@leeds.ac.uk, u.zannier@sns.it

† *AMS Classification* : 14K15, 12H05, 14K20, 11J95. *Key words* : semi-abelian varieties; Manin-Mumford, André-Oort and Zilber-Pink conjectures; differential Galois theory; polynomial Pell equations.

<b>3</b>	<b>The overall strategy</b>	<b>13</b>
3.1	Algebraic lower bounds . . . . .	13
3.2	Transcendental upper bounds . . . . .	15
3.3	What remains to be done . . . . .	18
<b>4</b>	<b>A warm up : the case of direct products</b>	<b>23</b>
<b>5</b>	<b>The general case</b>	<b>30</b>
5.1	Fields of periods and the Main Lemma . . . . .	30
5.2	Reducing the Main Theorem to the Main Lemma . . . . .	33
5.3	The role of $K$ -largeness . . . . .	34
<b>6</b>	<b>Proof of the Main Lemma</b>	<b>36</b>
<b>7</b>	<b>Appendix I</b>	<b>42</b>
7.1	Analytic description of the semi-abelian logarithm . . . . .	42
7.2	Analytic description of the Ribet sections . . . . .	44
<b>8</b>	<b>Appendix II : application to Pell equations</b>	<b>46</b>

# 1 Introduction

## 1.1 The viewpoint of group schemes : relative Manin-Mumford

Let  $\mathbb{Q}^{alg}$  be the algebraic closure of  $\mathbb{Q}$  in  $\mathbb{C}$ , let  $S$  be an irreducible algebraic curve over  $\mathbb{Q}^{alg}$ , and let  $G/S$  be a semi-abelian scheme over  $S$  of relative dimension 2 and toric rank 1. We write  $G_{tor}$  for the union of all the torsion points of the various fibers of  $G \rightarrow S$ . This set  $G_{tor}$  is also the set of values at all points of  $S$  of the various torsion-sections of the group scheme  $G/S$ . Let further  $s : S \rightarrow G$  be a section of  $G/S$ . The image of  $s$  is an irreducible algebraic curve  $s(S) = W$  in  $G$ , defined over  $\mathbb{Q}^{alg}$ . Pursuing the theme of “unlikely intersections” and relative versions of the Manin-Mumford conjecture (see [31]), we here study the following question, where “strict” means “distinct from  $G$ ”.

**Question 1** : *assume that  $W \cap G_{tor}$  is infinite (i.e. Zariski dense in  $W$ ). Must  $W$  then lie in a strict subgroup scheme of  $G/S$  ?*

Let us review some of the results known along this line :

i) the analogous question has a positive answer when  $G/S$  is replaced by an abelian scheme of relative dimension 2 : see [19], [31], Theorem III.12 and §5 for non-simple ones, and [20] for the general case.

ii) Assume that *the scheme  $G/S$  is isoconstant*, i.e. isomorphic, after a finite base extension, to a product  $G = G_0 \times S$  with  $G_0/\mathbb{Q}^{alg}$ . Then, the Zariski closure  $W_0$  of the projection of  $W$  to  $G_0$  is an algebraic curve (or a point) meeting  $G_{0,tor}$  Zariski-densely. By Hindry's generalization of Raynaud's theorem on the Manin-Mumford conjecture, see [16], §5, Thm. 2,  $W_0$  is a torsion-coset of a strict algebraic subgroup  $H_0$  of  $G_0$ , and  $W$  lies in a translate of  $H = H_0 \times S$  by a torsion section. So, a positive answer is known in this case.

Therefore, the first new case occurs when  $G$  is a non isoconstant extension over  $S$  of an isoconstant elliptic scheme by  $\mathbb{G}_m$ , i.e. when  $G$  is a *semi-constant extension* in the sense of [11]. But as in [11] (though for different reasons), this case turns out to be more delicate, and the question can then have a *negative* answer. A counterexample is given in [8], and we shall refer to the corresponding sections  $s_R$  as Ribet sections of  $G/S$  and to their images  $W_R = s_R(S)$  as Ribet curves. For a formal definition, see the end of this subsection (and [17] for its initial version); a more concrete description is given in Appendix I. In this paper, we will prove that in all other cases, our question has a positive answer; in other words :

**Main Theorem** . *Let  $E/S$  be an elliptic scheme over the curve  $S/\mathbb{Q}^{alg}$ , and let  $G/S$  be an extension of  $E/S$  by  $\mathbb{G}_{m/S}$ . Let further  $s : S \rightarrow G$  be a section of  $G/S$ , with image  $W = s(S)$ .*

**(A)** *Assume that  $W \cap G_{tor}$  is infinite. Then,*

*i) either  $s$  is a Ribet section;*

*ii) or  $s$  factors through a strict subgroup scheme of  $G/S$ .*

**(B)** *More precisely,  $W \cap G_{tor}$  is infinite if and only if  $s$  is a Ribet section, or a torsion section, or a non isoconstant section of a strict subgroup scheme of  $G/S$ .*

We point out that this statement is invariant under isogenies  $G \rightarrow G'$  of the ambient group scheme, and under finite base extensions  $S' \rightarrow S$ . Throughout the paper, we will allow ourselves, sometimes tacitly, to perform such isogenies and base extensions.

We now rephrase Part (A) of the theorem according to the various types of extensions  $G/S$  and elliptic schemes  $E/S$  which can occur, and explain in each case the meaning of “isoconstant” in its Part (B); a more concrete discussion of this array of cases is given in §2. Concerning the type of  $E/S$ , we recall that the scheme  $E/S$  is isoconstant if and only if the  $j$ -invariant of its various fibers is constant; performing a finite base extension, we will then assume that  $E$  is equal to  $E_0 \times S$  for some elliptic curve  $E_0/\mathbb{Q}^{alg}$ . As for the type of  $G/S$ ,

- either  $G$  is isogenous as a group scheme over  $S$  to a direct product  $\mathbb{G}_m \times E$ . We then say that *the extension  $G/S$  is isotrivial* and perform this isogeny. Since  $W$  is flat over  $S$ , Conclusion (ii) then reads :  $W$  lies in a translate of  $E/S$  or of  $\mathbb{G}_{m/S} = \mathbb{G}_m \times S$  by a torsion section of  $G/S$  (and Conclusion (i) does not occur); in this case, the isoconstant sections of the strict subgroup schemes are the translates by torsion sections of the constant sections of  $\mathbb{G}_{m/S}$ , or of the constant sections of  $E/S$  if  $E = E_0 \times S$  is constant.

- or the extension  $G/S$  is not isotrivial. Conclusion (ii) then reads :  $W$  lies in a translate of  $\mathbb{G}_{m/S}$  by a torsion section of  $G/S$ ; in this case, the isoconstant sections are the translates by torsion sections of the constant sections of  $\mathbb{G}_{m/S}$ .

Now, whether  $G/S$  is or is not an isotrivial extension,

- we automatically get Conclusion (ii) if either the scheme  $E/S$  is not isoconstant, or it is isoconstant, but  $E_0$  does not admit complex multiplications, or if  $E_0$  is CM, but  $G$  is isoconstant (a case already covered by [16], of course).

- in the remaining case where  $E/S$  is isoconstant, with a CM elliptic curve  $E_0/\mathbb{Q}^{alg}$ , and  $G$  is not isoconstant (hence in particular, not isotrivial), Ribet sections of  $G/S$  do exist, their images  $W$  do not lie in strict subgroup schemes of  $G/S$  but meet  $G_{tor}$  infinitely often, while not all sections  $s$  satisfying the hypotheses of the theorem are Ribet sections.

In other words, both conclusions (i) and (ii) of the Main Theorem occur in this last case and are then mutually exclusive. However, there is a way of reconciling them, through the setting of Pink’s extension of the André-Oort and Zilber conjectures to mixed Shimura varieties, which we turn to in the next subsection §1.2 (see, e.g., Corollary 1).

Another type of application of the Main Theorem is given in Appendix II of the paper : this concerns the solvability of Pell’s equations over polynomial rings, and extends some of the results of [20] to the case of non-separable

discriminants.

To conclude this introduction, here is the promised definition of a *Ribet section*  $s_R : S \rightarrow G$  (see [9], §1 - a more concrete, but analytic, characterization is given in Appendix I ; several other definitions are discussed in [10]). In the notations of the Main Theorem, let  $\hat{E}/S$  be the dual of  $E/S$ : the isomorphism class of the  $\mathbb{G}_m$ -torsor  $G$  over  $E$  is given by a section  $q : S \rightarrow \hat{E}$ . Let further  $\mathcal{P} \rightarrow E \times_S \hat{E}$  be the Poincaré biextension of  $E$  and  $\hat{E}$  by  $\mathbb{G}_m$  : by [13], 10.2.13, a section  $s : S \rightarrow G$  of  $G/S$  lifting a section  $p : S \rightarrow E$  of  $E/S$  is entirely described by a trivialisation of the  $\mathbb{G}_m$ -torsor  $(p, q)^*\mathcal{P}$  over  $S$ . Assume now that  $E = E_0 \times S$  isoconstant, and admits complex multiplications, and let  $f : \hat{E}_0 \rightarrow E_0$  be a non-zero antisymmetric isogeny (i.e., identifying  $E_0$  and  $\hat{E}_0$ , a purely imaginary complex multiplication) which for simplicity, we here assume to be divisible by 2. Then,  $(f(q), q)^*\mathcal{P}$  is a trivial torsor in a canonical way, and the corresponding trivialization yields a well-defined section  $s = s(f)$  of  $G/S$  above  $p = f(q)$ . When  $G/S$  is semiconstant (which means : when  $q$  is not constant), any section  $s_R$  of  $G/S$  a non-zero multiple of which is of the form  $s(f)$  for some antisymmetric  $f$  will be called a Ribet section of  $G/S$ . So, on such a semi-abelian scheme  $G/S$ , there exists essentially only one Ribet section  $s_R$  (more precisely, all are linearly dependent over  $\mathbb{Z}$ ), and by [8] (see also [10], and Appendix I), its image  $W_R = s_R(S)$  meets  $G_{tor}$  infinitely often. It follows from Hindry's theorem (see Corollary 3 of §4) that the latter property characterizes Ribet sections among those sections of  $G/S$  which project to a section of  $E/S$  of the form  $p = f(q)$ .

## 1.2 The viewpoint of mixed Shimura varieties : Pink's conjecture

The consequences of the Main Theorem described in this section are discussed in [9], which we here summarize for the convenience of the reader.

Let  $X$  be a modular curve, parametrizing isomorphism classes of elliptic curves with some level structure, let  $\mathcal{E}$  be the universal elliptic scheme over  $X$ , with dual  $\hat{\mathcal{E}}$ , and let  $\mathcal{P}$  be the Poincaré bi-extension of  $\mathcal{E} \times_X \hat{\mathcal{E}}$  by  $\mathbb{G}_m$ . This is a mixed Shimura variety of dimension 4, which parametrizes points  $P$  on extensions  $G$  of elliptic curves  $E$  by  $\mathbb{G}_m$ . A point of  $\mathcal{P}(\mathbb{C})$  can be represented by a triple  $(E, G, P)$ , and is called special if the attached Mumford-Tate group is abelian, which is equivalent to requiring that  $E$  has complex multiplications, that  $G$  is an isotrivial extension, and that  $P$  is a

torsion point on  $G$ . Denote by  $\mathcal{P}_{sp}$  the set of special points of  $\mathcal{P}$ . Following [27], we further say that an irreducible subvariety of  $\mathcal{P}$  is special if it is a component of the Hecke orbit of a mixed Shimura subvariety of  $\mathcal{P}$ . Given any irreducible subvariety  $Z$  of  $\mathcal{P}$ , the intersection of all the special subvarieties of  $\mathcal{P}$  containing  $Z$  is called the special closure of  $Z$ . The special subvarieties of  $\mathcal{P}$  of dimension 0 are the special points; the special curves of  $\mathcal{P}$  are described below; for the full list, see [9], section 3.

**Corollary 1.** *Let  $W/\mathbb{Q}^{alg}$  be an irreducible closed algebraic curve in  $\mathcal{P}$ . Assume that  $W \cap \mathcal{P}_{sp}$  is infinite. Then,  $W$  is a special curve.*

To prove this corollary, we distinguish the various cases provided by the projection  $\varpi : \mathcal{P} \rightarrow X$  and its canonical section (rigidification)  $\sigma : X \rightarrow \mathcal{P}$ , whose image  $\sigma(X)$  is made up of points of the type  $(E, \mathbb{G}_m \times E, 0) \in \mathcal{P}$  :

- either the restriction of  $\varpi$  to  $W$  is dominant : the corollary then says that  $W$  lies in the Hecke orbit of the curve  $\sigma(X)$ . Indeed, up to Hecke transforms,  $\sigma(X)$  is the only one-dimensional (mixed, but actually pure) Shimura subvariety of  $\mathcal{P}$  dominating  $X$ .

In this case where  $\varpi|_W$  is dominant, the corollary follows not from our Main Theorem, but from André's theorem [2], p. 12, on the special points of the mixed Shimura variety  $\mathcal{E}$  (see also [22], Thm. 1.2).

- or  $\varpi(W)$  is a point  $x_0$  of  $X$ , necessarily of CM type. In particular,  $W$  lies in the fiber  $\mathcal{P}_0$  of  $\varpi$  above  $x_0$ . This fiber  $\mathcal{P}_0$  is a 3-dimensional mixed Shimura subvariety of  $\mathcal{P}$ , which can be identified with the Poincaré biextension of  $E_0 \times \hat{E}_0$  by  $\mathbb{G}_m$ , where  $E_0$  denotes an elliptic curve in the isomorphism class of  $x_0$ . An analysis of the generic Mumford-Tate group of  $\mathcal{P}_0$  as in [6], p. 52, shows that up to Hecke transforms, there are exactly four types of special curves in  $\mathcal{P}_0$  : the fiber  $(\mathbb{G}_m)_{x_0}$  above  $(0, 0)$  of the projection  $\mathcal{P}_0 \rightarrow (\mathcal{E} \times_X \hat{\mathcal{E}})_{x_0} = E_0 \times \hat{E}_0$  and the images  $\psi_B(B)$  of the elliptic curves  $B \subset E_0 \times \hat{E}_0$  passing through  $(0, 0)$  such that the  $\mathbb{G}_m$ -torsor  $\mathcal{P}_{0|B}$  is trivial, under the corresponding (unique) trivialization  $\psi_B : B \rightarrow \mathcal{P}_{0|B}$ . There are, up to torsion translates, three types of such elliptic curves  $B$  : the obvious ones  $E_0 \times 0$  and  $0 \times \hat{E}_0$  (whose images we denote by  $\psi(E_0 \times 0)$ ,  $\psi(0 \times \hat{E}_0)$ ), and the graphs of antisymmetric isogenies from  $E_0$  to  $\hat{E}_0$ , in which case  $\psi_B$  corresponds precisely to a Ribet section of the semi-abelian scheme  $\mathcal{G}_0/\hat{E}_0$  defined below.

Corollary 1 now follows from the Main Theorem, by interpreting  $\mathcal{P}_0/\hat{E}_0$  as the “universal” extension  $\mathcal{G}_0$  of  $E_0$  by  $\mathbb{G}_m$ , viewed as a group scheme over

the curve  $S := \hat{E}_0$ , so that  $\mathcal{P}_{sp} \cap \mathcal{P}_0 \subset (\mathcal{G}_0)_{tor}$ . More precisely, suppose that  $W$  dominates  $\hat{E}_0$ : then, it is the image of a multisection of  $\mathcal{G}_0/\hat{E}_0$ , and after a base extension, the theorem implies that  $W$  is the Ribet curve  $\psi_B(B)$ , or that it lies in a torsion translate of  $\mathbb{G}_{m/\hat{E}_0} = \mathbb{G}_m \times \hat{E}_0$ , where a new application of the theorem (or more simply, of Hindry's) shows that it must coincide with a Hecke transform of  $\mathbb{G}_m = (\mathbb{G}_m)_{x_0}$  or of  $\psi(0 \times \hat{E}_0)$ . By biduality (i.e. reverting the roles of  $\hat{E}_0$  and  $E_0$ ), the same argument applies if  $W$  dominates  $E_0$ . Finally, if  $W$  projects to a point of  $E_0 \times \hat{E}_0$ , then, this point must be torsion, and  $W$  lies in the Hecke orbit of  $(\mathbb{G}_m)_{x_0}$ .

Although insufficient in the presence of Ribet curves, the argument devised by Pink to relate the Manin-Mumford and the André-Oort settings often applies (see the proof of Theorems 5.7 and 6.3 of [27], and a probable addition to [10] for abelian schemes). In the present situation, one notes that given a point  $(E, G, P)$  in  $\mathcal{P}(\mathbb{C})$ , asking that it be special as in Corollary 1 gives 4 independent conditions, while merely asking that  $P$  be torsion on  $G$  as in the Main Theorem gives 2 conditions. Now, unlikely intersections for a curve  $W$  in  $\mathcal{P}$  precisely means studying its intersection with the union of the special subvarieties of  $\mathcal{P}$  of codimension  $\geq 2$  (i.e. of dimension  $\leq 2$ ), and according to Pink's Conjecture 1.2 of [27], when this intersection is infinite,  $W$  should lie in a special subvariety of dimension  $< 4$ , i.e. a proper one. Similarly, if  $W$  lies in the fiber  $\mathcal{P}_0$  of  $\mathcal{P}$  above a CM point  $x_0$  and meets infinitely many special curves of this 3-fold, then, it should lie in a special surface of the mixed Shimura variety  $\mathcal{P}_0$ . In these directions, our Main Theorem, combined with [2], p. 12, and with the relative version of Raynaud's theorem obtained in [19], implies :

**Corollary 2.** *Let  $W/\mathbb{Q}^{alg}$  be an irreducible curve in the mixed Shimura 4-fold  $\mathcal{P}$ , and let  $\delta_W$  be the dimension of the special closure of  $W$ .*

- i) Suppose that  $\delta_W = 4$ ; then, the intersection of  $W$  with the union of all the special surfaces of  $\mathcal{P}$  dominating  $X$  is finite;*
- ii) Suppose that  $\delta_W = 3$ ; then, the intersection of  $W$  with the union of all the special non-Ribet curves of  $\mathcal{P}$  is finite.*

The proof goes along the same lines as that of Corollary 1; see [9] for more details, and for a discussion on the gap between these corollaries and the full statement of Pink's conjecture 1.2 of [27], which would give a positive answer to :

**Question 2** *Let  $W/\mathbb{Q}^{alg}$  be an irreducible curve in the mixed Shimura 4-fold  $\mathcal{P}$ , and let  $\delta_W$  be the dimension of the special closure of  $W$ . Is the intersection of  $W$  with the union of all the special subvarieties of  $\mathcal{P}$  of dimension  $\leq \delta_W - 2$  then finite ?*

The case where  $\delta_W = 2$  is covered by Corollary 1 . The remaining cases would consist in disposing of the restrictions “dominating  $X$ ” and “non-Ribet” in Corollary 2. These problems are out of the scope of the present paper.

Although the Shimura view-point will not be pursued further, the Poincaré biextension, which has already appeared in the definition of Ribet sections, plays a role in the proof of the Main Theorem. See §5.3, Remark 3.(iii), §3.3, Footnote (3), and §6, Case **(SC2)**, where  $s$  is viewed as a section of  $\mathcal{P}$ , rather than of  $G$ . See also the sentence concluding §4.

### 1.3 Plan of the paper

- In the first sections (§§ 2, 3.1, 3.2), we give a set of notations, present the overall strategy of the proof, borrowed from [19] and based on the same set of preliminary lemmas : large Galois orbits, bounded heights, Pila-Wilkie upper bounds. The outcome is that for a certain real analytic surface  $\mathcal{S}$  in  $\mathbb{R}^4$  attached to the section  $s \in G(S)$ :

$W \cap G_{tor}$  **infinite**  $\Rightarrow \mathcal{S}$  **contains a semi-algebraic curve.**

- The program for completing the proof is sketched in §3.3, and can be summarized by the following two steps. Here  $\log_G(s)$  is a local logarithm of  $s$ , and  $F_{pq}$  is a certain Picard-Vessiot extension of  $K$  attached to  $G$  and to the projection  $p$  of  $s$  to  $E$ . Then, under a natural assumption on  $p$  (see Proposition 4),

$\mathcal{S}$  **contains a semi-algebraic curve**  $\Rightarrow \log_G(s)$  **is algebraic over  $F_{pq}$ .**

The proof is thereby reduced to a statement of algebraic independence, which forms the content of our “Main Lemma”, cf. middle of §3.3. Notice the similarity between the statements of this Main Lemma and the Main Theorem, making it apparent that up to translation by a constant section,

$\log_G(s)$  **is algebraic over  $F_{pq}$**   $\Rightarrow s$  **is Ribet or factors**

as is to be shown.

- As a warm up, we realize these two steps in Section 4 when  $G$  is an isotrivial extension. In Section 5, we go back to the general case  $G/S$ , prove the first step, and comment on the use of Picard-Vessiot extensions.



- Section 6 is devoted to the proof of the Main Lemma. As in §4, we appeal to results of Ax type [3] to treat isoconstant cases, and to Picard-Vessiot theory, in the style of André’s theorem [1], for the general case.
- Finally, Appendix I gives a concrete description of the local logarithm  $\log_G(s)$ , including an analytic presentation of the Ribet sections  $s_R$ , and a proof, independent from those of [8], [10], that they do contradict RMM (relative Manin-Mumford), while Appendix II is devoted to an application to polynomial Pell’s equations, following the method of [20].

## 2 Restatement of the Main Theorem

### 2.1 Introducing $q$ and $p$

We first repeat the setting of the introduction, with the help of the fundamental isomorphism  $Ext_S(E, \mathbb{G}_m) \simeq \hat{E}$  to describe the various cases to be studied.

So, let  $\mathbb{Q}^{alg}$  be the algebraic closure of  $\mathbb{Q}$  in  $\mathbb{C}$ . Extensions of scalars from  $\mathbb{Q}^{alg}$  to  $\mathbb{C}$  will be denoted by a lower index  $\mathbb{C}$ . Let  $S$  be an irreducible algebraic curve over  $\mathbb{Q}^{alg}$ , whose generic point we denote by  $\lambda$ , and let  $K = \mathbb{Q}^{alg}(S) = \mathbb{Q}^{alg}(\lambda)$ , so  $K_{\mathbb{C}} := \mathbb{C}(S_{\mathbb{C}}) = \mathbb{C}(\lambda)$ . We use the notation  $\bar{\lambda}$  for the closed points of  $S_{\mathbb{C}}$ , i.e.  $\bar{\lambda} \in S(\mathbb{C})$ . Let  $G/S$  be a semi-abelian scheme over  $S$  of relative dimension 2 and toric rank 1. Making Question 1 more precise, we write  $G_{tor}$  for the union of all the torsion points of the various fibers of  $G_{\mathbb{C}} \rightarrow S_{\mathbb{C}}$ , i.e.  $G_{tor} = \cup_{\bar{\lambda} \in S(\mathbb{C})} (G_{\bar{\lambda}})_{tor} \subset G(\mathbb{C})$ , where  $G_{\bar{\lambda}}$  denotes the fiber of  $G$  above  $\bar{\lambda}$ . This set  $G_{tor}$  is also the set of values at all  $\bar{\lambda} \in S(\mathbb{C})$  of the various torsion-sections of  $G_{\mathbb{C}} \rightarrow S_{\mathbb{C}}$ . Let finally  $s : S \rightarrow G$  be a section of  $G/S$  defined over  $\mathbb{Q}^{alg}$ , giving a point  $s(\lambda) \in G_{\lambda}(K)$  at the generic point of  $S$ , and closed points  $s(\bar{\lambda}) \in G_{\bar{\lambda}}(\mathbb{C})$  at the  $\bar{\lambda}$ ’s in  $S(\mathbb{C})$ .

In the description which follow, we may have to withdraw some points of  $S$ , or replace  $S$  by a finite cover, but will still denote by  $S$  the resulting curve. After a base extension, the group scheme  $G/S$  can be presented in a unique way as an extension

$$0 \longrightarrow \mathbb{G}_{m/S} \longrightarrow G \xrightarrow{\pi} E \longrightarrow 0$$

by  $\mathbb{G}_{m/S} = \mathbb{G}_m \times S$  of an elliptic scheme  $E/S$  over  $S$ . We denote by  $\pi : G \rightarrow E$  the corresponding  $S$ -morphism.

The extension  $G$  is parametrized by a section

$$q \in \hat{E}(S)$$

of the dual elliptic scheme  $\hat{E}/S$ . We write

$$p = \pi \circ s \in E(S)$$

for the projection to  $E$  of the section  $s$ . Finally, we denote by  $k \subset \mathbb{Q}^{alg}$  the number field over which  $S$  is defined, and assume without loss of generality that the generic fibers of  $G$ , hence of  $q$ , and of  $s$ , hence of  $p$ , are defined over  $k(S) = k(\lambda)$ .

Since the algebraic curve  $W = s(S) \subset G$  is the image of a section, the minimal strict subgroup schemes  $H$  of  $G$  which may contain  $W$  are flat over  $S$ , with relative dimension 1. These can be described as follows :

- either  $q$  has infinite order in  $\hat{E}(S)$  : then,  $G/S$  is a non isotrivial extension, and  $H \subsetneq G$  is contained in a finite union of torsion translates of  $\mathbb{G}_{m/S}$ ; in particular,  $\pi(H)$  is finite over  $S$ , and  $H$  can contain  $W$  only if  $p = \pi(s)$  is a torsion section of  $E/S$ .

- or  $q$  has finite order : in this case,  $G$  is isogeneous to the direct product  $\mathbb{G}_{m/S} \times_S E$ . Since the answer to the question is invariant under isogenies, we can then assume that  $G$  is this direct product, i.e. that  $q = 0$ . Strangely enough, this (easy) case of Question 1 does not seem to have been written up yet. We present it in §4. The answer (cf. Theorem 1 below) is a corollary of Hindry's theorem when  $E/S$  is isoconstant, since  $G/S$  too is then isoconstant; in an apparently paradoxical way, we will use it to characterize the Ribet sections of semi-constant extensions, cf. Corollary 3.

So, from now on and apart from §4, we could assume that  $q$  has infinite order in  $\hat{E}(S)$ , i.e. that  $G$  is a non-isotrivial extension. However, the only hypothesis we will need in our general study of §3.3 and §5 concerns the section  $p = \pi(s)$  of  $E/S$ , see next subsection.

## 2.2 Isoconstant issues

In general, given our curve  $S/\mathbb{Q}^{alg}$ , we say that a scheme  $X/S$  is isoconstant if there exists a finite cover  $S' \rightarrow S$  and a scheme  $X_0/\mathbb{Q}^{alg}$  such that  $X_{S'} = X \times_S S'$  is isomorphic over  $S'$  to the constant scheme  $X_0 \times_{\mathbb{Q}^{alg}} S'$ . We then say that a section  $x$  of  $X/S$  is *constant* if after a base change  $S'/S$  which

makes  $X$  constant, the section of  $X_{S'}/S'$  which  $x$  defines comes from the constant part  $X_0$  of  $X_{S'}$ . This notion is indeed independent of the choice of  $S'$  (and often does not even require the base change  $S'/S$ ). See Footnote (4) of §4 below for further conventions in the isoconstant cases.

In these conditions, the hypothesis just announced about  $p$  reads :

**$p \in E(S)$  is not a torsion section, and is not constant if  $E/S$  is isoconstant,**

and will be abbreviated as “the section  $p$  is not torsion, nor constant”; in terms to be described in §5.3 and §6, it is better expressed as :

**$p \in E(S)$  does not lie in the Manin kernel  $E^\sharp$  of  $E$ .**

The relation to the Main Theorem is as follows. If  $p$  is a torsion section, then, a torsion translate of  $s \in G(S)$  lies in  $\mathbb{G}_m$ , so  $s$  satisfies Condition (ii) of the Main Theorem. And if  $p = p_0$  is a constant (and not torsion) section, then,  $p(S) = \{p_0\} \times S$  does not meet  $E_{tor}$  at all, so  $s(S) \cap G_{tor}$  is empty. In other words, the Main Theorem is trivial in each of these cases.

A precision on the expression “ $G$  is semi-constant” is now in order : it appears only if  $E/S$  is isoconstant, and means that there exists a finite cover  $S' \rightarrow S$  such that on the one hand, there exists an elliptic curve  $E_0/\mathbb{Q}^{alg}$  such that the pull-back of  $E/S$  to  $S'$  is isomorphic over  $S'$  to  $E_0 \times S'$ , and that on the other hand, the section  $q' \in \hat{E}'(S')$  which  $q$  defines is given by a section of  $\hat{E}_0 \times_{\mathbb{Q}^{alg}} S$  which does *not* come from  $\hat{E}_0(\mathbb{Q}^{alg})$ . Since the answer to our question is invariant under finite base extension of  $S$ , we will assume in this case that  $E/S$  is already constant, i.e.  $E = E_0 \times S$ , and that  $q \in \hat{E}_0(S) \setminus \hat{E}_0(\mathbb{Q}^{alg})$ ; indeed, as just said above, the second condition is then valid for any  $S'/S$ . Notice that the condition that  $q$  be non-constant forces it to be of infinite order. Consequently, semi-constant extensions are automatically both non isoconstant and non-isotrivial. On the other hand, if  $q \in \hat{E}_0(\mathbb{Q}^{alg})$  is constant, we are in the purely constant case of [16] already discussed in the Introduction.

**Remark 1 (traces and images)** : let  $\mathcal{E}_0/\mathbb{Q}^{alg}$  denote the constant part ( $K/\mathbb{Q}^{alg}$ -trace) of  $E/S$ . An innocuous base extension allows us to assume that if  $E/S$  is isoconstant, then  $\mathcal{E}_0 \neq \{0\}$  (and we then set  $\mathcal{E}_0 := E_0$ ), and  $E/S$  is actually isomorphic to  $E_0 \times S$ . Denote by  $G_0/\mathbb{Q}^{alg}$  the constant part ( $K/\mathbb{Q}^{alg}$ -trace) of  $G/S$ , and by  $G^0/\mathbb{Q}^{alg}$  the maximal constant quotient ( $K/\mathbb{Q}^{alg}$ -image) of  $G/S$ . Then :

-  $G/S$  is an isotrivial extension if and only if  $q$  is a torsion section of  $\hat{E}/S$ . In this case, an isogeny allows us to assume that  $q = 0$ , i.e.  $G = \mathbb{G}_m \times E$ .

We then have  $G_0 = G^0 = \mathbb{G}_m \times \mathcal{E}_0$ , and  $G/S$  is isoconstant if and only if  $E/S$  is isoconstant.

- assume now that  $G/S$  is not an isotrivial extension, and that  $E/S$ , hence  $\hat{E}/S$ , is not isoconstant. Then,  $G_0 = \mathbb{G}_m$ , while  $G^0 = \mathcal{E}_0 = \{0\}$ .

- finally, assume that  $G/S$  is not an isotrivial extension, but  $E = E_0 \times S$ , hence  $\hat{E} = \hat{E}_0 \times S$ , is (iso)constant. Then either  $q$  is a non constant section of  $\hat{E}/S$ , in which case  $G$  is semi-constant, and we have  $G_0 = \mathbb{G}_m, G^0 = \mathcal{E}_0 = E_0$ ; or  $q$  is constant, in which case  $G^0 = G_0 \in \text{Ext}_{\mathbb{Q}^{alg}}(E_0, \mathbb{G}_m)$ , and  $G = G_0 \times S$  is itself constant.

For the sake of brevity, we will henceforth say that a section of a group scheme  $H/S$  is (iso)constant if it is a constant section of the constant part  $H_0 \times S$  of  $H/S$ .

### 2.3 Antisymmetric relations and restatement

The answer to Question 1 of the Introduction, as well as the proofs, depend on possible relations between  $p$  and  $q$ . For ease of notations, we fix a principal polarization  $\psi : \hat{E} \simeq E$  of the elliptic scheme, and allow ourselves to identify  $q$  with its image  $\psi(q) \in E(S)$ . Also, we denote by  $\mathcal{O}$  the ring of endomorphisms of  $E$ . If  $E/S$  is not isoconstant,  $\mathcal{O}$  reduces to  $\mathbb{Z}$ . Otherwise,  $\mathcal{O}$  may contain complex multiplications, and we say that the non-torsion and non-constant sections  $p$  and  $q$  are *antisymmetrically related* if there exists  $\alpha \in \mathcal{O} \otimes \mathbb{Q}$  with  $\bar{\alpha} = -\alpha$  such that  $q = \alpha p$  in  $E(K)$  modulo torsion.

Notice that we reserve this expression for sections  $p, q$  which are non torsion (and non constant). Therefore, an antisymmetric relation between  $p$  and  $q$  necessarily involves a non-zero imaginary  $\alpha$ , hence  $\mathcal{O} \neq \mathbb{Z}$ , forcing  $E/S$  to be isoconstant. And since  $q$  is not constant, the corresponding semi-abelian scheme  $G = G_q$  is then semiconstant, and admits a Ribet section  $s_R$  projecting to  $p \in E(S)$ .

For any positive integer  $m$ , we set :

$$S_m^G = \{\bar{\lambda} \in S(\mathbb{C}), s(\bar{\lambda}) \text{ has order } m \text{ in } G_{\bar{\lambda}}(\mathbb{C})\},$$

$$S_m^E = \{\bar{\lambda} \in S(\mathbb{C}), p(\bar{\lambda}) \text{ has order } m \text{ in } E_{\bar{\lambda}}(\mathbb{C})\},$$

$$S_\infty^G = \cup_{m \in \mathbb{Z}_{>0}} S_m^G \simeq W \cap G_{tor}, \quad S_\infty^E = \cup_{m \in \mathbb{Z}_{>0}} S_m^E \simeq \pi(W) \cap E_{tor},$$

where the indicated bijections are induced by  $S \simeq s(S) = W$ ,  $S \simeq p(S) = \pi(W)$ . Clearly,  $\cup_{k|m} S_k^G \subset \cup_{k|m} S_k^E$  for all  $m$ 's, and the points of  $S_\infty^G$  can be

described as those points of  $\pi(W) \cap E_{tor}$  (“likely intersections”) that lift to points of  $W \cap G_{tor}$  (“unlikely intersections”).

Our Main Theorem can then be divided into the following three results. We first consider the case when  $q$  is torsion, which reduces after an isogeny to the case  $q = 0$ .

**Theorem 1.** *Let  $E/S$  be an elliptic scheme over the curve  $S/\mathbb{Q}^{alg}$ , and let  $G = \mathbb{G}_m \times E$  be the trivial extension of  $E/S$  by  $\mathbb{G}_{m/S}$ . Let further  $s : S \rightarrow G$  be a section of  $G/S$ , with image  $W = s(S)$ , such that  $p = \pi(s)$  has infinite order in  $E(S)$ . Then,  $S_\infty^G$  is finite (in other words,  $W \cap G_{tor}$  is finite) as soon as*

(o) *no multiple of  $s$  by a positive integer factors through  $E/S$  (i.e. the projection of  $s$  to the  $\mathbb{G}_m$ -factor of  $G$  is not a root of unity).*

The case when  $q$  is not torsion can be restated as follows.

**Theorem 2.** *Let  $E/S$  be an elliptic scheme over the curve  $S/\mathbb{Q}^{alg}$ , and let  $G/S$  be a non-isotrivial extension of  $E/S$  by  $\mathbb{G}_{m/S}$ , i.e. parametrized by a section  $q \in \hat{E}(S) \simeq E(S)$  of infinite order. Let further  $s : S \rightarrow G$  be a section of  $G/S$ , with image  $W = s(S)$ , such that  $p = \pi(s)$  has infinite order in  $E(S)$ . Then,  $S_\infty^G$  is finite (in other words,  $W \cap G_{tor}$  is finite) in each of the following cases :*

- (i)  *$E/S$  is not isoconstant;*
- (ii)  *$E/S$  is isoconstant, and  $p$  and  $q$  are not antisymmetrically related;*
- (iii)  *$E/S$  is isoconstant,  $p$  and  $q$  are non-constant antisymmetrically related sections, and no multiple of  $s$  is a Ribet section.*

For the sake of symmetry, we recall that in these two theorems, the hypothesis that  $p$  is not torsion is equivalent to requiring that no multiple of  $s$  by a positive integer factors through  $\mathbb{G}_{m/S}$ .

Since Ribet sections exist only in Case (iii) of Theorem 2, the conjunction of Theorems 1 and 2 is equivalent to Part (A) of the Main Theorem, giving necessary conditions for  $W \cap G_{tor} \simeq S_\infty^G$  to be infinite. That these conditions are (essentially) sufficient, i.e. that Part (B) holds true, is dealt with by the following statement, which we prove right now.

**Theorem 3.** *Let  $E/S$  be an elliptic scheme over the curve  $S/\mathbb{Q}^{alg}$ , and let  $G/S$  be an extension of  $E/S$  by  $\mathbb{G}_{m/S}$ . Let further  $s : S \rightarrow G$  be a section of  $G/S$ . Then,*

- i) if  $s$  is a Ribet section,  $S_\infty^G$  is infinite, and equal to  $S_\infty^E$ ;*
- ii) if  $s$  is a torsion section, then  $S_\infty^G = S(\mathbb{C})$ ;*
- iii) if  $s$  is a non-torsion section factoring through a strict subgroup scheme  $H/S$  of  $G/S$ , then  $S_\infty^G$  is empty if  $s$  is an isoconstant section of  $H/S$ , and infinite but strictly contained in  $S(\mathbb{Q}^{alg})$  if  $s$  is not isoconstant.*

*Proof :* (i) is proved in [8], [10], see also Appendix I below. The second statement is clear. As for (iii), this is an easy statement if the connected component of  $H$  is  $\mathbb{G}_m$ . If (for an isotrivial  $G$ ), it is isogenous to  $E$ , the isoconstant case is again clear, while the non-isoconstant one follows from “torsion values for a single point” as in [31], cf. Proposition 1.(iv) below.

So, we can now concentrate on Theorems 1 and 2.

### 3 The overall strategy

Our strategy will be exactly the same as in [19].

#### 3.1 Algebraic lower bounds

In this section, we denote by  $k$  a number field over which the algebraic curve  $S$ , the group scheme  $G/S$  and its section  $s$  are defined. We fix an embedding of  $S$  in a projective space over  $k$ , and denote by  $H$  the corresponding height on the set  $S(\mathbb{Q}^{alg})$  of algebraic points of  $S$ . We then have :

**Proposition 1.** *Let  $E/S$  be an elliptic scheme, and let  $p : S \rightarrow E$  be a section of  $E/S$  of infinite order. There exist positive real numbers  $C, C'$  depending only on  $S/k, E/S$  and  $p$ , with the following properties. Let  $\bar{\lambda} \in S(\mathbb{C})$  be such that  $p(\bar{\lambda})$  is a torsion point of  $E_{\bar{\lambda}}(\mathbb{C})$ , i.e.  $\bar{\lambda} \in S_\infty^E$ . Then,*

- i) the point  $\bar{\lambda}$  lies in  $S(\mathbb{Q}^{alg})$ , i.e. the field  $k(\bar{\lambda})$  is an algebraic extension of  $k$ ;*
- ii) the height  $H(\bar{\lambda})$  of  $\bar{\lambda}$  is bounded from above by  $C$ ;*
- iii) if  $n \geq 1$  denotes the order of  $p(\bar{\lambda})$ , then  $[k(\bar{\lambda}) : k] \geq C'n^{1/3}$ .*
- iv) the set  $S_\infty^E$  is infinite (assuming that if  $E/S$  is isoconstant,  $p$  is not constant).*

*Proof :* *i, ii, iii)* In the non-isoconstant case, one can reduce to the Legendre curve, where all is already written in [19], [20], [31], based on diophantine results of Silverman, David and the second author. Notice that the upper

bound (ii) on  $H(\bar{\lambda})$  is needed to deduce the degree lower bound (iii). In the isoconstant case, the proof is easier as (ii) is not needed, and one can sharpen the lower bound (iii) to  $n^2$  in the non-CM case (non effective), resp.  $n/\log n$  in the CM case. But as usual, any positive power of  $n$  will do.

*iv)* This is the issue of “torsion values for a single point”, an analytic proof of which is given in [31], p. 92, in the non-isoconstant case. If  $E/S$  is isoconstant, the second case of this analytic proof does not occur, since we assume that  $p$  is not constant.

**Corollary.** *Hypotheses and the notation  $C'$  being as in Proposition 1, let  $G/S$  be an extension of  $E/S$  by  $\mathbb{G}_{m/S}$ , and let  $s$  be a section of  $G/S$  lifting the non-torsion section  $p$  of  $E/S$ . Assume that  $\bar{\lambda} \in S_\infty^E$  actually lies in  $S_\infty^G$ , and let  $m$  be the order of the torsion point  $s(\bar{\lambda}) \in G_{\bar{\lambda}}(\mathbb{Q}^{alg})$ . Then,  $[k(\bar{\lambda}) : k] \geq C'm^{1/6}$ .*

*Proof:* if  $s(\bar{\lambda})$  has precise order  $m$  in  $G_{\bar{\lambda}}$  and projects to a point  $p(\bar{\lambda})$  of order  $n|m$  on  $E_{\bar{\lambda}}$ , then,  $ns(\bar{\lambda})$  is a primitive  $m/n$ -th root of unity, so has degree  $\gg (m/n)^{1-\varepsilon}$  over  $k$ , and we can assume that this is larger than  $C'(m/n)^{1/3}$  (assume  $C' < [k : \mathbb{Q}]^{-1}$ ). Since the fields of definition of  $ns(\bar{\lambda})$  and of  $p(\bar{\lambda})$  are contained in  $k(\bar{\lambda})$ , we get  $[k(\bar{\lambda}) : k] \geq C'(sup(n, m/n))^{1/3} \geq C'm^{1/6}$ .

The conclusion of this first step is summarized by the implications :

$$\forall \bar{\lambda} \in S(\mathbb{C}), \bar{\lambda} \in S_\infty^G \Rightarrow \bar{\lambda} \in S(\mathbb{Q}^{alg}) \text{ and } H(\bar{\lambda}) \leq C,$$

and

$$\forall \bar{\lambda} \in S(\mathbb{Q}^{alg}), \forall m \geq 1, \bar{\lambda} \in S_m^G \Rightarrow [k(\bar{\lambda}) : k] \geq C'm^{1/6}.$$

In particular, if  $W = s(S)$  contains a point  $w = s(\bar{\lambda})$  of order  $m$  (w.r.t. the group law of its fiber  $G_{\bar{\lambda}}$ ), then  $W$  contains at least  $C'm^{1/6}$  points of order  $m$  (w.r.t. the group laws of their respective fibers) : indeed, since  $W$  is defined over  $k$ , it contains the orbit of  $w$  under  $Gal(\mathbb{Q}^{alg}/k)$ . However, we will need a sharper version of this statement, involving the archimedean sizes of the conjugates of  $\bar{\lambda}$ , and the upper bound on  $H(\bar{\lambda})$  will again come to help at this stage.

## 3.2 Transcendental upper bounds

The next step is based on the following theorem of Pila. For the involved definitions and a short history on this type of results, leading to [21] and the accompanying references, we refer to [31], Remark 3.1.1, and to the Amer.

J. Math. version of [19]). Dimensions here refer to real dimensions. For any  $m \in \mathbb{Z}_{>0}$ , we set  $\mathbb{Q}_m = \frac{1}{m}\mathbb{Z} \subset \mathbb{Q}$ .

**Proposition 2.** *Let  $\mathcal{S}$  be a naive-compact-2-(dimensional)-analytic subset of an affine space  $\mathbb{R}^d$ . Assume that no semi-algebraic curve of  $\mathbb{R}^d$  is contained in  $\mathcal{S}$ . For any  $\varepsilon > 0$ , there exists a real number  $c = c(\mathcal{S}, \varepsilon) > 0$  with the following property. For each positive integer  $m$ , the set  $\mathcal{S} \cap \mathbb{Q}_m^d$  contains at most  $c m^\varepsilon$  points.*

*Proof :* see [19], Lemma 1.

In §3.3, we will give a precise description of the real surfaces  $\mathcal{S}$  to which Proposition 2 is to be applied. See the following §4.1 for a more easily recognizable form when  $G = \mathbb{G}_m \times E$ . Roughly speaking,

$$\mathcal{S} \sim \log_G^B(W) \subset (\mathbb{Z} \otimes \mathbb{C}) \oplus (\mathbb{Z}^2 \otimes \mathbb{R}) \simeq \mathbb{R}^4$$

is the set of logarithms of the various points  $s(\bar{\lambda})$ , when  $\bar{\lambda}$  runs through  $S(\mathbb{C})$ , but we express these logarithms in terms of a *basis attached to the  $\mathbb{Z}$ -local system of periods  $\Pi_G$* , i.e. in Betti terms rather than in the de Rham viewpoint provided by the Lie algebra, hence the upper index  $B$  above. In this basis, the logarithms of the torsion points  $s(\bar{\lambda})$  of order  $m$  are represented by vectors with coordinates in  $\mathbb{Q}_m$ , and so  $\log_G^B(s(S_m^G)) \subset \mathbb{Q}_m^4$ . Thanks to the “zero estimate” discussed at the end of the present subsection, which compares the **graph**  $\tilde{\mathcal{S}} \subset S \times \mathbb{R}^4$  of  $\log_G^B \circ s : S \rightarrow \mathbb{R}^4$  with its projection  $\mathcal{S}$  in  $\mathbb{R}^4$ , Proposition 2 then implies (with a proviso to be explained below for the first conclusion)

- either that for any positive integer  $m$ ,  $\exp_G(\tilde{\mathcal{S}}) \sim s(S) = W$  contains at most  $c m^\varepsilon$  points of order  $m$  w.r.t the group law of their respective fibers;
- or that  $\mathcal{S}$  contains a real semi-algebraic curve, where algebraicity refers to the real affine space  $\mathbb{R}^4$  associated to the above mentioned basis.

In §§4 and 5-6, we will prove that in all cases considered in Theorems 1 and 2, the surface  $\mathcal{S}$  contains no semi-algebraic curve. The first conclusion must then hold true. Combined with the conclusion of §3.1, this implies that the orders  $m$  of the torsion points lying on  $W$  are uniformly bounded, and so, there exists a positive integer  $N = N(k, S, G, s)$  such that  $W \cap G_{tor} \subset \bigcup_{\bar{\lambda} \in S(\mathbb{Q}^{alg})} G_{\bar{\lambda}}[N]$ . The latter set is the union of the values at all  $\bar{\lambda} \in S(\mathbb{Q}^{alg})$  of the torsion sections of  $G/S$  of order dividing  $N$ , which form a finite union of curves in  $G$ . As soon as  $p$  is not a torsion section, neither is  $s$  and  $W$



intersects this finite union in a finite number of points. Hence,  $W \cap G_{tor}$  is finite, and this concludes the proof of Theorems 1 and 2. (NB : as done in [19], this conclusion can be reached in a faster way via the inequalities  $m^{1/6} \ll [k(\bar{\lambda}) : k] \ll \sharp(S_m^G) \ll m^\epsilon$ ,  $H(\bar{\lambda}) \ll 1$ , and the Northcott property.)

However, two points must be modified for the above discussion to hold.

- We need a uniform determination of the logarithms of the points  $s(\bar{\lambda})$ , and this requires fixing from the start a (any) simply connected pointed subset  $(\Lambda, \bar{\lambda}_0)$  of the Riemann surface  $S^{an}$  attached to  $S(\mathbb{C})$ ; in particular, our surface

$$\mathcal{S} = \mathcal{S}_\Lambda := \log_{G, \bar{\lambda}_0}^B(s(\Lambda)),$$

and the graph  $\tilde{\mathcal{S}}_\Lambda \subset \Lambda \times \mathbb{R}^4$  of  $\log_{G, \bar{\lambda}_0}^B \circ s : \Lambda \rightarrow \mathbb{R}^4$  will depend on a choice of  $\Lambda$ . Furthermore, the surfaces  $\mathcal{S}$  studied by Proposition 2 must be compact, so,  $\Lambda$  too must be compact. Consequently,  $\exp_G(\tilde{\mathcal{S}}_\Lambda) = s(\Lambda) \subset W(\mathbb{C})$  is truly smaller than  $W(\mathbb{C})$ , and the desired “first conclusion” is reached in a slightly different way : as in [19] (see Lemma 6.2 of AJM version, Lemma 8.2 of Math. Ann.), one first attaches to the height bound  $C$  a finite union  $\Lambda_C$  of simply-connected compact pointed sets  $(\Lambda_i, \bar{\lambda}_i)$  such that for any point  $\bar{\lambda} \in S(\mathbb{Q}^{alg})$  of sufficiently large degree with  $H(\bar{\lambda}) \leq C$ , a positive proportion (say half, i.e. independent of  $\bar{\lambda}$ ) of the conjugates of  $\bar{\lambda}$  over  $k$  lies in  $\Lambda_C$ <sup>1</sup>. Letting  $\tilde{\mathcal{S}}_{\Lambda_C}$  be the finite union of the graphs of the maps  $\log_{G, \bar{\lambda}_i}^B \circ s$ , we deduce that for any  $\bar{\lambda} \in S_m^G$ , a similar positive proportion of the orbit of  $s(\bar{\lambda}) = w$  under  $Gal(\mathbb{Q}^{alg}/k)$  lies in  $\exp_G(\tilde{\mathcal{S}}_{\Lambda_C}) = s(\Lambda_C) \subset W$ . Proposition 2, combined with the subsequent zero estimate, then implies that this orbit has at most  $c'm^\epsilon$  points, and one can conclude as above (or via the Northcott

---

<sup>1</sup> In the present case, let us first remove from  $S$  the finite set consisting of the points of bad reduction and those where the section is not defined (or any finitely many ones that may cause trouble along the way, possibly none...). Now remove “small” open disks around each of these points; what remains is a compact set in  $S$ . We want them small enough so that at most half of the conjugates of the relevant  $\bar{\lambda}$  fall in their union: this may be achieved because  $\bar{\lambda}$  has bounded height. In fact, if “many” conjugates fall into a same small disk, then the corresponding contribution to the height is too big. In turn this follows for instance on looking at the difference  $f(\bar{\lambda}) - f(\bar{\lambda}_0)$  where  $\bar{\lambda}_0$  is the center of the disk and  $f$  is a suitable nonconstant coordinate on  $S$ . Using the coordinate reduces the verification to the case of algebraic numbers (rather than algebraic points). Having chosen these small enough disks, we cover the said compact set with finitely many simply connected domains where the logs are locally defined.

property on  $\bar{\lambda}$ ). In what follows, we fix one of the  $\Lambda_i$ 's, call it  $\Lambda$  and write  $\log_G^B \circ s$  for  $\log_{G, \bar{\lambda}_i}^B \circ s$ .

• Proposition 2 provides an upper bound for the image in  $\mathcal{S} \cap \mathbb{Q}_m^4 \subset \mathbb{R}^4$  of  $S_m^G \cap \Lambda$  under  $\log_G^B \circ s : \Lambda \rightarrow \mathbb{R}^4$ , while we need an upper bound for  $S_m^G \cap \Lambda$  itself. In other words, we must show that not too many points  $\bar{\lambda}$  of  $S_m^G$  can be sent by  $\log_G^B \circ s$  onto the same point of  $\mathbb{Q}_m^4$ . Clearly, it suffices to show that the projection

$$u(\lambda) := d\pi(\log_G(s(\lambda))) = \log_E(\pi(s(\lambda))) = \log_E(p(\lambda))$$

of  $\log_G(s)$  under the differential of  $\pi : G \rightarrow E$  satisfies this separation property. The gap can now be filled by appealing to the “zero estimate” of [19], Lemma 7.1 of AJM, or Lemma 9.1 of Math. Ann., as follows. The (elliptic) Betti notations introduced here will be developed in §3.3 and §4.

Let  $\Lambda$  be a compact and simply connected subset of the Riemann surface  $S(\mathbb{C})$ , which, without loss of generality, we may assume to be homeomorphic to a closed disk in  $\mathbb{C}$ . We recall that given an analytic sheaf  $\mathcal{F}$  on  $S(\mathbb{C})$ , a section  $\sigma \in \mathcal{F}(\Lambda)$  of  $\mathcal{F}$  over  $\Lambda$  is by definition analytic on a neighbourhood of  $\Lambda$  in  $S(\mathbb{C})$ . Let now  $E/S$  be an elliptic scheme over  $S$ , and let  $\omega_1(\lambda), \omega_2(\lambda)$  be the analytic functions on a neighbourhood of  $\Lambda$  in  $S(\mathbb{C})$  expressing its periods relative to a given global differential form of the first kind on  $E/S$ . Fix a determination  $\log_E$  of the corresponding elliptic logarithm on  $E(\Lambda)$ . For any analytic section  $\mathbf{p} \in E(\Lambda)$ , there then exists unique real analytic functions  $\beta_1, \beta_2 : \Lambda \rightarrow \mathbb{R}$  such that  $\log_E(\mathbf{p}(\lambda)) = \beta_1(\lambda)\omega_1(\lambda) + \beta_2(\lambda)\omega_2(\lambda)$ . We call  $\{\beta_1, \beta_2\}$  the Betti coordinates of  $\mathbf{p}$ , and (extending a well-known notion for the Legendre family) say that  $\mathbf{p}$  is a *Picard-Painlevé section* of  $E/\Lambda$  if its Betti coordinates  $\beta_1, \beta_2$  are *constant*. We then have :

**Proposition 3. (Zero estimate).** *Let  $E/S$  and the simply-connected compact subset  $\Lambda$  be as above, and let  $p$  be a regular section of  $E/S$ . Assume that  $p$  is not a torsion section, and if  $E/S$  is isoconstant, that it is not a constant section. There exists an integer  $C''$  depending only on  $E/S, \Lambda$  and  $p$  such that for any Picard-Painlevé section  $\mathbf{p}$  of  $E/\Lambda$ , the set*

$$\{\bar{\lambda} \in \Lambda, \log_E(p(\bar{\lambda})) = \log_E(\mathbf{p}(\bar{\lambda}))\}$$

*has at most  $C''$  elements.*

In other words, if we set  $u(\lambda) = \log_E(p(\lambda)) = b_1(\lambda)\omega_1(\lambda) + b_2(\lambda)\omega_2(\lambda)$ , then for any real numbers  $\beta_1, \beta_2$ , the equation  $u(\bar{\lambda}) = \beta_1\omega_1(\bar{\lambda}) + \beta_2\omega_2(\bar{\lambda})$

has at most  $C''$  solutions  $\bar{\lambda} \in \Lambda$ . So, the statement above is just a fancy translation of [19], Amer. JM, p. 12, and follows from Lemma 7.1 there in exactly the same way if  $E/S$  is not isoconstant. The constant case is even easier. Notice that we need Proposition 3 only for  $\beta_1, \beta_2$  running in  $\mathbb{Q}$ , i.e. for torsion Picard-Painlevé sections  $\mathbf{p}$ , and that the Painlevé equation may bring a new view point on the computation of the bound  $C''$ .

In our applications,  $p$  is not a torsion section of  $E/S$ . And in the isoconstant case, we have assumed without loss of generality that  $p$  is not constant. So, the map  $\log_E^B \circ p$  separates the points of  $\Lambda$  up to the bounded error  $C''$ ; a fortiori, so does its lift  $\log_G^B \circ s$ , and the gap between its image  $\mathcal{S}$  and its graph  $\hat{\mathcal{S}}$  is filled.

### 3.3 What remains to be done

In view of the previous discussion, the proof of Theorems 1 and 2 is now reduced to defining the real surface  $\mathcal{S}$  properly, and to showing that under each of their hypotheses,  $\mathcal{S}$  contains no semi-algebraic curve. This is dealt with as follows.

#### The real surface $\mathcal{S}$

Fix a simply connected and compact subset  $\Lambda \in S(\mathbb{C})$ , homeomorphic to a closed disk, as well as a point  $\bar{\lambda}_0$  in  $\Lambda$ , and a point  $U_0$  in  $Lie(G_{\bar{\lambda}_0}(\mathbb{C}))$  such that  $\exp_{G_{\bar{\lambda}_0}}(U_0) = s(\bar{\lambda}_0) \in G_{\bar{\lambda}_0}(\mathbb{C})$ . We henceforth denote by  $\lambda$  the general element<sup>2</sup> of  $\Lambda$ , and (sometimes) by an upper index  $^{an}$  the analytic objects over the Riemann surface  $S^{an}$  attached to our schemes over  $S$ .

We first repeat the definition of the real surface  $\mathcal{S}$  in more precise terms. The group scheme  $G/S$  defines an analytic family  $G^{an}$  of Lie groups over the Riemann surface  $S^{an}$ . Similarly, its relative Lie algebra  $(LieG)/S$  defines an

---

<sup>2</sup> A remark may be in order about the meaning of the notation  $\lambda$ . In the first paragraphs, it represented the generic point of  $S_{\mathbb{C}}$ , i.e. we set  $\mathbb{C}(S) = \mathbb{C}(\lambda)$  (by the way, from now on, we are over  $\mathbb{C}$ , so, dropping the lower index  $\mathbb{C}$ , we will write  $K = \mathbb{C}(S)$ ). But it now represents the “general element” of the simply connected domain  $\Lambda \subset S^{an}$ , which can have many analytic automorphisms. It is understood that we here consider only a global  $\lambda$ . Such a  $\lambda$  may require several algebraically dependent parameters to be expressed. Of course, when  $\Lambda$  is small enough, we can work with a chart  $t$  on  $\Lambda$  such that  $\mathbb{C}(\lambda)$  is an algebraic extension of  $\mathbb{C}(t)$ . The results of functional algebraic independence we appeal to do not require such reduction.

analytic vector bundle  $LieG^{an}$  over  $S^{an}$ , of rank 2. The  $\mathbb{Z}$ -local system of periods of  $G^{an}/\Lambda$  is the kernel of the exponential exact sequence

$$0 \longrightarrow \Pi_G \longrightarrow LieG^{an} \xrightarrow{exp_G} G^{an} \longrightarrow 0 ,$$

over  $S^{an}$ . Its sections over  $\Lambda$  form a  $\mathbb{Z}$ -module  $\Pi_G(\Lambda) \subset LieG^{an}(\Lambda)$  of rank 3. Indeed, on using similar notations for the group schemes  $E/S$  and  $\mathbb{G}_m \times S$ , the canonical projection  $\pi : G \rightarrow E$  over  $S$  induces at the Lie algebra level an exact sequence

$$0 \longrightarrow Lie\mathbb{G}_m^{an} \longrightarrow LieG^{an} \xrightarrow{d\pi} LieE^{an} \longrightarrow 0 .$$

From the compatibility of the exponential morphisms, we deduce an exact sequence of  $\mathbb{Z}$ -local systems of periods

$$0 \longrightarrow \Pi_{\mathbb{G}_m} \longrightarrow \Pi_G \xrightarrow{d\pi} \Pi_E \longrightarrow 0 ,$$

with  $\Pi_{\mathbb{G}_m}(\Lambda)$  and  $\Pi_E(\Lambda)$  of respective ranks 1 and 2 over  $\mathbb{Z}$ .

There exists a unique analytic section  $U := \log_{G, \bar{\lambda}_0}$  of  $Lie(G^{an})/\Lambda$  such that

$$U(\bar{\lambda}_0) = U_0 \quad \text{and} \quad \forall \lambda \in \Lambda, exp_{G^{an}}(U(\lambda)) = s(\lambda).$$

Since  $\Lambda$  is fixed and  $\bar{\lambda}_0$  plays no role in what follows, we will forget about them and will just write  $U = \log_{G, \Lambda} = \log_G$ , i.e.

$$\forall \lambda \in \Lambda, U(\lambda) = \log_G(s(\lambda)).$$

We call  $U = \log_G(s)$  “the” logarithm of the section  $s$ . Its projection  $p = \pi(s) \in E(S)$  admits as logarithm

$$\log_E(p) := u = d\pi(U) = d\pi(\log_G(s)) = \log_E(\pi(s)).$$

We describe these logarithms in terms of classical Weierstrass functions in §4 for the (iso)trivial case  $G = \mathbb{G}_m \times E$ , and in Appendix I for the general case. These explicit expressions are not needed, but will provide the interested reader with a translation of the algebraic independence results in more classical terms.

Now, we rewrite  $U$  in terms of a conveniently chosen basis of the  $\mathbb{Z}$ -local system of periods  $\Pi_G/\Lambda$  of  $G^{an}/\Lambda$ . We call  $U^B(\lambda) = \log_G^B(s(\lambda))$  the resulting expression. For this, we choose a generator  $\varpi_0 = 2\pi i$  of  $\Pi_{\mathbb{G}_m}$ , and a  $\mathbb{Z}$ -basis

$\{\omega_1, \omega_2\}$  of  $\Pi_E(\Lambda)$ . At each point  $\bar{\lambda} \in \Lambda$ , the latter generate over  $\mathbb{R}$  the  $\mathbb{C}$ -vector space  $Lie(E_{\bar{\lambda}})$ . Consequently (and as already said before Proposition 3), there exist uniquely defined real analytic functions  $b_1, b_2 : \Lambda \rightarrow \mathbb{R}^2$  such that

$$\forall \lambda \in \Lambda, u(\lambda) = b_1(\lambda)\omega_1(\lambda) + b_2(\lambda)\omega_2(\lambda). \quad (\mathfrak{R}_u)$$

We call  $u^B = (b_1, b_2) : \Lambda \rightarrow \mathbb{R}^2$  the Betti presentation of the logarithm  $u = \log_E(p)$ .

Now, choose at will lifts  $\{\varpi_1, \varpi_2\}$  of  $\{\omega_1, \omega_2\}$  in  $\Pi_G(\Lambda)$ . Then,  $U - b_1\varpi_1 - b_2\varpi_2$  lies in the kernel  $Lie\mathbb{G}_m(\Lambda)$  of  $d\pi$ , which is generated over  $\mathbb{C}$  by  $\varpi_0$ . Therefore, there exist a unique real analytic function  $a : \Lambda \rightarrow \mathbb{C} = \mathbb{R}^2$  such that

$$U = a\varpi_0 + b_1\varpi_1 + b_2\varpi_2.$$

In conclusion, there exist uniquely defined real-analytic functions  $a : \Lambda \rightarrow \mathbb{C}, b_1 : \Lambda \rightarrow \mathbb{R}, b_2 : \Lambda \rightarrow \mathbb{R}$  such that  $U = \log_G(s)$  satisfies the relation :

$$\forall \lambda \in \Lambda, U(\lambda) = a(\lambda)\varpi_0(\lambda) + b_1(\lambda)\varpi_1(\lambda) + b_2(\lambda)\varpi_2(\lambda). \quad (\mathfrak{R}_U)$$

We call the *real analytic* map

$$U^B = (a, b_1, b_2) : \Lambda \rightarrow \mathbb{C} \times \mathbb{R}^2 = \mathbb{R}^4,$$

the Betti presentation of the logarithm  $U$  of  $\log_G(s)$ . Its image  $\mathcal{S} = \mathcal{S}_\Lambda := U^B(\Lambda) = \log_G^B(s(\Lambda)) \subset \mathbb{R}^4$  is the real surface to be studied. Since  $\Pi_G$  is the kernel of the exponential morphism, it is clear that for any  $\bar{\lambda} \in S_m^G$ ,  $U^B(\bar{\lambda})$  lies in  $\mathbb{Q}_m \times \mathbb{Q}_m^2 \subset \mathbb{Q}_m^4 \subset \mathbb{R}^4$ .

### Reducing to algebraic independence.

To complete their proofs, we must show that under the hypotheses of Theorems 1 and 2, the surface  $\mathcal{S}$  contains no semi-algebraic curve of the ambient affine space  $\mathbb{R}^4$ . This will be done in two steps, as follows. But before we describe them, we point out that since  $\log_E(p), \log_E(s), \omega_i, \varpi_i, \dots$  are local sections of the globally defined vector bundles  $(LieG)/S, (LieE)/S$ , it makes sense to speak of the minimal extension  $K(\log_G(s)), \dots$  of  $K = \mathbb{C}(\lambda)$  they generate in the field of meromorphic functions over a neighbourhood  $\Lambda'$  of  $\Lambda$ . A similar remark applies to the field  $K(a), \dots$  generated by the real analytic functions  $a, \dots$  in the fraction field of the ring of real analytic functions over  $\Lambda'$ .

• (*First step*) Let  $F_{pq}^{(1)} = K(\omega_1, \omega_2, \log_E(p), \log_E(q))$  be the field generated over  $K$  by  $\omega_1, \omega_2, \log_E(p) = u$  and a logarithm  $v = \log_E(q)$  of the section  $q \in \hat{E}(S) \simeq E(S)$  parametrizing the extension  $G$ . As usual, we assume that if  $E/S$  is isoconstant, then  $p$  is not constant. We then claim that *if  $\mathcal{S}$  contains a semi-algebraic curve, and if  $q = 0$ , then,  $\log_G(s)$  is algebraic over  $F_{pq}^{(1)}$* . In fact, we will only need a corollary of this result, involving the universal vectorial extensions of  $E$  and of  $G$ , where the base field  $F_{pq}^{(1)}$  is replaced by a **differential field**  $F_{pq} := F_{pq}^{(2)}$  containing  $F_{pq}^{(1)}$ , which, inspired by the theory of one-motives, we can call the *field of generalized periods of  $\{E, p, q\}$* . (In more classical terms, the upper indexes (1) and (2) here stand for elliptic integrals of the first and second kinds.) In these conditions, and under *no* assumption on  $q$ , we will prove :

**Proposition 4.** *Assume that  $p$  is not torsion and not constant, and that  $\mathcal{S}$  contains a semi-algebraic curve. Then,  $\log_G(s)$  is algebraic over the field  $F_{pq}$  of generalized periods of  $\{E, p, q\}$ .*

• (*Second step*) The desired contradiction is then provided by the following Main Lemma, whose proof is the object of §6 (and §4 for  $q = 0$ ). This is a statement of “Ax-Lindemann” type, but with logarithms replacing exponentials, in the style of André’s theorem [1] (see also [7]) for abelian schemes. For results on semi-abelian surfaces close to this Main Lemma, see [6], Propositions 4.a, 4.b and Theorem 2. For a broader perspective on algebraic independence of relative periods, see Ayoub’s recent paper [4].

**Main Lemma.** *With  $S/\mathbb{C}$ , let  $G/S$  be an extension by  $\mathbb{G}_m$  of an elliptic scheme  $E/S$ , parametrized by a section  $q$  of  $\hat{E}/S$ , and let  $G_0$  be the constant part of  $G$ . Let further  $s$  be a section of  $G/S$ , with projection  $p = \pi \circ s$  to  $E/S$ , and let  $F_{pq}$  be the field of generalized periods of  $\{E, p, q\}$ .*

**(A)** *Assume that  $\log_G(s)$  is algebraic over  $F_{pq}$ . Then, there exists a constant section  $s_0 \in G_0(\mathbb{C})$  such that*

- i) either  $s - s_0$  is a Ribet section;*
- ii) or  $s - s_0$  factors through a strict subgroup scheme of  $G/S$ .*

**(B)** *More precisely,  $\log_G(s)$  is algebraic over  $F_{pq}$  if and only if there exists a constant section  $s_0 \in G_0(\mathbb{C})$  such that  $s - s_0$  is a Ribet section, or a torsion section, or factors through a strict subgroup scheme of  $G/S$  projecting onto  $E/S$ .*

The analogy with the Main Theorem is clear, except perhaps for the last conclusion of Part (B) of the Main Lemma (which forces an isotrivial  $G \simeq \mathbb{G}_m \times E$ ). This is due to the fact that even in this isotrivial case, the roles of  $\mathbb{G}_m$  and  $E$  are here not symmetric, because of the occurrence of  $p$  in the base fields  $F_{pq}^{(1)}, F_{pq}$ . On the contrary (“torsion values for a single point” on a group scheme of relative dimension 1 over a curve), they played similar roles for relative Manin-Mumford.

**Proposition 4 + Main Lemma (A)  $\Rightarrow$  Theorems 1 and 2.**

Let us first deal with Theorem 1, where  $G = \mathbb{G}_m \times E$ , with constant part  $G_0 = \mathbb{G}_m$ , resp.  $G_0 \times S = G$ , if  $E$  is not, resp. is, isoconstant. Assume for a contradiction that  $S_\infty^G$  is infinite. Then, the real surface  $\mathcal{S}$  must contain an algebraic curve, and since  $G$  admits no Ribet section, Proposition 4, combined with Part (A) of the Main Lemma, implies that a multiple by a non-zero integer of the section  $s$  factors through a translate of  $H = \mathbb{G}_m$  or of  $H = E$  by a constant (non-necessarily torsion) section  $s_0 \in G_0(\mathbb{C})$ . But the projection  $p$  of  $s$  to  $E$  is by assumption not torsion, and we know that it cannot be constant. So,  $H$  must be equal to  $E$ , and  $s$  projects on the  $\mathbb{G}_m$ -factor of  $G$  to a constant point  $\delta_0$ . Since  $s(S) = W$  contains torsion points,  $\delta_0$  must be a root of unity, and  $s$  factors through a torsion translate of  $E$ , as was to be proved.

In the direction of Theorem 2, we now assume that  $G$  is a non-isotrivial extension, so  $H = \mathbb{G}_m/S$  is the only connected strict subgroup scheme of  $G/S$ . The proof above easily adapts to the case when  $G$  is isoconstant, where again,  $G$  admits no Ribet section (in the sense of §1). Now, assume that  $G$  is not isoconstant, so  $G_0 = \mathbb{G}_m$ , and that  $S_\infty^G$  is infinite. If  $G$  is not semi-constant, Proposition 4, combined with Part (A) of the Main Lemma, implies that a multiple of  $s$  factors through a translate of  $H = \mathbb{G}_m$  by a constant section  $s_0 \in \mathbb{G}_m(\mathbb{C})$ , so  $s$  factors through a torsion translate of  $\mathbb{G}_m$ , and  $p = \pi(s)$  is torsion, contradicting the hypothesis of Theorem 2. So,  $G$  must be semi-constant,  $E = E_0 \times S$  must be isoconstant, and the argument just described shows that  $s$  must satisfy Conclusion (i) of the Main Lemma, for some  $s_0 \in \mathbb{G}_m(\mathbb{C})$ . The mere existence of a Ribet section  $s_R := s - s_0$  of  $G/S$  implies that  $p = \pi(s) = \pi(s_R)$  and  $q$  are antisymmetrically related. Moreover, by Theorem 3.i,  $s_R(\bar{\lambda})$  is a torsion point on  $G_{\bar{\lambda}}$  whenever  $s(\bar{\lambda})$  is so, since  $\pi(s_R(\bar{\lambda})) = \pi(s(\bar{\lambda})) = p(\bar{\lambda})$  is then a torsion point of  $E_0$ . There are infinitely many such  $\bar{\lambda}$ 's, so at least one. Consequently the constant section  $s_0 \in \mathbb{G}_m(\mathbb{C})$  is torsion, and (a multiple of)  $s$  is a Ribet section of  $G/S$ . This

concludes the argument for Theorem 2.

**Part (B) of the Main Lemma.**

Just as for the Main Theorem (cf. Theorem 3), let us right now deal with the “if” side of Part B of the Main Lemma.

The periods  $\Pi_G$  of  $G$  are defined over the subfield  $F_q$  of  $F_{pq}$  (see §5.1, or the explicit formula given in Appendix I, §7.1, or the footnote below), so, clearly,  $\log_G(s)$  lies in  $F_{pq}$  if  $s - s_0$  is a torsion section. When  $s - s_0 := s_R$  is a Ribet section, an explicit formula for  $\log_G(s_R)$  in terms of  $u$  and  $\zeta_\lambda(u)$  is given in Appendix I, §7.2, from which the rationality of  $\log_G(s_R)$  over  $F_{pq}$  immediately follows. In fact, we will prove this in a style closer to Manin-Mumford issues in Lemma 3 of §6 below<sup>3</sup>. The last case forces  $G$  to be an isotrivial extension. In the notations of §4, we then have  $s - s_0 = (\delta, p) \in G(S)$ , with  $\delta$  a root of unity, so  $\log_G(s)$  is rational over the field  $F_p$ .

As for the “only if” side of Part (B) not covered by Part (A), we must show that if (a multiple by a positive integer of)  $s - s_0$  is a non constant section  $\delta$  of  $\mathbb{G}_m(S)$ , then  $\ell := \log_{\mathbb{G}_m}(\delta)$  is transcendental over  $F_{pq}$ . But then,  $p - \pi(s_0)$  is a torsion section of  $E/S$ , so  $F_{pq} = F_q$ , and the statement follows from Lemma 1 of §4, with  $q$  playing the role of  $p$ .

In conclusion, we have reduced the proof of the Main Theorem (more specifically, of Theorems 1 and 2) to defining the field  $F_{pq}$ , proving Proposition 4, and proving Part (A) of the Main Lemma.

## 4 A warm up : the case of direct products

In this Section, we perform the above-mentioned tasks under the assumption that  $G$  is an isotrivial extension, thereby establishing Theorem 1, as stated in §2. Without loss of generality, we assume that  $G = \mathbb{G}_m \times E$ , i.e.  $q = 0$  (so,

---

<sup>3</sup> More intrinsically, concerning the field of definition of  $\Pi_G$  : the Cartier dual of the one-motive  $[0 \rightarrow G]$  is the one-motive  $[\mathbb{Z} \rightarrow \hat{E}]$  attached to  $q \in \hat{E}(S)$ , so their fields of (generalized) periods coincide, and  $K(\Pi_G) = F_G^{(1)} \subset F_G^{(2)} = F_q$ . Concerning  $\log_G(s_R)$  : in the notations of §1.2, it suffices to consider the generic Ribet section  $s_R$  of the semi-abelian scheme  $\mathcal{P}_0$ , viewed as an extension  $\mathcal{G}_0$  of  $E_0$  by  $\mathbb{G}_m$ , over the base  $\hat{E}_0$ . As mentioned there, its image  $W_R$  is a special curve of the mixed Shimura variety  $\mathcal{P}_0$ . Therefore, the inverse image of  $W_R$  in the uniformizing space of  $\mathcal{P}_0$  is an algebraic curve. In the notations of §6, the statement amounts to the vanishing of  $\tau_{s_R}$ , and could alternatively be deduced from the self-duality of the one-motive  $[M_{s_R} : \mathbb{Z} \rightarrow G]$  attached to the Ribet section, cf. [10].



the field  $F_{pq} = F_{p0}$  will coincide with  $F_p$ ). Of course, if  $E/S$  is isoconstant, say  $E = E_0 \times S$ , then  $G = G_0 \times S$  with  $G_0 = \mathbb{G}_m \times E_0/\mathbb{Q}^{alg}$ , and Theorem 1 follows from Hindry's theorem [16]; in this isoconstant case, the strategy we are here following reduces to that of [25].

We first rewrite in concrete terms the logarithms  $U, u$  and their Betti presentations, under no assumption on the elliptic scheme  $E/S$  nor on its section  $p$ . We fix global differential forms<sup>4</sup> of the first and second kind  $\omega, \eta$  for  $E/S$ , and for any  $\lambda \in \Lambda$ , we let

$$\wp_\lambda, \zeta_\lambda, \sigma_\lambda$$

be the usual Weierstrass functions attached to the elliptic curve  $E_\lambda/\mathbb{C}$  and its differential forms  $\omega_\lambda, \eta_\lambda$ . We also fix an elliptic logarithm of the point  $p(\bar{\lambda}_0)$ , and extend it to an analytic function  $u(\lambda) = \log_E(p(\lambda)) = \text{Arg}_{\wp_\lambda}(p(\lambda))$  on  $\Lambda$ . Similarly, we fix a basis of periods and quasi-periods for  $E_{\bar{\lambda}_0}$ , and extend them to analytic functions  $\omega_1(\lambda), \omega_2(\lambda), \eta_1(\lambda), \eta_2(\lambda)$  (of hypergeometric type if  $E/S$  is the Legendre curve). There then exist uniquely defined real-analytic functions  $b_1, b_2$  with values in  $\mathbb{R}$  such that

$$\forall \lambda \in \Lambda, u(\lambda) = b_1(\lambda)\omega_1(\lambda) + b_2(\lambda)\omega_2(\lambda), \quad (\mathfrak{R}_u)$$

and the Betti presentation of  $\log_E(p(\lambda))$  is given by

$$u^B(\lambda) := \log_E^B(p(\lambda)) = (b_1(\lambda), b_2(\lambda)) \in \mathbb{R}^2.$$

We now go to  $G = \mathbb{G}_m \times E$  over  $\Lambda$ . The section  $s : \Lambda \rightarrow G$  has two components  $(\delta, p)$ , where  $\delta : S \rightarrow \mathbb{G}_m/S$  is expressed by a rational function on  $S$ . We fix a classical logarithm of  $\delta(\bar{\lambda}_0)$  and extend it to an analytic function  $\ell(\lambda) := \log_{\mathbb{G}_m}(\delta(\lambda))$  on  $\Lambda$ . With these notations, the section  $\log_G \circ s$  of  $(\text{Lie}G^{an})/\Lambda$  is represented by the analytic map

$$\Lambda \ni \lambda \mapsto \log_G(s(\lambda)) := U(\lambda) = \begin{pmatrix} \ell(\lambda) \\ u(\lambda) \end{pmatrix} \in \mathbb{C}^2 = (\text{Lie}G)_\lambda.$$

---

<sup>4</sup> When the modular invariant  $j(\lambda)$  is constant, i.e. when  $E/S$  is isoconstant, we tacitly assume that  $E = E_0 \times S$ , with  $E_0/\mathbb{C}$ , and that the chosen differential of first and second kind  $\omega, \eta$  are *constant* (i.e. come from  $E_0/\mathbb{C}$ ). In particular, the periods  $\omega_1, \omega_2$  and quasi-periods  $\eta_1, \eta_2$  are constant. The Weierstrass functions are those of  $E_0$ , and we drop the index  $\lambda$  from their notation.

The  $\mathbb{Z}$ -local system of periods  $\Pi_G$  admits the basis

$$\varpi_0(\lambda) = \begin{pmatrix} 2\pi i \\ 0 \end{pmatrix}, \varpi_1(\lambda) = \begin{pmatrix} 0 \\ \omega_1(\lambda) \end{pmatrix}, \varpi_2(\lambda) = \begin{pmatrix} 0 \\ \omega_2(\lambda) \end{pmatrix},$$

and the Betti presentation of  $\log_G(s(\lambda))$  is given by

$$\Lambda \ni \lambda \mapsto U^B(\lambda) := \log_G^B(s(\lambda)) = (a(\lambda), b_1(\lambda), b_2(\lambda)) \in \mathbb{C} \times \mathbb{R}^2 = \mathbb{R}^4,$$

where  $a, b_1, b_2$  are the unique real analytic functions on  $\Lambda$  satisfying

$$\forall \lambda \in \Lambda, \begin{pmatrix} \ell(\lambda) \\ u(\lambda) \end{pmatrix} = a(\lambda) \begin{pmatrix} 2\pi i \\ 0 \end{pmatrix} + b_1(\lambda) \begin{pmatrix} 0 \\ \omega_1(\lambda) \end{pmatrix} + b_2(\lambda) \begin{pmatrix} 0 \\ \omega_2(\lambda) \end{pmatrix}. \quad (\mathfrak{R}_{\ell, u})$$

We then set  $\mathcal{S} = U^B(\Lambda) \subset \mathbb{R}^4$  as usual.

Since  $q = 0$ , the tower of function field extensions of  $K = \mathbb{C}(\lambda)$  to be considered take here the following simple forms :

$$F^{(1)} = K(\omega_1, \omega_2) \quad , \quad F_{p_0}^{(1)} := F_p^{(1)} = F^{(1)}(u) = K(\omega_1, \omega_2, u),$$

while their differential extensions

$$F^{(2)} = F^{(1)}(\eta_1, \eta_2) \quad , \quad F_{p_0}^{(2)} := F_p^{(2)} = F_p^{(1)}(\zeta_\lambda(u)) = K(\omega_1, \omega_2, \eta_1, \eta_2, u, \zeta_\lambda(u))$$

involve the Weierstrass  $\zeta$  function, and can be rewritten as

$$F := F^{(2)} = K(\omega_1, \omega_2, \eta_1, \eta_2) \quad , \quad F_p := F_p^{(2)} = F(u, \zeta_\lambda(u)).$$

We point out that since it contains the field of definition  $F$  of the periods of  $\omega$  and  $\eta$ , the field  $F_p$  depends only on the section  $p$ , not on the choice of its logarithm  $u$ , so, the notation is justified. Furthermore, let  $\alpha \in \mathcal{O} = \text{End}(E)$  be a non-zero endomorphism of  $E$ , and let  $p_0 \in \mathcal{E}_0(\mathbb{C})$  be a constant section of  $E$ . Then, the section  $p' = \alpha p + p_0$  yields the same field  $F_{p'} = F_p$  as  $p$ . In particular,  $F_p = F$  if  $p$  is a torsion or a constant section of  $E/S$ .

#### **Proof of Proposition 4 when $q = 0$ .**

Suppose for a contradiction that  $\mathcal{S}$  contains a real semi-algebraic curve  $\mathcal{C}$ , and denote by  $\Gamma \subset \Lambda$  the inverse image of  $\mathcal{C}$  in  $\Lambda$  under the map  $U^B$  (all we will need is that  $\Gamma$  has an accumulation point inside  $\Lambda$ , but it is in fact a real curve). We are going to study the restrictions to  $\Gamma$  of the functions

$$a, b_1, b_2, u, \ell, \omega_1, \omega_2.$$

Recall that all these are functions of  $\lambda \in \Lambda$ . In view of the defining relation  $(\mathfrak{R}_{\ell,u})$ , the transcendence degree of the functions  $u, \ell$  over the field  $\mathbb{C}(\omega_1, \omega_2, a, b_1, b_2)$  is at most 0. When restricted to  $\Gamma$ , the latter field has transcendence degree  $\leq 1$  over  $\mathbb{C}(\omega_1, \omega_2)$ , since  $U^B(\Gamma) = (a, b_1, b_2)(\Gamma)$  is the algebraic curve  $\mathcal{C}$ . So, the restrictions to  $\Gamma$  of the two functions  $u, \ell$  generate over  $\mathbb{C}(\omega_{1|\Gamma}, \omega_{2|\Gamma})$  a field of transcendence degree  $\leq 1 + 0 = 1$ , and are therefore algebraically dependent over  $\mathbb{C}(\omega_{1|\Gamma}, \omega_{2|\Gamma})$ . Since  $\Gamma$  is a real curve of the complex domain  $\Lambda$ , the complex analytic functions  $u, \ell$  are still algebraically dependent over the field of  $\Lambda$ -meromorphic functions  $\mathbb{C}(\omega_1, \omega_2)$ , i.e.

$$\text{tr.deg.}_{\mathbb{C}(\omega_1(\lambda), \omega_2(\lambda))} \mathbb{C}(\omega_1(\lambda), \omega_2(\lambda), u(\lambda), \ell(\lambda)) \leq 1.$$

Now, assume as in Proposition 4 that  $p$  has infinite order, and if  $E/S$  is isotrivial, that  $p$  is not constant. Then, André's Theorem 3 in [1] (see also [5], Thm. 5) implies that  $u(\lambda)$  is transcendental over the field  $F^{(1)} = K(\omega_1(\lambda), \omega_2(\lambda))$ . The previous inequality therefore says that the function  $\ell(\lambda)$  is algebraic over the field  $F_p^{(1)} = K(\omega_1(\lambda), \omega_2(\lambda), u(\lambda))$ , or equivalently, that the field of definition of  $\log_G(s(\lambda)) = (\ell(\lambda), u(\lambda))$  is algebraic over  $F_{p0}^{(1)}$ , hence also over  $F_{p0} = F_{p0}^{(2)}$ , and Proposition 4 is proved when  $q = 0$ .

### Proof of Main Lemma (A) when $q = 0$

Before giving this proof, let us point out that the advantage of the fields  $F = F^{(2)}, F_{pq} = F_{pq}^{(2)}$  over their first kind analogues is that they are closed under the derivation  $' = \partial/\partial\lambda$ . Moreover, by Picard-Fuchs theory, they are Picard-Vessiot (i.e. differential Galois) extensions of  $K$ . Since  $\ell(\lambda)$  satisfies a  $K$ -rational DE of order 1,  $K(\ell)$  and  $F_p(\ell) = F_p(\log_G(s))$  too are Picard-Vessiot extension of  $K$ .

**Lemma 1.** *Let  $\Lambda$  be a ball in  $\mathbb{C}$ , let  $\{\wp_\lambda, \lambda \in \Lambda\}$  be a family of Weierstrass functions, with invariants  $g_2, g_3$  algebraic over  $\mathbb{C}(\lambda)$  and periods  $\omega_1, \omega_2$ , and let  $u$  be an analytic function on  $\Lambda$  such that  $u, \omega_1, \omega_2$  are linearly independent over  $\mathbb{Q}$ . If  $j(\lambda)$  is constant, we assume that  $g_2, g_3$  too are constant, and that  $u$  is not constant. Let further  $\ell$  be a non-constant analytic function on  $\Lambda$ . Assume that  $\wp_\lambda(u(\lambda))$  and  $e^{\ell(\lambda)} := \delta(\lambda)$  are algebraic functions of  $\lambda$ , and consider the tower of differential fields  $K \subset F \subset F_p$ , where  $K = \mathbb{C}(\lambda), F = K(\omega_1, \omega_2, \eta_1, \eta_2), F_p = F(u, \zeta_\lambda(u))$ . Then,*

$$\text{tr.deg.}_{F} F_p(\ell(\lambda)) = 3.$$

In particular,  $\ell(\lambda)$  is transcendental over  $F_p$ , i.e. Part (A) of the Main Lemma holds true when  $q = 0$ .

This last statement is indeed equivalent to Part (A) of the Main Lemma when  $G \simeq \mathbb{G}_m \times E$  is a trivial extension (or more generally, an isotrivial one), with constant part  $G_0 = \mathbb{G}_m$ , resp.  $G_0 \times S = G$ , if  $E$  is not, resp. is, isoconstant. Indeed, with  $s = \exp_G(\ell, u) = (\delta, p)$  as above, we then have  $F_p = F_{pq}$ , and  $F_p(\ell) = F_p(\log_G(s))$ . Lemma 1 then says that if  $\log_G(s)$  is algebraic over  $F_{pq}$ , then either  $p$  is a torsion point (so, a multiple of  $s$  factors through  $\mathbb{G}_m$ ), or  $E$  isoconstant and  $p = p_0$  is constant (so, the constant section  $s_0 = (1, p_0) \in G_0(\mathbb{C})$  satisfies  $s - s_0 \in \mathbb{G}_m(S)$ ), or  $\delta = \delta_0$  is constant (so, the constant section  $s_0 = (\delta_0, 0) \in G_0(\mathbb{C})$  satisfies  $s - s_0 \in E(S)$ ). In all cases, we therefore derive Conclusion (ii) of the Main Lemma.

*Proof* (of Lemma 1): this is essentially due to André, cf. [1], but not fully stated there (nor in [5]). It is proven in full generality in [7], but one must look at the formula on top of p. 2786 to see that  $K$  can be replaced by  $F$  in Theorem L ... So, it is worth giving a direct proof.

We first treat the case when  $E/S$  is not isoconstant. By Picard-Lefschetz, the Picard-Vessiot extension  $F = K(\omega_1, \omega_2, \eta_1, \eta_2)$  of  $K = \mathbb{C}(\lambda)$  has Galois group  $SL_2$ . By [5], the Galois group of  $F_p = F(u(\lambda), \zeta_\lambda(u(\lambda)))$  over  $F$  is a vector group  $\mathcal{V} \simeq \mathbb{C}^2$  of dimension 2 (i.e. these two functions are algebraically independent over  $F$ ), while the Galois group of  $K(\ell(\lambda))$  over  $K = \mathbb{C}(\lambda)$  is  $\mathbb{C}$ . Since  $\mathbb{C}$  is not a quotient of  $SL_2$ , the Galois group of  $F(\ell(\lambda))$  over  $F$  is again  $\mathbb{C}$ . Now,  $SL_2$  acts on the former  $\mathcal{V} = \mathbb{C}^2$  by its standard representation, and on the latter  $\mathbb{C}$  via the trivial representation, so the Galois group of  $F(u(\lambda), \zeta_\lambda(u(\lambda)), \ell(\lambda))$  over  $F$  is a subrepresentation  $\mathcal{W}$  of  $SL_2$  in  $\mathbb{C}^2 \oplus \mathbb{C}$  projecting onto both factors. Since the standard and the trivial representations are irreducible and non isomorphic, we must have  $\mathcal{W} = \mathbb{C}^2 \oplus \mathbb{C}$ . Therefore,

$$tr.deg.F F(u(\lambda), \zeta_\lambda(u(\lambda)), \ell(\lambda)) = dim\mathcal{W} = 3.$$

We now turn to the case of an isoconstant  $E = E_0 \times S$ . The field of periods  $F$  then reduces to  $K$ , and  $SL_2$  disappears. But since the ambient group  $G = \mathbb{G}_m \times E$  is now isoconstant, we can appeal to Ax's theorem on the functional version of the Schanuel conjecture. More precisely, since the result we stated involves the  $\zeta$ -function, we appeal to its complement on vectorial extensions, see [12], Thm. 2.(iii), or more generally, [6], Proposition 1.b, which implies that  $tr.deg_K K(u, \zeta(u), \ell) = 3$  as soon as  $u$  and  $\ell$  are not

constant. (Actually, André's method still applies to the isoconstant case, but requires a deeper argument, involving Mumford-Tate groups : cf. [1], Theorem 1, and [7], §8.2).

**Remark 2 :** (i) Concerning the proof of Theorem 1 : in fact, as pointed out by the fourth author in [31], p. 79, Comment (v), since we are here dealing with a direct product, the torsion points yield torsion points on  $\mathbb{G}_m$ , which lie on the unit circle, a real-curve. So, in the argument of §3.2, the dimension decreases by 1 a priori, and Bombieri-Pila (real-curves) rather than Pila (real-surfaces) as in Proposition 2 would suffice.

(ii) As shown by the proof of Proposition 4 ( $q = 0$ ) above, it would have sufficed to prove that the (non constant) logarithm  $\ell$  is transcendental over the field  $\mathbb{C}(\omega_1, \omega_2, u)$ . Adjoining  $\lambda$  leads to  $F_p^{(1)}$  as base field, and as already said, differential algebra then forces to consider  $F_p^{(2)}$ . For a broader perspective on these extensions of the base field, see §5.3 below.

(iii) For the last statement of Lemma 1 to hold, the only necessary hypothesis is that  $\ell$  be non-constant. Indeed, if  $p$  is torsion or constant, then  $F_p = F$ , and  $u$  plays no role. But we have preferred to present Lemma 1 and its proof in this way, as an introduction to the general proofs of §5 and §6.

(iv) Conversely, let  $q$  be any (non-necessarily torsion or constant) section of  $E/S$ , and set  $v = \log_E(q)$ ,  $F_q = F^{(2)}(v, \zeta_\lambda(v))$  and  $F_{pq} = F_p \cdot F_q$ , as will be done in §5. The same proof as above shows that

$\forall p, q \in E(S), \forall \delta \in \mathbb{G}_m(S), \delta \notin \mathbb{G}_m(\mathbb{C}), \ell := \log_{\mathbb{G}_m}(\delta)$  is transcendental over  $F_{pq}$ .

Indeed, the only new case is when  $p$  and  $q$  are linealy independent over  $\text{End}(E)$  modulo the constant part of  $E/S$ . From the same references and argument as above, replacing  $\mathcal{V} = \mathbb{C}^2$  by  $\mathcal{V} \oplus \mathcal{V}$ , we deduce that the transcendence degree of  $F_{pq}(\ell)$  over  $F$  is equal to 5, yielding the desired conclusion on the transcendency of  $\ell$ .

## A characterization of Ribet sections

We close this section on isotrivial extensions by a corollary to Theorem 1, which plays a useful role in checking the compatibility of the various definitions of Ribet sections : see the equalities  $\beta_R = \beta_J$  in [8], and  $s_{\tilde{R}} = s_R$  in Appendix I, Proposition 5.(i) below.

**Corollary 3.** *Let  $G/S$  be an extension of  $E/S$  by  $\mathbb{G}_m$ , and let  $p$  be a section of  $E/S$  of infinite order and not constant if  $E/S$  is isoconstant (equivalently, by Proposition 1.iv, such that the set  $S_\infty^E$  attached to  $p$  is infinite). Let further  $s^\dagger$  and  $s$  be two sections of  $G/S$  such that  $\pi \circ s^\dagger = \pi \circ s = p$ . Assume that for all but finitely many (resp. for infinitely many) values of  $\bar{\lambda}$  in  $S_\infty^E$ , the point  $s^\dagger(\bar{\lambda})$  (resp.  $s(\bar{\lambda})$ ) lies in  $G_{tor}$ , i.e. that  $s^\dagger$  (resp.  $s$ ) “lifts almost all (resp. infinitely many) torsion values of  $p$  to torsion points of  $G$ ”. Then, there exists a torsion section  $\delta_0 \in \mathbb{G}_m(\mathbb{C})$  (i.e. a root of unity) such that  $s = s^\dagger + \delta_0$ .*

*Proof :* Let  $\delta_0 := s - s^\dagger \in \mathbb{G}_m(S)$ . We know that  $s^\dagger$  lifts almost all torsion points  $p(\bar{\lambda}) \in E_{tor}$  to points in  $G_{tor}$ . If  $s$  does so for infinitely many of them, then, so does the section  $s_1 := (\delta_0, p)$  of the direct product  $G_1 = \mathbb{G}_m \times E/S$ . This contradicts Theorem 1, unless the projection  $\delta_0(S)$  of  $s_1(S)$  to  $\mathbb{G}_m$  is reduced to a root of unity.

It is interesting to note that in this way, Theorem 1 on the trivial extension  $G_1$  has an impact on extensions  $G$  which need not at all be isotrivial. For instance, if  $G$  is semi-constant and  $E_0$  has CM, Corollary 3, applied to the Ribet section  $s^\dagger = s_R$ , shows that up to isogenies,  $s_R$  is the only section which lifts infinitely many torsion values of  $\pi(s_R)$  to torsion points of  $G$ . We also point out that since the elliptic scheme  $E \simeq E_0 \times S$  is here constant, Hindry’s theorem on the constant semi-abelian variety  $\mathbb{G}_m \times E_0$  suffices to derive this conclusion.

On the other hand, assume that  $G$  is an isotrivial extension. Then, there exists a subgroup scheme  $E^\dagger/S$  of  $G/S$  such that the restriction  $\pi^\dagger$  of  $\pi : G \rightarrow E$  to  $E^\dagger$  is an  $S$ -isogeny. Any section  $s^\dagger$  of  $G/S$  a non zero multiple of which factors through  $E^\dagger$  then satisfies the lifting property of the corollary, since  $p(\bar{\lambda}) := \pi^\dagger \circ s^\dagger(\bar{\lambda})$  is a torsion point of  $E_{\bar{\lambda}}$  if and only if  $s^\dagger(\bar{\lambda})$  is a torsion point of  $G_{\bar{\lambda}}$ . By Corollary 3, such sections  $s^\dagger$  are, up to a root of unity, the only section  $s$  above  $p = \pi \circ s^\dagger$  for which  $s(S) \cap G_{tor}$  is infinite. Of course, this is (after an isogeny) just a rephrasing of Theorem 1, but it shows the analogy between these “obvious” sections and the Ribet sections. This is a reflection of the list of special curves of the mixed Shimura variety described in §1.2.

## 5 The general case

### 5.1 Fields of periods and the Main Lemma

Apart from the statement of Lemma 2 below, we henceforth make no assumption on the extension  $G$  of  $E/S$  by  $\mathbb{G}_m$ . So the section  $q \in \hat{E}(S)$  which parametrizes  $G$  is arbitrary. Concerning the elliptic scheme  $E/S$ , we recall the notations of §4, and in particular, the fields of periods  $F^{(1)} = K(\omega_1, \omega_2)$  of  $E$  and its differential extension  $F := F^{(2)} = F^{(1)}(\eta_1, \eta_2)$ . We identify  $\tilde{E}$  and  $E$  in the usual fashion, and denote by  $v = \log_E(q)$  a logarithm of the section  $q$  over  $\Lambda$ . We recall that the field  $F_q := F_q^{(2)} = F^{(2)}(v, \zeta_\lambda(v))$  depends only on  $q$ , and coincides with  $F^{(2)} = F$  when  $q$  is a torsion section, i.e. when  $G$  is isotrivial. We will also use the notations of §3.3 on the local system of periods  $\Pi_G = \mathbb{Z}\varpi_0 \oplus \mathbb{Z}\varpi_1 \oplus \mathbb{Z}\varpi_2 \subset \text{Lie}G^{an}(\Lambda)$  of  $G^{an}/\Lambda$ .

Consider the extension  $F_G^{(1)} = K(\varpi_0, \varpi_1, \varpi_2) = K(\varpi_1, \varpi_2)$  of  $K = \mathbb{C}(\lambda)$  generated by the elements of  $\Pi_G$ . Since  $\Pi_G$  projects onto  $\Pi_E$  under  $d\pi$ , whose kernel has relative dimension 1, this field is an extension of  $F^{(1)} = K(\omega_1, \omega_2)$  of transcendence degree  $\leq 2$ . So, the field  $F_G^{(2)}$  generated by  $\Pi_G$  over  $F := F^{(2)}$  has transcendence degree  $\leq 2$ . In fact, the duality argument mentioned in Footnote (3), or more explicitly, the computation given in Appendix I, shows that

$$F_G^{(2)} = F^{(2)}(v, \zeta_\lambda(v)) := F_q,$$

i.e.  $F_G^{(2)}$  coincide with the differential extension  $F_q$  of  $F^{(2)} = F$  attached to  $q$ . So,  $F_G^{(2)}$  is in fact a Picard-Vessiot extension of  $K$ .

A more intrinsic way to describe these “fields of the second kind” is to introduce the **universal vectorial extension**  $\tilde{E}/S$  of  $E/S$ , cf. [11]. This is an  $S$ -extension of  $E/S$  by the additive group  $\mathbb{G}_a$ , whose local system of periods  $\Pi_{\tilde{E}}$  generates the field  $F^{(2)}$ . The universal vectorial extension  $\tilde{G}/S$  of  $G/S$  is the fiber product  $G \times_E \tilde{E}$ , and its local system of periods  $\Pi_{\tilde{G}}$  generates the field  $F_G^{(2)}$ . Now, for both  $\tilde{E}$  and  $\tilde{G}$  (and contrary to  $E$  and  $G$ ), these local systems generate the spaces of horizontal vectors of connections  $\partial_{\text{Lie}\tilde{E}}, \partial_{\text{Lie}\tilde{G}}$  on  $\text{Lie}\tilde{E}/S, \text{Lie}\tilde{G}/S$ . This explains why the fields  $K(\Pi_{\tilde{E}}) = F^{(2)} = F$  and  $K(\Pi_{\tilde{G}}) = F_G^{(2)} = F_q$  are Picard-Vessiot extensions of  $K$ .

Let now  $s$  be a section of  $G/S$ , and let  $U = \log_G(s) \in \text{Lie}G^{an}(\Lambda)$  be a logarithm of  $s$  over  $\Lambda$ . As usual, set  $p = \pi(s) \in E(S), u = \log_E(p) = d\pi(U) \in \text{Lie}E^{an}(\Lambda)$ . Since  $\text{Ker}(d\pi)$  has relative dimension 1, the field generated over

$K$  by  $\log_G(s)$  is an extension of  $K(\log_E(p))$  of transcendence degree  $\leq 1$ , so  $\log_G(s)$  has transcendence degree  $\leq 1$  over  $F_p = F(u, \zeta_\lambda(u))$ . Finally, set

$$F_{pq} := F_p \cdot F_q, \quad L = L_s := F_{pq}(\log_G(s)).$$

The field  $F_{pq} = F_{pq}^{(2)}$  is the field of generalized periods of  $\{E, p, q\}$  promised in §3.3. Since it contains  $F_G^{(2)}$ , the field  $L = L_s$  depends only on  $s$ , not on the choice of its logarithm  $U = \log_G(s)$ , and is an extension of  $F_{pq}$ , of transcendence degree  $\leq 1$ . In fact, the explicit formulae of Appendix I show that for  $q \neq 0$  and  $p \neq 0, -q$ :

$$L_s = F_{pq}(\ell_s - g_\lambda(u, v)), \quad \text{where } g_\lambda(u, v) = \log_{G_m} \frac{\sigma_\lambda(v+u)}{\sigma_\lambda(v)\sigma_\lambda(u)}$$

is a Green function attached to the sections  $\{p, q\}$  of  $E/S$ , and  $\ell_s = \log_{G_m}(\delta_s)$  for some rational function  $\delta_s \in K^*$  attached to the section  $s$  of  $G/S$ . This formula implies that  $L$  is a differential field, but  $L$  is even a Picard-Vessiot extension of  $K$ . One way to check this is to relate  $\log_G(s)$  to an integral of a differential of the 3rd kind on  $E$ , with integer, hence *constant* residues, and to differentiate under the integral sign. Another way consists in lifting  $s$  to a section  $\tilde{s}$  of  $\tilde{G}/S$ , projecting to  $\tilde{p} \in \tilde{E}(S)$ . Then, for any choices  $\tilde{u} = \log_{\tilde{E}}(\tilde{p})$  and  $\tilde{U} = \log_{\tilde{G}}(\tilde{s})$  of logarithms of  $\tilde{p}$  and  $\tilde{s}$ , the field  $F^{(2)}(\tilde{u}) = F(u, \zeta_\lambda(u)) = F_p$  is contained in  $F_q(\tilde{U})$ , which can therefore be written as  $F_{pq}(\tilde{U})$ , and the latter field  $F_{pq}(\tilde{U})$  coincides with  $F_{pq}(U) = L$ , since  $\tilde{U}$  lifts  $U$  and  $\tilde{u}$  in the fiber product  $Lie\tilde{G} = LieG \times_{LieE} Lie\tilde{E}$ . Now, in the notations of [11], [7],  $\tilde{U}$  is a solution of the inhomogenous linear system  $\partial_{Lie\tilde{G}}(\tilde{U}) = \partial \ell n_{\tilde{G}}(\tilde{s})$ , which, on the one hand, is defined over  $K$ , and on the other hand, admits  $F_q(\tilde{U})$  as field of solutions. So,  $L = F_q(\tilde{U})$  is indeed a Picard-Vessiot extension of  $K$ . By the same argument, applied to the differential equation  $\partial_{Lie\tilde{E}}(\tilde{u}) = \partial \ell n_{\tilde{E}}(\tilde{p})$ , we see anew that  $F_p$  is a Picard-Vessiot extension of  $K$ . Notice, on the other hand, that  $F_q(U)$  is in general not a differential extension of  $F_q$  (it contains  $u$ , but not  $\zeta_\lambda(u)$ ).

The following diagram summarizes these notations (and proposes other



natural ones...):

$$\begin{array}{ccc}
& L & \\
& \uparrow & \\
& F_{pq} & \\
F_q \nearrow & & \nwarrow F_p \\
& F & \\
& \uparrow & \\
& K & 
\end{array}
\quad
\begin{array}{l}
L = F_{pq}(\log_G(s)) = F_{\tilde{G}}(\log_{\tilde{G}}(\tilde{s})) = F_{pq}(\ell_s - g(u, v)) \\
F_{pq} = F_p \cdot F_q = F(u, \zeta(u), v, \zeta(v)) \\
F_q = F_G^{(2)} := F_{\tilde{G}} = F_{\tilde{E}}(\log_{\tilde{E}}(\tilde{q})) \quad F_p = F_{\tilde{E}}(\log_{\tilde{E}}(\tilde{p})) \\
F = F^{(2)} := F_{\tilde{E}} = K(\omega_1, \omega_2, \eta_1, \eta_2) \\
K = \mathbb{C}(\lambda)
\end{array}$$

All the notations of the Main Lemma have now been discussed, and we can restate its Part (A) in the non-isotrivial case, as follows :

**Lemma 2.** (= Main Lemma for  $q$  non-torsion) *With  $S/\mathbb{C}$ , let  $G/S$  be a non isotrivial extension by  $\mathbb{G}_m$  of an elliptic scheme  $E/S$ , parametrized by a section  $q$  of  $\hat{E}/S$ , and let  $G_0$  be the constant part of  $G$ . Let further  $s$  be a section of  $G/S$ , with projection  $p = \pi \circ s$  to  $E/S$ , and let  $F_{pq} = F_p \cdot F_q \supset F$  be the field of generalized periods of  $\{E, p, q\}$ . Assume that  $\log_G(s)$  is algebraic over  $F_{pq}$ . Then, there exists a constant section  $s_0 \in G_0(\mathbb{C})$  such that*

- i) either  $s - s_0$  is a Ribet section;*
- ii) or  $s - s_0$  is a torsion section.*

*In other words, if  $s$  is not a constant translate of a Ribet or a torsion section of  $G/S$ , then  $g_\lambda(u, v) - \ell_s$  is transcendental over  $F_{pq}$ .*

Conclusion (ii) of Lemma 2 appears to be stronger than Conclusion (ii) of Part (A) of the Main Lemma, but is in fact equivalent to it when  $G$  is not isotrivial. Indeed, in this case,  $\mathbb{G}_{m/S}$  is the only connected strict subgroup scheme of  $G$ . Now, if a multiple by a non-zero integer  $N$  of the section  $s - s_0$  factors through  $\mathbb{G}_m$ , i.e. is of the form  $\delta$  for some section  $\delta \in \mathbb{G}_m(S)$ , then,  $p = \pi(s)$  is a torsion or a constant section of  $E/S$ , so  $F_{pq} = F_q$ , and  $F_{pq}(\log_G(s)) = F_q(\ell)$ , where  $\ell = \log_{\mathbb{G}_m}(\delta)$ . By assumption,  $\ell$  is then algebraic over  $F_q$ , and Lemma 1 of §4 implies that  $\delta = \delta_0 \in \mathbb{G}_m(\mathbb{C})$  is a constant section. Considering the constant section  $s'_0 = s_0 - \frac{1}{N}\delta_0$  of  $G/S$ , we derive that  $s - s'_0$  is a torsion section, i.e. that Conclusion (ii) of Lemma 2 is fulfilled.

## 5.2 Reducing the Main Theorem to the Main Lemma

In view of §4, we could now restrict to the case of a non isotrivial extension  $G$ , prove Proposition 4 in this case, and finally prove Lemma 2, thereby concluding the proof of Theorem 2. However, as announced above, we will remain in the general case, and make no assumption on  $q$ .

We now prove Proposition 4, extending the pattern of proof of §4 to the general case. We recall the notations of Proposition 4, including the fundamental assumption that  $p = \pi(s)$  is neither a torsion nor a constant section of  $E/S$ . By Lemma 1, this condition implies that  $F_p$  has transcendence degree 2 over  $F$ , hence that  $u(\lambda) = \log_E(p(\lambda))$  is transcendental over the field  $F$ .

Consider the following tower of fields of functions on  $\Lambda$ , where the lower left (resp. upper right) ones are generated by complex (resp. real) analytic functions. The inclusions which the NE-arrows represent come from the definition of  $a, b_1, b_2$  in terms of  $\log_E(p(\lambda)) = u(\lambda), \log_G(s(\lambda)) = U(\lambda)$ , cf. Relations  $(\mathfrak{R}_u), (\mathfrak{R}_U)$  of §3.3; the inclusions of the NW-arrows on the left come from the definition of the fields of periods  $F, F_{pq}$ ; those of the NW-arrows on the right are obvious.

$$\begin{array}{ccccc}
 & & F_{pq}(a, b_1, b_2) & & (\mathfrak{R}_U) : \quad \log_G s = a\varpi_0 + b_1\varpi_1 + b_2\varpi_2 \\
 & \nearrow & & \nwarrow & \\
 F_{pq}(\log_G(s)) & & & & F_{pq}(b_1, b_2) \\
 & \nwarrow & & \nearrow & \\
 & & F_{pq} & & F(b_1, b_2) \\
 & & & \nwarrow & \nearrow \\
 & & & F(\log_E(p)) & (\mathfrak{R}_u) : \quad u := \log_E p = b_1\omega_1 + b_2\omega_2 \\
 & & & \uparrow & \\
 & & & F & 
 \end{array}$$

Now, assume for a contradiction that the real surface  $\mathcal{S}$  contains a semi-algebraic curve  $\mathcal{C}$ . As in §4, consider the real curve  $\Gamma = (U^B)^{-1}(\mathcal{C}) \subset \Lambda \subset S(\mathbb{C})$ , and denote by a lower index  $\Gamma$  the restrictions to  $\Gamma$  of the various functions of  $\lambda$  appearing above; similar notation  $F|_\Gamma, F_{pq}|_\Gamma$ , etc, for the fields they generate. For instance, since  $(a|_\Gamma, b_1|_\Gamma, b_2|_\Gamma)$  parametrize the algebraic

curve  $\mathcal{C}$ , these three functions generate a field of transcendence degree 1 over  $\mathbb{R}$ , and  $tr.deg_{F_\Gamma} F_\Gamma(b_{1|\Gamma}, b_{2|\Gamma}) \leq 1$ . But by the result recalled above and the principle of isolated zeroes,  $u_\Gamma$  is transcendental over  $F_\Gamma$ . Therefore, the restriction to  $\Gamma$  of the field  $F(b_1, b_2)$  is an algebraic extension of the restriction to  $\Gamma$  of the field  $F(u)$ . We may abbreviate this property by saying that  $F(u)$  and  $F(b_1, b_2)$  are essentially equal over  $\Gamma$ . Going up north-west in the tower, we deduce that the fields  $F_{pq}$  and  $F_{pq}(b_1, b_2)$  are essentially equal over  $\Gamma$ .

Notice that  $b_{1|\Gamma}$  and  $b_{2|\Gamma}$  are not both constant since  $F_\Gamma(b_{1|\Gamma}, b_{2|\Gamma}) := (F(b_1, b_2))_\Gamma \supset (F(u))_\Gamma$  is a transcendental extension of  $F_\Gamma$ . So,  $a_\Gamma$  must be algebraic over  $\mathbb{R}(b_{1|\Gamma}, b_{2|\Gamma})$ . Therefore,  $(F_{pq}(a, b_1, b_2))_\Gamma$  is an algebraic extension of  $(F_{pq}(b_1, b_2))_\Gamma$ , hence of the essentially equal field  $(F_{pq})_\Gamma$ , and we deduce that the intermediate field  $(F_{pq}(\log_G(s)))_\Gamma$  is algebraic over  $(F_{pq})_\Gamma$ . But  $\log_G(s(\lambda)) = U(\lambda)$  is a complex analytic map, so, by isolated zeroes,  $F_{pq}(\log_G(s(\lambda)))$  must also be algebraic over  $F_{pq}$ . This concludes the proof of Proposition 4.

### 5.3 The role of $K$ -largeness

The change of base field from  $K$  to  $F_{pq}$  can be viewed as the “logarithmic” equivalent of the passage from  $K$  to the field  $K_{\mathbf{G}}^\sharp$  generated by the Manin kernel  $\mathbf{G}^\sharp := Ker(\partial \ell n_{\mathbf{G}})$  of an algebraic  $D$ -group  $\mathbf{G}/K$ , which one encounters in the study of the exponentials of algebraic sections of  $Lie(\mathbf{G})/S$ , as in [11]. A Manin kernel has in fact already appeared in the present paper at the level of the  $D$ -group  $\tilde{E}$ : in this case, the group of  $K^{alg}$ -points of  $\tilde{E}^\sharp$  projects onto  $E_{tor} \oplus \mathcal{E}_0(\mathbb{C})$ , and the recurrent hypothesis made on the section  $p$  (that it be neither torsion nor constant) exactly means that none of its lifts  $\tilde{p}$  to  $\tilde{E}(K)$  lies in  $\tilde{E}^\sharp$ .

For elliptic curves, the field of definition  $K_{\tilde{E}}^\sharp$  of  $\tilde{E}^\sharp$  is always algebraic over  $K$ , and one says that the  $D$ -group  $\tilde{E}$  is  $K$ -large. This is the hypothesis required on  $\mathbf{G}$  for the Galois theoretic approach to the proof of the relative Lindemann-Weierstrass theorem of [11], §6 (see also [7], §8.1 and §4.3 of S. & C. for the abelian case). But the third author [26] has checked that it can be extended to non  $K$ -large groups (those with  $K_{\mathbf{G}}^\sharp$  transcendental over  $K$ ), as follows :

**Theorem** : *Let  $\mathbf{G}$  be an almost semi-abelian algebraic  $D$ -group over  $K$ , let  $a \in Lie\mathbf{G}(K)$ , and let  $y = exp_{\mathbf{G}}(\int a) \in \mathbf{G}(K^{diff})$  be a solution of the*

equation  $\partial \ln_{\mathbf{G}}(y) = a$ . Then,  $\text{tr.deg.}(K_{\mathbf{G}}^{\sharp}(y)/K_{\mathbf{G}}^{\sharp})$  is the smallest among dimensions of connected algebraic  $D$ -groups  $H$  of  $\mathbf{G}$  defined over  $K$  such that  $a \in \text{Lie}H + \partial \ln_{\mathbf{G}}(\mathbf{G}(K))$ , equivalently such that  $y \in H + \mathbf{G}(K) + \mathbf{G}^{\sharp}$ . Moreover,  $H^{\sharp}(K^{\text{diff}})$  is the Galois group of  $K_{\mathbf{G}}^{\sharp}(y)$  over  $K_{\mathbf{G}}^{\sharp}$ .

By  $\int a$ , we here mean any  $x \in \text{Lie}\mathbf{G}(K^{\text{diff}})$  such that  $\partial_{\text{Lie}\mathbf{G}}x = a$ . When  $x$  lies in  $\text{Lie}\mathbf{G}(K)$ , this leads to results of ‘‘Ax-Lindemann’’ type (as used in [25], [23]), whereas the transcendence results required by the present strategy (= that of [19]) concern the equation  $\partial_{\text{Lie}\mathbf{G}}x = b$ , with  $b = \partial \ln_{\mathbf{G}}(y)$  for some  $y \in \mathbf{G}(K)$ . Notice that when  $\mathbf{G}$  is the universal vectorial extension  $\tilde{G}$  of our semi-abelian scheme  $G = G_q$ , the analogous field  $K_{\text{Lie}\mathbf{G}}^{\sharp}$  is precisely the field of periods  $F_q = F_{\tilde{G}}$  of  $\tilde{G}$ ; see [7], §2 of S & C, for a justification of this analogy, and Remark 3.ii below for the adjunction of  $F_p$  in the base field.

More generally, given  $(x, y) \in (\text{Lie}\mathbf{G} \times \mathbf{G})$ , analytic over a ball  $\Lambda \subset S(\mathbb{C})$  and linked by the relation  $\partial \ln_{\mathbf{G}}(y) = \partial_{\text{Lie}\mathbf{G}}x$  (i.e. essentially  $y = \exp_{\mathbf{G}}(x)$ ), one may wonder which extension of the Ax-Schanuel theorem holds for the transcendence degree over  $K_{\text{Lie}\mathbf{G}}^{\sharp} \cdot K_{\mathbf{G}}^{\sharp}$  of the point  $(x, y)$ . The case when  $x$  is algebraic over  $K_{\text{Lie}\mathbf{G}}^{\sharp}$  includes the study of Picard-Painlevé sections; in this direction, see [14], Lemma 3.4. The case when  $x$  has transcendence degree  $\leq 1$  over  $K_{\text{Lie}\mathbf{G}}^{\sharp}$  would be of particular interest, as it occurs when the Betti coordinates of  $x$  parametrize an algebraic curve, and this may pull the present strategy back into the ‘‘exponential’’ framework of Ax-Lindemann-Weierstrass.

**Remark 3.**- i) Just as in §4, all we need to know for relative Manin-Mumford statements is the transcendence degree of  $\log_G \circ s$  over the field of generalized periods  $F_{pq}$  of  $\{E, p, q\}$ . More clearly put, the transcendence degree (i.e. the differential Galois group) of  $F_{pq}$  over  $K$  plays no role. Of course,  $\text{Gal}(F_{pq}/K)$  will come as a help during the proof of the Main Lemma, in parallel with the role of  $SL_2$  in §4. But in the notations of the diagram of §6, we must merely compute  $\text{Gal}(L/F_{p,q}) = \text{Im}(\tau_s) \subset \mathbb{C}$ , and show that under the hypotheses of the Main Lemma,  $\tau_s$  vanishes only if one of its conclusions (i), (ii) is satisfied.

ii) Adjoining the field  $F_p$  to the base field  $F_q$  comes in naturally, since the Picard-Vessiot extension  $F_q(\tilde{U})$  of  $K$  automatically contains it. But another advantage of the compositum  $F_{pq}$  is that the roles of  $p$  and  $q$  become symmetric in the statement of the Main Lemma. The explicit formula for  $L$  given by the Green function makes this apparent. More intrinsically, the field  $L_s = F_{pq}(\log_G(s))$  is the field of periods of the smooth one-motive

$M = [M_s : \mathbb{Z} \rightarrow G]$  over  $S$ , in the sense of [13], attached to the section  $s \in G(S)$ , with  $p = \pi(s)$  and  $q \in \hat{E}(S)$  parametrizing the extension  $G$ . By biduality,  $p$  parametrizes an extension  $G'$  of  $\hat{E}$  by  $\mathbb{G}_m$ , and  $s$  may be viewed as a section  $s'$  of  $G'$  above the section  $q$  of  $\hat{E}(S)$ . The Cartier dual of  $M$  is the one-motive  $M' = [M'_{s'} : \mathbb{Z} \rightarrow G']$  attached to this section  $s'$ , and its field of periods  $F_{M'} = L'_{s'} = F_{qp}(\log_{G'}(s'))$  coincides with  $F_M$ , since they are the Picard-Vessiot extensions of two adjoint differential systems. So, although  $\log_G$  and  $\log_{G'}$  have no direct relations, the fields which  $\log_G(s)$  and  $\log_{G'}(s')$  generate **over**  $F_{pq}$  are the same. Similarly, the structures of  $G$  and  $G'$  usually differ a lot, but the conclusions (i) and (ii) of the Main Lemma turn out to be invariant under this duality. See Case **(SC2)** of §6 for a concrete implementation of this remark.

iii) The above symmetry is best expressed in terms of the Poincaré bi-extension  $\mathcal{P}$ , resp.  $\mathcal{P}'$ , of  $E \times_S \hat{E}$ , resp.  $\hat{E} \times_S E$ , by  $\mathbb{G}_m$ . As recalled in the Introduction (cf. [13]), a section  $s$  of  $G = G_q$  above  $p$  corresponds to a trivialization of the  $\mathbb{G}_m$ -torsor  $(p, q)^*\mathcal{P} \simeq (q, p)^*\mathcal{P}'$ . Then, the inverse image  $\zeta$  of this trivialization under the uniformizing map

$$\mathbb{C}^3 \times \tilde{S} \rightarrow \mathcal{P}^{an}$$

of  $\mathcal{P}$  generates  $L$  over  $F_{pq}$  (here,  $\tilde{S}$  denotes the universal cover of  $S^{an}$ , for instance the Poincaré half-plane when  $S = X$  is a modular curve). This view-point turns the Main Lemma into a statement about the transcendency of  $\zeta$  over  $F_{pq}$ , and explain why the various types of special curves of the mixed Shimura variety  $\mathcal{P}/X$ , as encountered in §1.2, occur in its conclusion.

iv) (*Autocritique* on differential extensions) It would be interesting to pursue the study of this uniformizing map further, as it may lead to a simplification of the present proof of the Main Theorem, where the appeal to differential extensions would be replaced by an Ax-Lindemann statement, extending the recent results of Ullmo-Yafaev [30] and Pila-Tsimerman [24] to mixed Shimura varieties. See [28] for a perspective on both approaches.

## 6 Proof of the Main Lemma

The arguments will be of the same nature as in §4, appealing to Ax type results for constant groups, and to representation theory otherwise. As mentioned before the enunciation of the Main Lemma in §3.3, similar results

appear in [6], Propositions 4.a, 4.b and Theorem 2, but it seems better to gather them here into a full proof.

Consider the tower of Picard-Vessiot extensions drawn on the left part of the following picture :

$$\begin{array}{c}
 L \\
 \uparrow \\
 F_{pq} \\
 \swarrow \quad \searrow \\
 F_q \quad \quad F_p \\
 \nwarrow \quad \nearrow \\
 F \\
 \uparrow \\
 K
 \end{array}
 , \rho_{G,s}(\gamma) = \begin{pmatrix} 1 & {}^t\xi_q(\gamma) & \tau_s(\gamma) \\ 0 & \rho_E(\gamma) & \xi_p(\gamma) \\ 0 & 0 & 1 \end{pmatrix}, \quad \begin{array}{l} \tau_s : Gal_{\partial}(L/F_{pq}) \hookrightarrow \mathbb{C} \\ {}^t\xi_q : Gal_{\partial}(F_q/F) \hookrightarrow \mathbb{C}^2 \simeq \hat{\mathcal{V}} \\ \xi_p : Gal_{\partial}(F_p/F) \hookrightarrow \mathbb{C}^2 \simeq \mathcal{V} \\ \rho_E : Gal_{\partial}(F/K) \hookrightarrow SL_2(\mathbb{C}) \end{array}$$

For convenience, we recall from §5.1 that the field  $F = K(\omega_1, \omega_2, \eta_1, \eta_2)$  is the Picard-Vessiot extension of  $K$  given by the Picard-Fuchs equation  $\partial_{Lie\tilde{E}}(*) = 0$  for  $E/S$ , whose set of solutions we denote by  $\mathcal{V} \simeq \mathbb{C}^2$ . If  $E/S$  is isoconstant,  $F = K$ , while  $Gal_{\partial}(F/K) = SL_2(\mathbb{C})$  otherwise. The field  $F_p = F(u, \zeta(u))$  corresponds to the inhomogeneous equation attached to  $p$  (given by  $\partial_{Lie\tilde{E}}(\tilde{u}) = \partial \ell n_{\tilde{E}}(\tilde{p})$ , for any choice of a lift  $\tilde{p} \in \tilde{E}(S)$  of  $p$ ), while  $F_q = F(v, \zeta(v))$  is the field of periods of the semi-abelian scheme  $G/S$ , generated by the solutions of  $\partial_{Lie\tilde{G}}(*) = 0$ . As already said, its resemblance with  $F_p$  reflects a duality, witnessed by the dual  $\hat{\mathcal{V}}$  of  $\mathcal{V}$ . We fix a polarization of  $E/S$ , allowing us to identify  $E$  and  $\hat{E}$  (and in particular,  $q$  to a section of  $E/S$ ), but will keep track of this duality. The field of generalized periods of  $\{E, p, q\}$  is the compositum  $F_{pq}$  of  $F_p$  and  $F_q$ . Finally,

$$L = F_{pq}(\log_G s(\lambda)) = F_{pq}(\log_{\tilde{G}}(\tilde{s}(\lambda)))$$

is the Picard-Vessiot extension generated by the solutions of the 3rd order inhomogeneous equation  $\partial_{Lie\tilde{G}}(\tilde{U}) = \partial \ell n_{\tilde{G}}(\tilde{s})$ , where  $\tilde{s}$  is the pullback of  $\tilde{p}$  to  $\tilde{G}$  over  $s$ . The corresponding 4th order homogeneous system can be described as the Gauss-Manin connection attached to the smooth one-motive  $M$  over  $S$  given by the section  $s \in G(S)$ . The drawing in the middle is a representation  $\rho_{G,s}$  of the differential Galois group of  $L/K$ . The right part expresses that the coefficients of this representation become injective group homomorphisms on the indicated subquotients of  $Gal_{\partial}(L/K)$ .

We will again distinguish between several cases, depending on the position of  $p$  and  $q$  with respect to the projection  $E^\sharp$  to  $E$  of the Manin kernel  $\tilde{E}^\sharp$  of  $\tilde{E}$ . So,

$$E^\sharp = E_{tor} + \mathcal{E}_0(\mathbb{C}) = \begin{cases} E_{tor} & \text{if } E \text{ is not isoconstant;} \\ E_0(\mathbb{C}) & \text{if } E \simeq E_0 \times S \end{cases}$$

depending on whether the  $K/\mathbb{C}$ -trace  $\mathcal{E}_0$  of  $E$  vanishes or not. (In fact,  $E^\sharp$  is the Kolchin closure of  $E_{tor}$ , and is also called the *Manin kernel of  $E$* .) We denote by  $\hat{p}, \hat{q}$  the images of  $p, q$  in the quotient  $E/E^\sharp$ . Notice that the ring  $\mathcal{O} = \text{End}(E/S)$  still acts on this quotient.

We recall that we must here merely prove Part (A) of the Main Lemma. By contraposition, we assume that no constant translate of  $s$  is a Ribet section, or factors through a strict subgroup scheme of  $G/S$ , and must deduce that  $\log_G(s)$ , or equivalently,  $\log_{\tilde{G}}(\tilde{s})$ , is transcendental over  $F_{pq}$ .

**Case (SC1) :**  $\hat{q} = 0$

Assume first that  $E/S$  is not isoconstant. Then, this vanishing means that  $q$  is a torsion section, and after an isogeny,  $G = G_q$  is isomorphic to  $\mathbb{G}_m \times E$ . We have already proven Part (A) of the Main Lemma in this case, see Lemma 1 and the lines which follow. So, we can assume that  $E = E_0 \times S$  is constant, and the relation  $\hat{q} = 0$  now means that  $q$  is constant. So,  $G = G_0 \times S$  is a constant semi-abelian variety, and we can apply to its (constant) universal vectorial extension  $\tilde{G}_0$  the slight generalization of Ax's theorem given in [6], Proposition 1.b. Since we are assuming that no constant translate  $s - s_0, s_0 \in G_0(\mathbb{C})$ , of  $s$  factors through a strict subgroup scheme  $H$  of  $G$ , the relative hull  $G_s$  of  $s$  in the sense of [6], §1, is equal to  $G$ , and Proposition 1.b of *loc. cit.* implies that

$$\text{tr.deg.}(K(\log_{\tilde{G}}(\tilde{s}))/K) = \dim(\tilde{G}) = 3.$$

Now,  $F_q = F = K$  since  $E$  and  $q$  are constant, while  $K(\tilde{U}) = K(\tilde{u}, U) = F_p(\log_G(s))$  has transcendence degree  $\leq 1$  over  $F_p$ , which has transcendence degree  $\leq 2$  over  $K$ . So, both transcendence degrees must be maximal, and  $\log_G(s)$  is indeed transcendental over  $F_p = F_{pq}$ .

**Case (SC2) :**  $\hat{p} = 0$

This case is dual to the previous one, and the following preliminary remarks will simplify its study. The hypothesis made on  $s$  implies that  $p$  is not a

torsion section (otherwise, a multiple of  $s$  factors through  $\mathbb{G}_m$ ). So, we can assume that  $E = E_0 \times S$  is constant, and that  $p = p_0$  is a constant non-torsion section of  $E$ . In view of Case **(CS1)**, we can also assume that  $\hat{q} \neq 0$ , i.e. that  $q$  is not constant<sup>5</sup> (and in particular, not torsion). We now consider the smooth one-motive  $M = [M_s : \mathbb{Z} \rightarrow G]$  attached to the section  $s$  above  $p = p_0$ , and its Cartier dual  $M' = [M'_{s'} : \mathbb{Z} \rightarrow G']$ , where  $G'$  is the extension of  $\hat{E}$  by  $\mathbb{G}_m$  parametrized by  $p_0$ . In particular,  $G'$  is a constant and non isotrivial semi-abelian variety. By Remark 3.ii of §5, the field  $F_{p_0q}(\log_G(s)) = F_q(\log_G(s))$  coincides with the field  $F_{qp_0}(\log_{G'}(s')) = F_q(\log_{G'}(s'))$ . But the section  $s'$  of  $G'$  project to  $q$  in  $\hat{E}$ , which is not constant, so no constant translate of  $s'$  factors through  $\mathbb{G}_m$ . Finally,  $\mathbb{G}_m$  is the unique connected subgroup scheme of  $G'$ , since  $G'$  is non isotrivial. So, the constant semi-abelian variety  $G'$  and its section  $s'$  satisfy all the hypotheses of Case **(SC1)**. Therefore,  $\log_{G'}(s')$  is transcendental over  $F_q = F_{qp}$ , or equivalently,  $\log_G(s)$  is transcendental over  $F_{pq}$ .

In the next two cases, the proof of our transcendence claim can be derived from the following simple observation : the Lie algebra  $\mathfrak{u}_s$  of the unipotent radical of the image of  $\rho_{G,s}$  consists of matrices of the form  $X$  indicated below, where  $({}^t y, x) \in \text{Im}({}^t \xi_q, \xi_p) \subset \hat{\mathcal{V}} \times \mathcal{V}$ , and  $t \in \mathbb{C}$ , and for two such matrices

$$X = \begin{pmatrix} 0 & {}^t y & t \\ 0 & 0 & x \\ 0 & 0 & 0 \end{pmatrix}, X', \text{ we have } [X, X'] = \begin{pmatrix} 0 & 0 & t(X, X') \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

where  $t(X, X') = \langle y|x' \rangle - \langle y'|x \rangle$  depends only on the vectors  $x, y, x', y'$ . Here, the transposition and the scalar product represent the canonical anti-symmetric pairing  $\mathcal{V} \times \mathcal{V} \rightarrow \mathbb{C}$  provided by the chosen principal polarisation on  $E/S$ . Now,

### Case **(SC3)** : $\hat{p}$ and $\hat{q}$ are linearly independent over $\mathcal{O}$

As mentioned in Remark 2.(iv) of §4, the argument leading to Lemma 1, i.e. the sharpened form of André's theorem [1] given in [7] (or alternatively,

---

<sup>5</sup> It is worth noticing that this case **(SC2)** is the logarithmic analogue of the counterexample studied in [11], §5.3. It does not provide a counterexample to the Main lemma, whose "exponential" analogue would amount, in the notations of [11], to the equality  $\text{tr.deg.}(K(y)/K) = 1$ . In fact, the Theorem stated in §4, combined with Lemma 1 and with the conclusion of Case **(SC2)**, implies that  $y$  is transcendental over  $K_G^\sharp$ .



if  $E/S$  is isoconstant, the sharpened forms of Ax's theorem given in [12], Thm. 2, and in [6], Prop. 1.b) implies in this case that  $u, \zeta(u), v, \zeta(v)$  are algebraically independent over  $F$ . In other words, the homomorphism  $({}^t\xi_q, \xi_p) : Gal_{\partial}(F_{pq}/F) \rightarrow \hat{\mathcal{V}} \times \mathcal{V} \simeq \mathbb{C}^4$  is bijective, and any couple  $({}^ty, x)$  occurs in the Lie algebra  $\mathfrak{u}_s$ . Consequently, there exists  $X, X' \in \mathfrak{u}_s$  such that  $t(X, X') \neq 0$ , and  $\mathfrak{u}_s$  contains matrices all of whose coefficients, *except* the upper right one, vanish. Therefore, the homomorphism  $\tau_s$  is bijective,  $Gal_{\partial}(L/F_{pq}) \simeq \mathbb{C}$ , and  $tr.deg(L/F_{pq}) = 1$ . (Notice that this yields  $tr.deg(L/F) = 1 + 4 = 5$ .)

So, from now on, we can assume that  $\hat{q}$  and  $\hat{p}$  are linked by a unique relation over  $\mathcal{O}$ , which, considering multiples if necessary, we write in the shape

$$\hat{q} = \alpha\hat{p}, \alpha \in \mathcal{O}, \hat{p} \neq 0, \alpha \neq 0.$$

We denote by  $\bar{\alpha}$  the complex conjugate of  $\alpha$ , which represents the image of  $\alpha \in End(E/S)$  under the Rosati involution attached to the chosen polarization. We first deal with non antisymmetric relations, in the sense of §2.3.

**Case (SC4) :**  $\hat{q} = \alpha\hat{p}$ , where  $\bar{\alpha} \neq -\alpha$  and  $\hat{p} \neq 0$

Lifted to  $E$  and up to an isogeny, this relation reads  $q = \alpha p + p_0$ , where  $p_0 \in E_0(\mathbb{C})$  is a constant section, equal to 0 if  $E/S$  is not isoconstant. Then,  $F_q = F_p$ , and more precisely,  $\xi_q = \alpha \circ \xi_p$ . In other words, the coefficients of the matrices  $X$  in  $\mathfrak{u}_s$  satisfy the relation  $y = \alpha x$ , for the natural action of  $\mathcal{O}$  on  $\mathcal{V}$ . Since  $p \notin E_0(\mathbb{C})$ , Lemma 1 implies that  $\xi_p$  is bijective, and any  $x \in \mathcal{V}$  occurs in the Lie algebra  $\mathfrak{u}_s$ . Recall that  $\langle | \rangle$  is antisymmetric, and that the adjoint of the endomorphism of  $\mathcal{V}$  induced by an isogeny  $\alpha \in \mathcal{O}$  is its Rosati image  $\bar{\alpha}$ . For any  $x, x'$  in  $\mathcal{V}$  occurring in matrices  $X, X'$  and such that  $\langle x|x' \rangle \neq 0$ , we then have :

$$\begin{aligned} t(X, X') &= \langle \alpha x|x' \rangle - \langle \alpha x'|x \rangle = \langle \alpha x|x' \rangle - \langle x'|\bar{\alpha}x \rangle \\ &= \langle \alpha x|x' \rangle + \langle \bar{\alpha}x, x' \rangle = \langle (\alpha + \bar{\alpha})x|x' \rangle \neq 0 \end{aligned}$$

since  $\alpha + \bar{\alpha}$  is a non-zero integer. We conclude as in Case (SC3) that  $Gal_{\partial}(L/F_{pq}) \simeq \mathbb{C}$ , and  $tr.deg.(L/F_{pq}) = 1$ . (Here, this yields  $tr.deg(L/F) = 1 + 2 = 3$ .)

The remaining cases concerns antisymmetric relations of the type  $\hat{q} = \alpha\hat{p}$ , with  $\hat{p} \neq 0$  and a non zero purely imaginary  $\alpha = -\bar{\alpha}$ . (In particular, the CM

elliptic scheme  $E/S$  must be isoconstant and so, Theorem 2 is now already proven under its Conditions (i) or (ii)). We first treat the case when  $q$  and  $p$  themselves are antisymmetrically related.

**Case (SC5) :**  $q = \alpha p$ , where  $\bar{\alpha} = -\alpha \neq 0$  and  $\hat{p} \neq 0$

This is the only case where a Ribet section of  $G/S$  exists above the section  $p \in E(S), p \notin E_0(\mathbb{C})$ . Denote by  $s_R$  this (essentially unique) Ribet section.

**Lemma 3.** *Let  $s_R$  be the Ribet section of  $G/S$ . Then,  $\log_G(s_R)$  is defined over  $F_{pq}$ .*

*Proof :* let  $L_R = F_{pq}(\log_G(s_R))$  be the field generated over  $F_{pq}$  by  $\log_G(s_R)$ . Since the differential Galois group  $Gal_{\partial}(L_R/F_{pq})$  injects via  $\tau_{s_R}$  into a vectorial group  $\mathbb{C}$ ,  $\log_G(s_R)$  is either transcendental or rational over  $F_{pq}$ . Assume that it is transcendental. Then, by Proposition 4, the surface  $\mathcal{S}$  attached to  $s_R$  contains no algebraic curve, and the whole reduction of the Main Theorem to the study of  $\mathcal{S}$  given in §3.1 and §3.2 implies that  $s_R$  admits only finitely many torsion values. But this contradicts the main result of [8], an analytic proof (or version) of which will be found in Appendix I, §7.2. In fact, the explicit formulae given there directly show that  $L_R = F_{pq}$ , cf. Proposition 5.(iii).

We now come back to our section  $s$ , which we assumed *not* to be a Ribet section of  $G$ , and more accurately, such that no constant translate  $s - s_0$  of  $s$  is a Ribet section. Since  $s$  and  $s_R$  project to the same section  $p$  of  $E/S$ , there exists a section  $\delta \in \mathbb{G}_m(S)$ , i.e. a rational function in  $K^*$ , such that  $s = s_R + \delta$ . The assumption on  $s$  implies that  $\delta \notin \mathbb{C}^*$  is not constant. Set  $\ell = \log_{\mathbb{G}_m}(\delta)$ . Then,  $\log_G(s) = \log_G(s_R) + \ell$ , and the lemma implies that that  $F_{pq}(\log_G(s)) = F_{pq}(\ell)$ , which is equal to  $F_p(\ell)$ , since  $\hat{q} = \alpha\hat{p}$ . By Lemma 1,  $F_p(\ell)$  has transcendence degree 1 over  $F_p$ , so  $\log_G(s)$  too is transcendental over  $F_{pq}$ .

**Case (SC6) :**  $q = \alpha p + p_0$ , where  $\bar{\alpha} = -\alpha \neq 0$  and  $p_0 \in E_0(\mathbb{C}), p_0 \notin E_{0,tor}$

Fixing an  $\alpha$ -division point  $p'_0$  of  $p_0$  in  $E_0(\mathbb{C})$ , we have  $q = \alpha(p - p'_0)$ , and there exists an (essentially unique) Ribet section  $s'_R$  of  $G/S$  above  $p' := p - p'_0$ . Then, the section  $s' := s - s'_R$  of  $G/S$  projects to  $\pi(s') = p - p' = p'_0$  in  $E_0(\mathbb{C}) \subset E(S)$ . Furthermore,  $F_p = F_{p'} = F_q$  since  $p'_0$  is constant, and since  $\log_G(s'_R)$  is defined over  $F_{p'q} = F_{pq} = F_q$  by Lemma 3, we deduce that

$\log_G(s') = \log_G(s) - \log_G(s'_R)$  generates over the field  $F_{pq} = F_{p'q} = F_q = F_{p'_0q}$  the same field as  $\log_G(s)$ . We are therefore reduced to showing that given  $\hat{q} \neq 0$  and a section  $s'$  of  $G/S$  projecting to a constant non-torsion section  $p'_0 \in E_0(\mathbb{C})$ , then  $\log_G(s')$  is transcendental over  $F_{p'_0q}$ . But this is exactly what we proved in Case **(SC2)** !

This concludes the proof of the Main Lemma, hence of the Main Theorem.

## 7 Appendix I

### 7.1 Analytic description of the semi-abelian logarithm

Let  $G/S$  be a *non isotrivial* extension of an elliptic scheme  $E/S$  by  $\mathbb{G}_m$  and let  $s$  be a section of  $G/S$ . The aim of this Appendix is to give an explicit formula for its local logarithm  $\log_G(s)$  in terms of the Weierstrass functions  $\wp_\lambda, \zeta_\lambda, \sigma_\lambda$ , in parallel with that of §4 in for products. We recall the notations of §5.1. In particular, we set  $p = \pi(s) \in E(S)$ ,  $u = \log_E(p)$ , For simplicity, we will work over the generic point of  $S$ , consider  $G$  as a semi-abelian variety over the field  $K = \mathbb{C}(S)$ , and often drop the variable  $\lambda$  indexing the Weierstrass functions and their (quasi-)periods  $\omega_1, \omega_2, \eta_1, \eta_2$ .

By Weil-Rosenlicht-Barsotti, the algebraic group  $G$ , viewed as a  $\mathbb{G}_m$ -torsor, defines a line bundle over  $E$  of degree 0, admitting a rational section  $\beta$  with divisor  $(-q) - (0) \in \hat{E}$ , which we identify with the point  $q \in E$  (the sign is admittedly not standard, but it will make the formulae symmetric in  $p$  and  $q$ ). By assumption,  $q$  is not a torsion point, and we set  $v = \log_E(q)$ . We further assume that  $p \neq 0$  and  $p + q \neq 0$ .

The rational section  $\beta$  provides a birational isomorphism  $G \dashrightarrow \mathbb{G}_m \times E$  and (after a shift away from 0) an isomorphism  $LieG \simeq Lie\mathbb{G}_m \oplus LieE$ . The 2-cocycle which describes the group law on the product ([29], VII.5) is a rational function on  $E \times E$ , expressed in terms of  $\sigma$ -functions by  $\frac{\sigma(z+z'+v)\sigma(z)\sigma(z')\sigma(v)}{\sigma(z+z')\sigma(z+v)\sigma(z'+v)}$ . Therefore, the exponential morphism  $exp_G$  is represented by the map

$$(LieG)^{an}(\Lambda) \ni \begin{pmatrix} t(\lambda) \\ z(\lambda) \end{pmatrix} \mapsto \begin{pmatrix} f_{v(\lambda)}(z(\lambda)) e^{t(\lambda)} \\ \wp_\lambda(z(\lambda)) \end{pmatrix} \in G^{an}(\Lambda)$$

where

$$f_v(z) = \frac{\sigma(v+z)}{\sigma(v)\sigma(z)} e^{-\zeta(v)z}$$

is a meromorphic theta function for the line bundle  $\mathcal{O}_E((-q) - (0))$ , whose factors of automorphy are given by  $e^{-\kappa_v(\omega_i)}$  (opposite of the multiplicative quasi-periods), with

$$\kappa_v(\omega_i) = \zeta(v)\omega_i - \eta_i v, \text{ for } i = 1, 2.$$

The occurrence of the trivial theta function  $e^{-\zeta(v)z}$  in  $f_v$  is due to the condition  $d_0(\exp_G) = id_{LieG}$ . The logarithmic form  $\frac{df_v}{f_v} = (\zeta(v+z) - \zeta(v) - \zeta(z))dz = \frac{1}{2} \frac{\wp'(z) - \wp'(v)}{\wp(z) - \wp(v)} dz$  is the pullback under  $\exp_E$  of the standard differential form of the 3rd kind on  $E$  with residue divisor  $-1.(0) + 1.(-q)$ .

Under this description, the section  $s$  of  $G/S$  under study and its logarithm  $\log_G(s)$  are given by

$$s = \begin{pmatrix} \delta_s \\ p \end{pmatrix}, \quad U := \log_G(s) = \begin{pmatrix} -g(u, v) + \zeta(v)u + \ell_s \\ u \end{pmatrix}$$

where  $\delta_s := s - \beta(p) \in K^*$  is a rational function on  $S$ , depending only on  $s$  (and on the choice of the section  $\beta$ ), for which we set  $\ell_s = \log_{\mathbb{G}_m}(\delta_s)$ , and (cf. the formulae in [6], up to signs) :

$$g_\lambda(u, v) = \log\left(\frac{\sigma_\lambda(u+v)}{\sigma_\lambda(v)\sigma_\lambda(u)}\right).$$

This is the Green function mentioned in §5.1.

The  $\mathbb{Z}$ -local system of periods  $\Pi_G$  of  $G^{an}/\Lambda$  which was introduced in §3.3 admits the basis

$$\varpi_0(\lambda) = \begin{pmatrix} 2\pi i \\ 0 \end{pmatrix}, \quad \varpi_1(\lambda) = \begin{pmatrix} \kappa_v(\lambda)(\omega_1(\lambda)) \\ \omega_1(\lambda) \end{pmatrix}, \quad \varpi_2(\lambda) = \begin{pmatrix} \kappa_v(\lambda)(\omega_2(\lambda)) \\ \omega_2(\lambda) \end{pmatrix}.$$

We can now describe the various extensions of  $F = K(\omega_1, \omega_2, \eta_1, \eta_2)$  appearing in §5 for a non isotrivial extension  $G$ . In view of the Legendre relation

$$2\pi i = \eta_1 \omega_2 - \eta_2 \omega_1 \in K^*,$$

the periods of  $G$  generate over  $F$  the field

$$F_G = F_G^{(2)} = F(\kappa_v(\omega_1), \kappa_v(\omega_2)) = F(v, \zeta(v)) := F_q,$$

while the field generated over  $F_{pq}$  by  $\log_G(s)$  satisfies

$$L = F_{pq}(\log_G(s)) = F(u, \zeta(u), v, \zeta(v), -g(u, v) + \zeta(v)u + \ell_s) = F_{pq}(\ell_s - g(u, v))$$

## 7.2 Analytic description of the Ribet sections

We now present the analytic description of the Ribet sections promised in §1.1. Assume that  $E = E_0 \times S$  is a *constant elliptic scheme with complex multiplications*, that  $q$  is not constant (i.e.  $G$  is semi-constant), and that  $p$  and  $q$  are *antisymmetrically related* in the sense of §2. So, their logarithms  $u(\lambda), v(\lambda)$  are non constant holomorphic functions on  $\Lambda$ , and satisfy :  $v = \alpha u$  modulo  $\mathbb{Q}\omega_1 \oplus \mathbb{Q}\omega_2$  for a totally imaginary non zero complex multiplication  $\alpha \in \mathcal{O} \otimes \mathbb{Q}$ . For simplicity, we will assume that this relation takes the form

$$v = \alpha u, \alpha \in \mathcal{O}, \alpha = -\bar{\alpha} \neq 0, u, v \notin \mathbb{C},$$

and that  $\alpha$  lies in  $2\mathcal{O} \subset 2(\mathbb{Z} \oplus \mathbb{Z}\tau)$ , where  $\tau := \frac{\omega_2}{\omega_1} \in \mathcal{O} \otimes \mathbb{Q}$ ,  $\Pi_E = \omega_1(\mathbb{Z} \oplus \mathbb{Z}\tau)$ .

We fix a rational section  $\beta$  of  $\pi : G \rightarrow E$  as above, so, any section  $s$  of  $G/S$  projecting to  $p \in E(S)$  is uniquely expressed as a couple  $(\delta_s, p)$ , for some rational function  $\delta_s \in K^*$ . In particular, the Ribet section  $s_R$  of  $G/S$  is given by  $(\delta_R, p)$ , where  $\delta_R \in K^*$  is well defined, at least up to a root of unity. Since  $E$  is constant, the index  $\lambda$  truly disappears from the notations  $\omega_{1,2}, \sigma, g, \dots$ , which are now attached to  $E_0$ . We must express  $\delta_R$  in terms of these functions, evaluated at  $u(\lambda)$ .

By the theory of complex multiplication (see for instance [18], Appendix I), there exists an explicit complex number  $s_2$  (actually algebraic over  $\mathbb{Q}$ , and given by Hecke's non-holomorphic modular form of weight 2), such that the quasi-periods of  $\zeta$  satisfy :

$$\eta_2 - s_2\omega_2 = \bar{\tau}(\eta_1 - s_2\omega_1), \text{ where } \omega_2 = \tau\omega_1.$$

From this and the functional equation of  $\sigma$ , one infers that the function  $\theta(z) = \sigma(z)e^{-\frac{1}{2}s_2z^2}$  satisfies, for  $N(\gamma) = \gamma\bar{\gamma}$  :

$$\forall \gamma \in \mathcal{O}, \left( \frac{\theta(\gamma z)}{\theta(z)^{N(\gamma)}} \right)^2 = \gamma^2 \prod_{e \in E[\gamma], e \neq 0} (\wp(z) - \wp(e)).$$

Since  $N(\alpha+1) - N(\alpha) - 1 = \alpha + \bar{\alpha} = 0$ , we deduce from the parity assumption on  $\alpha$  that the meromorphic function

$$\delta_{\bar{R}} := e^{g(u,v) - s_2uv} = \frac{\sigma(u+v)}{\sigma(v)\sigma(u)} e^{-s_2uv}, \text{ where } v = \alpha u,$$

is a rational function of  $\wp(u(\lambda)), \wp'(u(\lambda))$ , hence lies in  $K^*$ . We then have:

**Proposition 5.** For  $q = \alpha p$ , consider the section  $s_{\bar{R}} = (\delta_{\bar{R}}, p)$  of  $G = G_q$  above  $p$ . Then,

- i) up to a root of unity,  $s_{\bar{R}}$  is equal to the Ribet section  $s_R$  of  $G$  above  $p$ ;
- ii) for any  $\bar{\lambda} \in S$  such that  $p(\bar{\lambda})$  is a torsion point of  $E_0$ , say of order  $n$ , the point  $s_{\bar{R}}(\bar{\lambda})$  is a torsion point of  $G_{\bar{\lambda}}$ , of order dividing  $n^2$ ;
- iii) the logarithm of  $s_{\bar{R}}$  is given by  $(\zeta(v)u - s_2uv, u)$ ; in particular,  $F_{pq}(\log_G(s_R)) = F_{pq}$ .

(Following the conventions of §2.1, the bar over a  $\lambda$  indicates a value of  $\lambda$ , not complex conjugation as for  $\tau, \alpha$  and  $\gamma$ .)

*Proof.* - i) In view of Corollary 3, applied to  $s = s_{\bar{R}}$  and  $s^\dagger = s_R$ , this first assertion is an immediate corollary of the second statement (ii). Indeed, by Proposition 1.(iv), the set  $S_\infty^E$  is infinite since  $p$  is not constant, so (ii) states in particular that  $s_{\bar{R}}$  lifts infinitely many torsion values of  $p$  to torsion points of  $G$ .

iii) By the formulae of the previous subsection, the first coordinate of  $\log_G(s_{\bar{R}})$  is equal to

$$-g(u, v) + \zeta(v)u + \log(\delta_{\bar{R}}) = \zeta(v)u - s_2uv.$$

The last assertion then immediately follows from (i). It is timely to point out here that the logarithmic derivative  $\tilde{\zeta}(z) = \zeta(z) - s_2z$  of  $\theta$  satisfies :  $\tilde{\zeta}(\alpha z) - \bar{\alpha}\tilde{\zeta}(z) \in \wp'(z)\mathbb{C}(\wp(z))$ .

ii) We must show that for any  $\bar{\lambda}$  such that  $u(\bar{\lambda}) = \frac{1}{n}\omega$  for some integer  $n$  and some period  $\omega$  of  $E$ , with corresponding additive and multiplicative quasi-periods  $\eta(\omega), \kappa_v(\omega)$ , and with  $v(\bar{\lambda}) = \alpha u(\bar{\lambda}) = \frac{1}{n}\alpha\omega$ , the number

$$\zeta(v(\bar{\lambda}))u(\bar{\lambda}) - s_2u(\bar{\lambda})v(\bar{\lambda}) - \frac{1}{n}\kappa_{v(\bar{\lambda})}(\omega) = \frac{1}{n^2}\eta(\omega)\alpha\omega - \frac{1}{n^2}s_2\alpha\omega^2 = \frac{1}{n^2}\alpha(\eta(\omega) - s_2\omega)\omega$$

is a rational multiple of  $2\pi i$ , and more precisely an integral multiple of  $\frac{1}{n^2} \cdot 2\pi i$ . Indeed, this number is the first coordinate of the vector  $\log_G(s_{\bar{R}}(\bar{\lambda})) - \frac{1}{n}\varpi$ , where  $\varpi$  lies in the  $\mathbb{Z}$ -module  $\Pi_G$  of periods of  $G$  above the period  $\omega \in \Pi_E$ . Since  $\log_E(p(\bar{\lambda})) = u(\bar{\lambda}) = \frac{1}{n}\omega$ , the second coordinate of this vector vanishes, so  $\log_G(s_{\bar{R}}(\bar{\lambda}))$  itself will lie  $\frac{1}{n^2}\mathbb{Z}\varpi_0 + \frac{1}{n}\Pi_G$ .

By additivity, it suffices to check the assertion when  $\omega = \omega_1$  and  $\omega_2 = \tau\omega_1$ . By the classical Legendre relation, combined with the CM relation

$\eta_2 - s_2\omega_2 = \bar{\tau}(\eta_1 - s_2\omega_1)$ , the number under study is equal to  $\frac{1}{n^2} \frac{\alpha}{\bar{\tau}-\tau} 2\pi i$  when  $\omega = \omega_1$ . Since  $\alpha$  is a purely imaginary element of  $\mathbb{Q}(\tau)$ , this is indeed a rational multiple of  $2\pi i$ , and even an integral one since  $\alpha \in \mathbb{Z} \oplus 2\mathbb{Z}\tau$ . For  $\omega = \omega_2$ , the same computation yields  $\frac{1}{n^2} \tau \bar{\tau} \frac{\alpha}{\bar{\tau}-\tau} 2\pi i$ , with same conclusion since  $\alpha\tau \in 2\mathbb{Z} \oplus \mathbb{Z}\tau$ .

**Remark 4.** i) Proposition 5.(ii) by itself suffices to give a counterexample to the Relative Manin-Mumford conjecture, independently of [8]. So, we could have used  $s_{\bar{R}}$  as a definition of the Ribet section. Granted the equality  $s_{\bar{R}} = s_R$ , this gives a new proof of Edixhoven's sharpening in [8], Appendix, according to which the Ribet section  $s_R$  lifts torsion points of order  $n$  to torsion points of order dividing, and often equal to,  $n^2$ .

ii) Instead of a parity assumption, one can merely assume that  $\alpha = \beta - \bar{\beta}$  for some  $\beta \in \mathcal{O}$ . As shown in greater generality in [10], the  $n$ -th root of unity  $ns_R(\bar{\lambda})$ , for  $\bar{\lambda} \in S_n^E$ , can then be expressed in terms of the Weil pairing  $e_n(\beta p(\bar{\lambda}), p(\bar{\lambda}))$ . The formula above also implies this sharpening, since it shows that  $ns_{\bar{R}}(\bar{\lambda})$  is then equal to  $\exp(\frac{1}{n}(\eta(\omega)\omega' - \eta(\omega')\omega))$ , where  $\omega' = \beta\omega$ .

## 8 Appendix II : application to Pell equations

In [20], the relative Manin-Mumford conjecture is proven for simple abelian surface schemes, and this implies the following corollary : consider the family of sextic polynomials  $D_\lambda(x) = x^6 + x + \lambda$ , where  $\lambda$  is a complex parameter. Then, there are only finitely many  $\bar{\lambda} \in \mathbb{C}$  such that the functional Pell equation  $X^2 - D_{\bar{\lambda}}(x)Y^2 = 1$  admits a solution in polynomials  $X, Y \in \mathbb{C}[x]$ ,  $Y \neq 0$ ; see also [31], III. 4.5 for connections with other problems and a proof of the deduction from RMM . The involved abelian surface  $A/\mathbb{C}(\lambda)$  is the jacobian of the (normalized) relative hyperelliptic curve  $C : y^2 = x^6 + x + \lambda$ , and RMM is applied to the section  $s$  of  $A$  defined by the linear equivalence class of the relative divisor  $(\infty_+) - (\infty_-)$  on  $C$ .

Following a suggestion of the second author, we may treat in the same way the case of a sextic  $D_\lambda(x) = (x - \rho(\lambda))^2 Q_\lambda(x)$  having a *squared linear factor*, i.e. a generic double root  $\rho(\lambda)$  for some algebraic function  $\rho(\lambda)$ , now applying the Main Theorem of the present paper to a quotient  $G = G_\rho$  of the generalized jacobian of the corresponding semi-stable relative sextic curve  $C$ . This  $G_\rho$  is an extension by  $\mathbb{G}_m$  of an elliptic curve  $E$  (over  $\mathbb{C}(\lambda)$ ), where  $E$  is the jacobian of the (normalized) relative quartic  $\tilde{C}$  with equation  $v^2 = Q_\lambda(u)$ ,

and RMM may be applied to the section  $s$  of  $G_\rho$  defined by the class of the relative divisor  $(\infty_+) - (\infty_-)$  on  $\tilde{C}$ , for the strict linear equivalence attached to the node of  $C$  at  $x = \rho(\lambda)$  (see [29], V.2, [8], Appendix). When the quartic  $Q_\lambda(x)$  does not depend on  $\lambda$  (or more generally, when  $E$  is isoconstant), no result of interest seems to appear. As an illustration of the opposite case, we will here consider the family of quartics

$$Q_\lambda(x) = x^4 + x + \lambda.$$

From the analysis in [20], [31] recalled below, we derive that the set  $\Lambda_Q$  of complex numbers  $\bar{\lambda}$  such that the Pell equation  $X^2 - Q_{\bar{\lambda}}(x)Y^2 = 1$  has a solution in polynomials  $X, Y \in \mathbb{C}[x]$ ,  $Y \neq 0$ , is infinite. The solutions for each such  $\bar{\lambda}$  form a sequence  $(X_{\bar{\lambda},n}, Y_{\bar{\lambda},n})_{n \in \mathbb{Z}}$  of polynomials in  $\mathbb{C}[x]$ . Our result is that for the  $\rho(\lambda)$  considered in the following statements, only finitely many of the polynomials  $Y_{\bar{\lambda},n}(x)_{\bar{\lambda} \in \Lambda_Q, n \in \mathbb{Z}}$  admit  $x = \rho(\bar{\lambda})$  among their roots. In other words :

**Theorem 4.** *i) (Case  $\rho(\lambda) = 0$ .) There are only finitely many complex numbers  $\bar{\lambda}$  such that the equation  $X^2 - x^2 Q_{\bar{\lambda}}(x)Y^2 = 1$  admits a solution in polynomials  $X, Y \in \mathbb{C}[x]$ ,  $Y \neq 0$ .*

*ii) (Case  $\rho(\lambda) = \frac{4y(\frac{1}{2}p_W + e_3) - 1}{8x(\frac{1}{2}p_W + e_3)} \in \mathbb{C}(\lambda)^{alg}$ , with notations explained below.)*

*There are only finitely many complex numbers  $\bar{\lambda}$  such that the equation  $X^2 - (x - \rho(\bar{\lambda}))^2 Q_{\bar{\lambda}}(x)Y^2 = 1$  admits a solution in polynomials  $X, Y \in \mathbb{C}[x]$ ,  $Y \neq 0$ .*

In spite of their similarity, these two statements cover different situations : in (i), the extension  $G_\rho$  is not isotrivial, and the theorem is a corollary of Theorem 2. On the contrary, (ii) illustrates the case of an isotrivial extension  $G_\rho$ , and follows from Theorem 1. In fact, we believe that on combining these two cases of our Main Theorem, Theorem 4 will hold for *any* choice of the algebraic function  $\rho(\lambda)$ .

The algebraic function  $\rho(\lambda)$  of Case (ii) can be described as follows. Consider the Weierstrass model  $(W_E) : y^2 = 4x^3 - \lambda x + (1/16)$  of the elliptic curve  $E/\mathbb{C}(\lambda)$ , with its standard group law, and the relative point  $p_W = (0, -\frac{1}{4})$  on  $(W_E)$ , which is generically of infinite order (it can be rewritten as the point  $p_{\tilde{W}} = (0, -1)$  on the curve  $(\tilde{W}_E) : Y^2 = X^3 - 4\lambda X + 1$ , and is not torsion at  $\lambda = \frac{1}{4}$ ). Then, the 2-division points of  $p_W$  are the four points of  $(W_E)$  :

$$\frac{1}{2}p_W = \left( \frac{1}{8}m_\lambda^2, -\frac{1}{8}m_\lambda^3 + \frac{1}{4} \right), \quad \text{where } m_\lambda \text{ is a root of } m^4 - 8m + 16\lambda = 0.$$



Choose one of the two roots  $m_\lambda$  which is real when  $\bar{\lambda} = \frac{1}{4}$ , and call the corresponding point  $\frac{1}{2}p_W(\lambda)$ . Further, choose one of the two points of order 3 on  $W_E$  which is real when  $\bar{\lambda}$  is real, and call it  $e_3(\lambda)$ . Computing the  $x$  and  $y$  coordinates of the relative point  $\frac{1}{2}p_W + e_3$  on  $W_E$  then provides the function  $\rho(\lambda)$  appearing in Case (ii).

In a more enlightening way, let in general  $p(\lambda)$  be the section of  $E$  defined by the class of the divisor  $(\infty_+) - (\infty_-)$  on  $\tilde{C}$ , for the standard linear equivalence of divisors. Then,  $p$  is the projection to  $E$  of the section  $s$  of  $G_\rho$  defined (via strict equivalence) above, and one checks that  $p$  is not a torsion section. By [20], [31] (see also [15], Prop. 3.1), the Pell equation for  $Q_{\bar{\lambda}}(x)$  has a non trivial solution if and only if  $p(\bar{\lambda})$  is a torsion point on  $E_{\bar{\lambda}}$ , i.e.  $\bar{\lambda} \in S_\infty^E$  in the notations of §2. Similarly, the Pell equation for  $(x - \rho(\bar{\lambda}))^2 Q_{\bar{\lambda}}(x)$  has a non trivial solution if and only if  $s(\bar{\lambda})$  is a torsion point of  $G_{\rho(\bar{\lambda})}$ , i.e.  $\bar{\lambda} \in S_\infty^{G_\rho}$ . Furthermore, the section  $q(\lambda)$  of  $\hat{E}$  parametrizing the extension  $G_\rho$  is represented the (standard) equivalence class of the divisor  $(q_+) - (q_-)$  on  $\tilde{C}$ , where  $q_\pm(\lambda)$  is the section  $(\rho(\lambda), \pm Q_\lambda^{1/2}(\rho(\lambda)))$  of  $\tilde{C}$ . Now, in the first case  $\rho(\lambda) = 0$ ,  $q$  is a non torsion section, i.e.  $G_\rho$  is a non isotrivial extension of the non isoconstant elliptic scheme  $E$ , and since  $p$  is not torsion, Theorem 2 implies that  $S_\infty^{G_\rho}$  is finite. On the other hand, Case (ii) is built up in such a way that  $q$  has finite order (equal to 3), so that  $G_\rho$  is now isogenous to  $\mathbb{G}_m \times E$ . But one can check (by specializing at the real number  $\bar{\lambda} = \frac{1}{4}$ ) that the projection of the section  $s$  to the  $\mathbb{G}_m$  factor is not a root of unity, so  $s$  does not factor through a translate of  $E$ . Since  $p$  is not torsion either, Theorem 1 now provides the finiteness of  $S_\infty^{G_\rho}$ .

## References

- [1] Y. André : Mumford-Tate groups of mixed Hodge structures and the theorem of the fixed part; *Compositio Math.*, 82 (1992), 1-24.
- [2] Y. André : Shimura varieties, subvarieties and CM points; lectures notes, Taiwan, 2001.
- [3] J. Ax : Some topics in differential algebraic geometry I : Analytic subgroups of algebraic groups; *Amer. J. Maths*, 94, 1972, 1195-1204.
- [4] J. Ayoub : Une version relative de la conjecture des périodes de Kontsevich-Zagier; *Annals of Math.*, to appear.

- [5] D. Bertrand : series of papers, starting with: Extensions de  $D$ -modules et groupes de Galois différentiels; Springer LN 1454, 1990 (Trento 1989), 125–141; Un analogue différentiel de la théorie de Kummer, W. de Gruyter 1992 (Luminy 1990), 39-49.
- [6] D. Bertrand : Schanuel’s conjecture for non-isoconstant elliptic curves over function fields; London MS LN 349, 2008, 41-62.
- [7] D. Bertrand : Théories de Galois différentielle et transcendance; Ann. Inst. Fourier, 59, 2009, 2773–2803. See also: Galois descent in Galois theories, Sémin. & Congrès 23, SMF, 2011, 1-24.
- [8] D. Bertrand : Special points and Poincaré bi-extensions; with an Appendix by Bas Edixhoven; ArXiv 1104.5178v1 .
- [9] D. Bertrand : Unlikely intersections in Poincaré biextensions over elliptic schemes; to appear in NDJFL, 2013 (= Oléron Proceedings).
- [10] D. Bertrand, B. Edixhoven : Pink’s conjecture, Poincaré biextensions and generalized Jacobians; in preparation.
- [11] D. Bertrand, A. Pillay : A Lindemann-Weierstrass theorem for semialbanian varieties over function fields; J. Amer. MS, 23, 2010, 491-533.
- [12] D. Brownawell, K. Kubota : Algebraic independence of Weierstrass functions; Acta Arithm. 33, 1977, 111-149.
- [13] P. Deligne : Théorie de Hodge III; Publ. math. IHES, 44, 1974, 5-77.
- [14] P. Habegger : Special points on fibered powers of elliptic surfaces; ArXiv 1110.1908, to appear in J. r. angew. Math.
- [15] F. Hazama : Twists and generalized Zolotarev polynomials; Pacific J. Maths; 203, 2002, 379-393.
- [16] M. Hindry : Autour d’une conjecture de Serge Lang; Invent. math. 94, 1988, 575-603.
- [17] O. Jacquinet, K. Ribet : Deficient points on extensions of abelian varieties by  $\mathbb{G}_m$ ; J. Number Th., 25, 1987, 133-151.

- [18] D. Masser : *Elliptic functions and transcendence*; Springer LN 437 (1975).
- [19] D. Masser, U. Zannier : series of papers, starting with : Torsion anomalous points and families of elliptic curves (CRAS, 346, 2008, 491-494, Amer. J. Math, 132, 2010, 1677-1691), Torsion points on families of squares of elliptic curves, Math. Ann., 2012, 352, 453-484.
- [20] D. Masser, U. Zannier : Torsion points on families of simple abelian surfaces and Pell's equation over polynomial rings; preprint, Nov. 2012.
- [21] J. Pila : Rational points on a subanalytic surface; Ann. Fourier 55, 2005, 1501-1515. See also E. Bombieri - J. Pila (Duke MJ 59, 1989, 491-533) for curves, and J. Pila - A. Wilkie (Duke MJ, 133, 2006, 591-616) in higher dimensions.
- [22] J. Pila : Rational points of definable sets and results of André-Oort–Manin-Mumford type; IMRN 13, 2009, 2476-2507.
- [23] J. Pila : O-minimality and the André-Oort conjecture for  $\mathbb{C}^n$ ; Annals Math.173 (2011), 1779-1840.
- [24] J. Pila, J. Tsimerman : Ax-Lindemann for  $\mathcal{A}_g$  ; ArXiv 1206.2663v1 .
- [25] J. Pila, U. Zannier : Rational points in periodic analytic sets and the Manin- Mumford conjecture; Rend. Lincei, Mat. Appl. 19 , 2008, 149–162.
- [26] A. Pillay : Notes on log-differential equations; preprint, May 2012.
- [27] R. Pink : A common generalization of the conjectures of André-Oort, Manin-Mumford, and Mordell-Lang; preprint, 2005.
- [28] T. Scanlon : The Zilber-Pink conjecture: diophantine geometry, functional transcendence, differential algebra and o-minimality; lecture notes, Ravello, 2013.
- [29] J-P. Serre : *Groupes algébriques et corps de classes*; Hermann, 1959.
- [30] E. Ullmo, A. Yafaev : Hyperbolic Ax-Lindemann theorem in the cocompact case; ArXiv 1209.0939v1 .

- [31] U. Zannier : *Some Problems of Unlikely Intersections in Arithmetic and Geometry*, (with appendices by David Masser); Ann. Maths St. 181, 2012, Princeton UP.

*Adresses of authors:* D. B. (Paris), D. M. (Basel), A. P. (Leeds), U. Z. (Pisa).