


Math 10250 Activity 23: Second Derivative Tests (Section 4.2)

GOAL: To study how the graph of a given $f(x)$ "bends", and how these features of the graph are described by $f'(x)$ and $f''(x)$.

► **The second derivative test for concavity**

Example 1 Water is filling up each of the following vessels at a constant rate of $1 \text{ cm}^3/\text{sec}$.



Let h be the height of the water level in the vessel at time t .

a. Sketch the graphs of h versus t for Vessels A and B in the axes for Figure 1. Indicate which graph belong to A and which to B.



Figure 1





Figure 2

b. Sketch the graph of h versus time t for Vessel C in the axes for Figure 2.

c. Comment on how the "bending" (up or down) of the graph changes with $h'(t)$. Mark on the graph where the "bending" changes.



We now introduce terminologies that describe the "bending" of a graph.

Case 1: For $a < x < b$, slope of the graph $f(x)$ is **increasing** as x increases i.e. $f'(x)$ is increasing. So $f''(x)$ is _____ for $a < x < b$. (Portions of u-shape)



We say that the graph of $f(x)$ is _____ for $a < x < b$.

Case 2: For $a < x < b$, slope of the graph $f(x)$ is **decreasing** as x increases i.e. $f'(x)$ is decreasing. So $f''(x)$ is _____ for $a < x < b$. (Portions of n-shape)




We say that the graph of $f(x)$ is _____ for $a < x < b$.

Second derivative test for concavity

Let $f(x)$ be a function that has a second derivative in an interval.

The above gives us:

• If $f''(x) > 0$ for all x then its graph is _____



• If $f''(x) < 0$ for all x then its graph is _____




Note: The places where the graph of $f(x)$ changes its concavity are called _____

Example 2 Using the graph of the derivative of $f(x)$ below, determine the concavity of $f(x)$.

Concave up:

Concave down:

Inflection points:



Q1: Where can $f''(x)$ change signs ($f(x)$ change concavity)?

A1: At the points where (i) _____, or (ii) _____ is undefined (e.g., f' has a sharp corner).

Example 3 The position of an object moving on a straight line is given by $s(t) = 2t^3 + 3t^2 - 36t + 7$. Determine (a) where the graph of $s(t)$ is concave up, (b) where it is concave down, and (c) where there are any inflection points, if any. Give physical interpretations for each of (a), (b), and (c).

Example 4 Determine where the graph of $f(x) = x^{5/3}$ is concave up, where it is concave down, and where there are any inflection points, if any. Sketch the graph of $f(x)$.