



Phenyl Glycopyranoside Recognition in Water Using Stoddart's Cyclobis(paraquat-*p*-phenylene) Receptor


Sandra A. Staley[†] and Bradley D. Smith^{*}

Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556, USA

Key Words: Molecular recognition; Cyclophane; Phenyl glycopyranoside; Association constants; Anomeric center.


Abstract: The cyclobis(paraquat-*p*-phenylene) receptor associates with various phenoxy-substituted guest compounds in water, including phenyl glycopyranosides. The receptor was found to bind phenyl β -D-glucopyranoside 0.6 kcal/mol more strongly than phenyl α -D-glucopyranoside. In addition, there was a general trend of increased affinity for anionic guests compared to structurally related neutral compounds.

Stoddart and coworkers have shown that cyclobis(paraquat-*p*-phenylene), **1⁴⁺**, is capable of binding electron-rich aromatic residues in organic and aqueous solution.¹ In particular, Stoddart and Kaifer have observed high associations with electron-rich aromatic amino acids (*e.g.*, tyrosine, tryptophane),² and neurotransmitters (*e.g.*, catecholamines)³ in aqueous solution. Cyclophane **1⁴⁺** has a number of structural attributes that simplify the determination and interpretation of its binding ability:¹ (i) Association with an electron-rich aromatic residue produces a colored, charge-transfer complex. Binding constants are readily determined by monitoring the appearance of this complex using standard spectrometric methods. (ii) The internal cavity of cyclophane **1⁴⁺** has a shape (10.3 x 6.8 Å) and hydrophobicity that make it a selective inclusion receptor for aromatic residues. (iii) Cyclophane **1⁴⁺** has a rigid structure that hardly changes upon inclusion of an aromatic guest, thus differences in guest association constants can be attributed directly to differences in guest structures. (iv) The solubility of tetracation **1⁴⁺** is strongly dependent on the identity of its counter-ions. For example, the tetrachloride salt is water-soluble, whereas the tetrakis(hexafluorophosphate) salt is soluble in acetonitrile. This allows binding affinities to be determined in solvents with substantially different properties. A consideration of the above features suggests cyclophane **1⁴⁺** can be employed as a binding probe for studies in molecular recognition.



Our interest lies in saccharide recognition in aqueous solution.⁴ The past few years have seen an increasing number of reports of small synthetic receptors designed to bind sugars using non-covalent

interactions.⁵⁻⁷ Most have been hydrogen bonding systems operating in non-polar solvents. Of particular relevance to the work described here is glycochane **2**, described by Penadés and coworkers.⁶ Host **2** was found to bind *p*-nitrophenyl glycopyranosides in water using a combination of hydrophobic forces, aromatic donor-acceptor interactions, and hydrogen bonding. Association constants were less than 300 M⁻¹, with a general binding preference for the α -anomer. Here we report on the ability of **14+** (tetrachloride salt) to bind various phenoxy-substituted organic compounds in water, including a range of phenyl D-glycopyranosides. In short, we find that host **14+** binds phenyl β -D-glycopyranosides more strongly than phenyl α -D-glycopyranosides.



Association constants, K_{assn} , and molar absorptivities, ϵ_{obs} , were determined from spectrometric titration studies at wavelength, λ_{obs} , using the method described by Stoddart.^{3,8} The values reported in Table 1 are an average of at least five independent trials, each using freshly made solutions.

Table 1: Binding of Guests With Host **14+** in Aqueous Buffer, pH 7.4, at 295 K.^a

Entry	Guest	K_{assn} 10 ³ (M ⁻¹) ^b	ΔG_{assn} (kcal mol ⁻¹)	ϵ_{obs} 10 ² (M ⁻¹ cm ⁻¹) ^a	λ_{obs} (nm)	¹³ C δ for phenoxy C4 (ppm) ^c
1	phenyl β -D-glucoside	4.1 \pm 0.5	4.9	2.3 \pm 0.6	326	121.7
2	phenyl α -D-glucoside	1.4 \pm 0.3	4.3	2.2 \pm 0.3	327	121.8
3	phenyl β -D-galactoside	5.3 \pm 0.7	5.1	2.0 \pm 0.1	326	121.7
4	phenyl α -D-mannoside	1.5 \pm 0.1	4.3	2.2 \pm 0.2	327	121.9
5	phenyl N-acetyl- β -D-glucosaminide	1.0 \pm 0.1	4.1	1.8 \pm 0.1	327	122.0
6	phenyl N-acetyl- α -D-glucosaminide	0.2 \pm 0.04	3.3	1.8 \pm 0.3	327	122.2
7	phenyl β -D-glucuronide	12.1 \pm 0.5	5.6	2.6 \pm 0.1	325	-
8	phenoxyacetic acid, 3	4.9 \pm 0.2	5.0	2.1 \pm 0.1	327	-
9	3-phenoxypropionic acid, 4	12.5 \pm 0.1	5.6	2.0 \pm 0.1	327	-
10	4-methoxyphenylacetic acid, 5	7.0 \pm 0.4	5.2	2.1 \pm 0.1	328	-
11	phenoxyethylpenicillanic acid, 6	1.1 \pm 0.3	4.1	2.5 \pm 0.1	324	-
12	phenoxyacetamide, 7	1.2 \pm 0.1	4.2	2.1 \pm 0.1	325	121.1
13	3-phenoxy-1,2-propanediol, 8	2.8 \pm 0.1	4.7	2.1 \pm 0.1	328	120.5

^aSee reference 8 for further details. ^bAverages and standard deviations calculated from at least five independent runs. ^cIn DMSO-*d*₆

As shown in Table 1, host **14+** bound phenyl β -D-glycopyranoside 0.6 kcal/mol more strongly than the α -anomer (entries 1 and 2). The trend also held for phenyl β -D-galactopyranoside and phenyl α -D-mannopyranoside (entries 3 and 4), as well as the anomers of phenyl N-acetyl-glucosaminide (entries 5 and 6), although in the latter case both binding affinities were reduced. In general, 1:1 binding stoichiometries for the **14+**:phenyl glycopyranoside complexes were confirmed from Job plots.⁹ Strong evidence that the guest phenyl ring was binding inside the host cavity was gained from ¹H NMR titration experiments. Incremental addition of host **14+** to the phenyl glycopyranoside guests produced a strong upfield movement of the guest phenyl

resonances with concomitant signal broadening.¹⁰ After the addition of about five molar equivalents of **14+**, the guest phenyl signals had disappeared into the baseline. The signals for the hydrogens on the pyranoside ring, on the other hand, remained essentially unchanged. These observations are in general agreement with previous binding studies using host **14+**.^{2,3}

To better understand the factors that affect host / guest binding, association constants were determined for a series of control compounds (Table 1). The strong binding observed with phenyl glucuronide (entry 7) suggested that coulombic attractions between tetracationic host **14+**, and an anionic guest can increase K_{assn} . In agreement with this suggestion is the trend of increased binding for phenoxy-substituted carboxylic acids **3 - 5** (entries 8 - 10) compared to the neutral controls **7** and **8** (entries 12 and 13). The proximity and orientation of the anionic carboxylate relative to the phenoxy group appears to be important. In particular, if the guest structure rigidly fixes the two groups with a large separation, no increase in K_{assn} is observed (compare entries 11 and 12).

The most intriguing trend within Table 1 is an apparent 0.7 kcal/mol increase in binding energies for phenyl β -glycopyranosides compared to their α -anomers. Possible explanations for this binding diastereoselectivity include arguments based on sterics, aromatic donor-acceptor interactions, and/or solvation effects. The steric argument focuses on the difference in steric accessibilities between the α - and β -anomeric positions.¹¹ The β -anomer allows the phenoxy substituent to adopt an equatorial position on the pyranoside ring, whereas, the α -anomer forces the phenoxy group into the sterically more-crowded axial position which may hinder access to the phenoxy-binding host **14+**. One example where a steric effect is clearly operating is the case of the phenyl *N*-acetyl-glucosaminides which have reduced binding affinities compared to the corresponding anomers of phenyl glucopyranoside (compare entries 5 and 6, with entries 1 and 2 respectively).

An explanation based on aromatic donor-acceptor interactions focuses on the amount of electron density within each glycoside's phenoxy group. Stoddart has established that the highly electron-deficient host **14+** associates more strongly with electron-rich aromatic guests.¹⁻³ To see if such a trend exists for the phenyl glycopyranosides in Table 1, ¹³C NMR was used to determine the relative order of phenoxy group electron densities. We reasoned that the ¹³C chemical shift for the phenoxy *para* carbon was a measure of relative electron density; the more upfield the chemical shift the more electron-rich the phenyl ring. As indicated in Table 1, there was no significant difference in ¹³C chemical shifts between glycoside anomers (the spectra were acquired in DMSO so as to eliminate ambiguities due to aqueous solvent effects). Therefore, as judged by this NMR criterion, there was no difference in phenoxy group electron densities.

An argument based on solvation effects would require the change in guest solvation upon association with host **14+** to be different for each of the phenyl glycopyranoside anomers. Since inclusion of the phenoxy groups into the cavity of host **14+** is driven, in part, by solvophobic effects in aqueous media, the difference in binding constants for the two anomers may be due to differences in solvophobic binding energies. A test of this explanation is to examine host / guest binding in a solvent where the solvophobic effect is diminished. Thus the association between host **14+** (as its tetrakis(hexafluorophosphate) salt), and the two anomers of phenyl D-glucopyranoside were determined in acetonitrile. As expected, the associations were found to be substantially weaker (for phenyl β -D-glucopyranoside $K_{\text{assn}} = 543 \text{ M}^{-1}$; for phenyl α -D-glucopyranoside $K_{\text{assn}} = 173 \text{ M}^{-1}$), due to the loss of the solvophobic driving force; however, a 0.7 kcal/mol binding preference in favor of the β -anomer was maintained. This result is evidence against a solvation effect as the source of the difference in anomer binding energies.

In conclusion, host 14^+ associates quite strongly with phenyl glycopyranosides in aqueous solution. Association is driven by inclusion of the guest phenoxy group inside the cavity of the cyclophane host. Phenyl β -D-glucopyranoside was found to bind 0.6 kcal/mol more strongly than the α -anomer. At present, the experimental evidence does not provide an unambiguous explanation for this binding diastereoselectivity. Chemical intuition suggests that steric differences between the α - and β -anomers may be the controlling factor.^{11,12} The very similar amounts of electron density within the glycoside phenyl groups indicate that differences in aromatic donor-acceptor interactions are not likely to be influential. Evidence against a solvation effect is the observation that the aqueous binding diastereoselectivity is maintained in acetonitrile.

Acknowledgment: This work was supported by a grant from the NSF (CHE 93-11584). We thank Dr. P. Duggan, James Cook University, Australia, for helpful discussions.

Notes and References

†Current address: Medicinal Chemistry Division, Upjohn Company, Kalamazoo, Michigan.

1. Odell, B.; Reddington, M. V.; Slawin, A. M. Z.; Spencer, N.; Stoddart, J. F.; Williams, D. J. *Angew. Chem. Int. Engl.* **1988**, *27*, 1547-1550.
2. Goodnow, T. T.; Reddington, M. V.; Stoddart, J. F.; Kaifer, A. E. *J. Am. Chem. Soc.* **1991**, *113*, 4335-4337.
3. Bernado, A. R.; Stoddart, J. F.; Kaifer, A. E. *J. Am. Chem. Soc.* **1992**, *114*, 10624-10631.
4. Bien, J. T.; Shang, M.; Smith, B. D. *J. Org. Chem.* **1995**, *60*, 2147-2152 and references cited therein.
5. Recent reports on sugar recognition using non-covalent interactions include: (a) Cuntze, J.; Owens, L.; Alcázar, V.; Seiler, P.; Diederich, F. *Helv. Chim. Acta* **1995**, *78*, 367-389. (b) Bonar-Law, R. P.; Sanders, J. K. M. *J. Am. Chem. Soc.* **1995**, *117*, 259-271. (c) Huang, C.; Cabell, L. A.; Anslyn, E. V. *J. Am. Chem. Soc.* **1994**, *116*, 2778-2792. (d) Das, G.; Hamilton, A. D. *J. Am. Chem. Soc.* **1994**, *116*, 11139-11140. (e) Eliseev, A. V.; Schneider, H. -J. *J. Am. Chem. Soc.* **1994**, *116*, 6081-6088. (f) Poh, B.; Tan, C. M. *Tetrahedron* **1993**, *49*, 9581-9592. (g) Savage, P. B.; Gellman, S. H. *J. Am. Chem. Soc.* **1993**, *115*, 10448-10449. (h) Liu, R.; Still, W. C. *Tetrahedron Lett.* **1993**, *34*, 2573-2576. (i) Kobayashi, K.; Asakawa, Y.; Kato, Y.; Aoyama, Y. *J. Am. Chem. Soc.* **1992**, *114*, 10307-10313.
6. Coterón, J. M.; Vicent, C.; Bosso, C.; Penadés, S. *J. Am. Chem. Soc.* **1993**, *115*, 10066-10076.
7. Recent reports on sugar recognition using covalent interactions include: (a) James, T. D.; Samankumara Sandanayake, K. R. A.; Shinkai, S. *Nature* **1995**, *374*, 345-347 and references cited therein. (b) Czarnik, A. W. *Acc. Chem. Res.* **1994**, *27*, 302-308. (c) Westmark, P. R.; Smith, B. D. *J. Am. Chem. Soc.* **1994**, *116*, 9343-9344.
8. Briefly, a solution of 14^+ (1 mM in 300 mM potassium phosphate, pH 7) was spectrometrically titrated with aliquots of 10 - 15 mM guest also containing 1 mM 14^+ . Usually the titration covered a guest / host range of 0.1 to 5, and corrections were made for background absorption by host and guest. The binding constant, K_{assn} , and the molar absorptivity, ϵ_{obs} , were determined using the equation: $A = (H K_{\text{assn}} \epsilon_{\text{obs}} G) / (1 + K_{\text{assn}} G)$; where A = absorbance at λ_{obs} , H = concentration of host, K_{assn} = association constant, ϵ_{obs} = molar absorptivity at λ_{obs} , G = concentration of guest.
9. (a) Conners, K. A. *Binding Constants, The Measurement of Molecular Complex Stability*, Wiley: New York, 1987. (b) Gil, V. M. S.; Olivera, N. C. *J. Chem. Ed.* **1990**, *67*, 472-478. (c) Ingham, K. C. *Anal. Biochem.* **1975**, *68*, 660-663.
10. Upon addition of one molar equivalent of host 14^+ , $\delta\Delta$ for the phenyl ^1H NMR resonances were -0.80 ppm for phenyl β -D-glucopyranoside and -0.65 ppm for phenyl α -D-glucopyranoside.
11. (a) Simmonds, R. J. *Chemistry of Biomolecules: An Introduction*, Royal Society: Cambridge, 1992. (b) Juaristi, E.; Cuevas, G. *The Anomeric Effect*, CRC Press: Baton Rouge, 1995.
12. Lipton, M. A.; the following paper in this journal.

(Received in USA 28 August 1995; revised 6 November 1995; accepted 8 November 1995)