

Reverse mathematics and the equivalence of definitions for well and better quasi-orders

Peter Cholak

Department of Mathematics, University of Notre Dame, USA

Alberto Marcone

Dipartimento di Matematica e Informatica, Università di Udine, Italy

Reed Solomon

Department of Mathematics, University of Connecticut, USA

March 30, 2004

1 Introduction

In reverse mathematics, one formalizes theorems of ordinary mathematics in second order arithmetic and attempts to discover which set theoretic axioms are required to prove these theorems. Often, this project involves making choices between classically equivalent definitions for the relevant mathematical concepts. In this paper, we consider a number of equivalent definitions for the notions of well quasi-order and better quasi-order and examine how difficult it is to prove the equivalences of these definitions.

As usual in reverse mathematics, we work in the context of subsystems of second order arithmetic and take RCA_0 as our base system. RCA_0 is the subsystem formed by restricting the comprehension scheme in second order arithmetic to Δ_1^0 formulas and adding a formula induction scheme for Σ_1^0 formulas. For the purposes of this paper, we will be concerned with fairly weak extensions of RCA_0 (indeed strictly weaker than the subsystem ACA_0 which is formed by extending the comprehension scheme in RCA_0 to cover all arithmetic formulas) obtained by adjoining certain combinatorial principles to RCA_0 . Among these, the most widely used in reverse mathematics is Weak König's Lemma; the resulting theory WKL_0 is extensively documented in [11] and elsewhere.

We give three other combinatorial principles which we use in this paper. In these principles, we use k to denote not only a natural number but also the finite set $\{0, \dots, k-1\}$. For any set X and any $n \in \mathbb{N}$, we let $[X]^n$ denote the set of all subsets of X of size n . Similarly, $[X]^{<\omega}$ denotes the

Cholak's research was partially supported by NSF Grants DMS 99-88716 and DMS 02-45167. Marcone's research was partially supported by INdAM of Italy and he thanks the Math Department of the University of Notre Dame for its kind hospitality. Solomon's research was partially supported by an NSF Postdoctoral Fellowship.




Diagram 1: none of the missing arrows hold except possibly the arrow from $RCA_0 + CAC$ to $WKL_0 + CAC$.

set of all finite subsets of X and $Y \in [X]^\omega$ is an abbreviation for the statement that Y is an infinite subset of X . The Pigeonhole principle is the statement

$$\forall k \forall f : \mathbb{N} \rightarrow k \exists A \in [\mathbb{N}]^\omega \exists i < k \forall j \in A (f(j) = i)$$

and is denoted $RT_{<infty}^1$. (This notation comes from thinking of the principle as a version of Ramsey's Theorem for singletons and finitely many colors.) Ramsey's Theorem for pairs and two colors is the statement

$$\forall f : [\mathbb{N}]^2 \rightarrow 2 \exists A \in [\mathbb{N}]^\omega \exists i < 2 \forall j \in [A]^2 (f(j) = i)$$

and is denoted RT_2^2 . The chain-antichain principle is denoted CAC and says that every infinite partial order has either an infinite chain (that is, an infinite linearly ordered subset) or an infinite antichain.

It is known that none of these principles is provable in WKL_0 and that over RCA_0 , RT_2^2 implies CAC which implies $RT_{<infty}^1$. $RT_{<infty}^1$ is the only one of these principles which is true in REC , the ω -model of the computable sets, and therefore, the implication from CAC to $RT_{<infty}^1$ cannot be reversed. It is an open question in reverse mathematics whether the implication from RT_2^2 to CAC can be reversed, even over WKL_0 . Diagram 1 illustrates the relationship of these principles below the system $WKL_0 + CAC$. (For more information on reverse mathematics and Ramsey's Theorem, see Cholak, Jockusch and Slaman [1].)

In Section 2, we give some background on well quasi-orders including five classically equivalent definitions for this concept. We show exactly how these equivalences are related in each of the systems presented in Diagram 1 as well as in the ω model REC , with the exception of one potential implication in the system $RCA_0 + CAC$. (This potentially missing implication depends on the open question of determining whether $RCA_0 + CAC$ implies $WKL_0 + CAC$.) Our main results are summarized in Diagrams 2, 3 and 4 and in Table 2. The proofs of these results are given in Section 3. Many of our results show that some implication between these equivalent definitions is not provable in RCA_0 by producing an appropriate "computable counterexample" to the classical equivalence. In some cases, these counterexamples can be extended to show that the implication is not even provable in WKL_0 . However, they do not lead to a reversal to a system stronger than RCA_0 in the usual reverse mathematics style. Therefore, although we can describe the exact relationships between the definitions in most of the systems above, we fall short of the ideal goal of giving an exact classification of the proof-theoretic strength for most of these equivalences.

In Section 4, we take a break from studying the reverse mathematics of equivalent definitions for well quasi-orders to look at the difficulty of proving that the notion of well quasi-order is closed

under products and intersections. Of course, we can check the difficulty of proving the closure properties for each of our five definitions for a well quasi-order. It is known that for any of these definitions, RCA_0 is strong enough to prove that the closure under either the operation of intersection or product implies the closure under the other operation. For the strongest definition of well quasi-order, RCA_0 suffices to prove the closure under these operation, but for the other four definitions, we show that there is an ω -model of WKL_0 which is not closed under either of these operations. Furthermore, we can mix-and-match these definitions by asking how difficult it is to prove that if we start with a well quasi-order under one definition, then the intersection or product satisfies another one of the definitions. We classify exactly which combinations are provable in RCA_0 and which combinations are not provable in WKL_0 .

In Section 5, we return to studying the reverse mathematics of equivalent definitions by considering the notion of a better quasi-orders. We present some background on better quasi-orders and examine the relationship between the definition of better quasi-order in terms of blocks and the definition in terms of barriers. Because it is shown in [7] that the closure of better quasi-orders under products and intersections implies ACA_0 (which is a strictly stronger subsystem than any mentioned in Diagram 1), we leave the project of classifying the proof-theoretic complexity of these operations as an open problem.

For a general survey of the classical theory of well quasi-orders and better quasi-orders, including important examples and more equivalent definitions, the reader is referred to Milner [8] and Simpson [9]. For more background in reverse mathematics, the reader is referred to Simpson [11]. For more on wqo and bqo theory from the reverse mathematics point of view, the reader is referred to Marcone [7].

Before beginning the mathematical part of the paper, we fix some notation for sequences. Given any set A , RCA_0 is strong enough to form $A^{<\omega}$, the set of all finite sequences of elements of A . If $a_1, \dots, a_n \in A$, then we write the sequence with these elements as $\langle a_1, \dots, a_n \rangle$. If $s, t \in A^{<\omega}$, then $|s|$ denotes the length of s and $s * t$ denotes the concatenation of s and t . If $a \in A$, we sometimes write $s * a$ in place of $s * \langle a \rangle$. We write $s \sqsubseteq t$ to denote that s is an initial segment of t and $s \sqsubset t$ to denote that s is a proper initial segment of t . Finally, if $n < |s|$, then the restriction of s to n , written $s|_n$, is the initial segment of s of length n .

2 Well quasi-orders

We begin with the basic definitions associated with quasi-orders and well quasi-orders.

Definition 2.1. (RCA_0) A *quasi-order* is a pair (Q, \preceq) such that Q is a set and \preceq is a transitive reflexive relation on Q .

Partial orders are the simplest examples of quasi-orders as a partial order is a quasi-order which also satisfies antisymmetry. We can transform a quasi-order Q into a partial order using the equivalence relation defined by $n \sim m$ if and only if $n \preceq m$ and $m \preceq n$. The quotient structure Q / \sim is a partial order which can be formed using Δ_1^0 comprehension in RCA_0 . Much of the standard terminology and notation for partial orders is used in the context of quasi-orders. For example, we write $i \perp j$ to indicate that i and j are incomparable under \preceq and we write $x < y$ if $x \preceq y$ and $y \not\preceq x$. Furthermore, we sometimes use $\preceq_{\mathbb{N}}$ to denote the order relation given by the symbol \preceq in

the language of second order arithmetic. This notation helps to emphasize when we are comparing elements of a quasi-order via the quasi-order relation and when we are comparing them via the underlying structure of arithmetic. We use this notation when the distinction between these orders is not immediately clear from the context.

$A \subseteq Q$ is an *antichain* if for all $i \neq j \in A$, $i \perp j$. An *infinite ascending chain* in Q is a function $f : \mathbb{N} \rightarrow Q$ such that for all $i <_{\mathbb{N}} j$, $f(i) < f(j)$. Similarly, a function f is an *infinite descending chain* in Q if for all $i <_{\mathbb{N}} j$, $f(j) < f(i)$. Any antichain in Q gives rise to a corresponding antichain in Q / \sim and vice versa. Infinite ascending chains and infinite descending chains in Q and Q / \sim are similarly related. Therefore, we frequently work with partial orders rather than quasi-orders.

A quasi-order (Q, \preceq) is called *linear* if for all $i, j \in Q$, either $i \preceq j$ or $j \preceq i$. For a linear quasi-order Q , Q / \sim is a linear order. If \preceq is a quasi-order on Q and \preceq_L is a linear quasi-order on Q , then we say \preceq_L is a *linear extension* of \preceq if for all $i, j \in Q$, $i \preceq j$ implies $i \preceq_L j$ and $i \sim_L j$ implies $i \sim j$. Notice that these conditions imply that $i \sim j$ if and only if $i \sim_L j$. Therefore, the linear extensions of a quasi-order Q correspond exactly to the linear extensions of the partial order Q / \sim . A linear quasi-order \preceq_L on Q is said to be *well ordered* if the induced linear order on Q / \sim_L is a well order. (In RCA_0 , the definition of a well order stating that each nonempty subset has a least element and the definition stating that there are no infinite descending chains are equivalent. We will use the definition in terms of no infinite descending chains.)

Definition 2.2. (RCA_0) A quasi-order (Q, \preceq) is a *well quasi-order* (or *wqo*) if for every function $f : \mathbb{N} \rightarrow Q$ there exist $m <_{\mathbb{N}} n$ such that $f(m) \preceq f(n)$.

As we shall see, there are many classically equivalent definitions for a well quasi-order, but the definition above is frequently used as the “official” definition for a well quasi-order in reverse mathematics. The simplest examples of well quasi-orders are well founded partial orders which have only finite antichains. For example, the m -fold cartesian product \mathbb{N}^m ordered by

$$\langle a_0, \dots, a_{m-1} \rangle \leq_{\mathbb{N}^m} \langle b_0, \dots, b_{m-1} \rangle \Leftrightarrow \forall i < m (a_i \leq b_i)$$

is a well quasi-order.

One natural question to ask about well quasi-orders is how difficult it is to prove that a given wqo is in fact well quasi-ordered. Simpson [10] showed that (given a suitable set of notations for the ordinals below ϵ_0) the system RCA_0 can prove that for all $m \in \mathbb{N}$, the statement that \mathbb{N}^m (ordered as above) is a wqo is equivalent to the statement that ω^m is a well order. It follows that for each standard m , RCA_0 can prove that \mathbb{N}^m is a well quasi-order, but that RCA_0 cannot prove the uniform statement that for all m , \mathbb{N}^m is a wqo.

Rather than pursue the task of describing the quasi-orders which are well quasi-orders in RCA_0 , we will look at various classically equivalent definitions for the notion of a well quasi-order and examine how difficult it is to prove their equivalence. An important notion for this direction of study is the finite basis property.

Definition 2.3. (RCA_0) A quasi-order \preceq on Q has the *finite basis property* if for every $X \subseteq Q$ there exists a finite $F \subseteq X$ such that $\forall x \in X \exists y \in F (y \preceq x)$.

Theorem 2.4 lists several classically equivalent definitions for a well quasi-order. Each of these equivalences can be proved using either Ramsey’s Theorem for pairs or Weak König’s Lemma, and the proofs can be formalized in $\text{WKL}_0 + \text{RT}_2^2$.

\rightarrow	wqo	wqo(set)	wqo(anti)	wqo(ext)
wqo	$\text{RCA}_0 \vdash$	$\rightarrow \text{RT}_{<\infty}^1$ $\text{WKL}_0 \not\vdash$ (*)	$\text{RCA}_0 \vdash$	
wqo(set)	$\text{RCA}_0 \vdash$	$\text{RCA}_0 \vdash$	$\text{RCA}_0 \vdash$	
wqo(anti)		$\rightarrow \text{RT}_{<\infty}^1$ $\text{WKL}_0 \not\vdash$	$\text{RCA}_0 \vdash$	
wqo(ext)				$\text{RCA}_0 \vdash$

Table 1: Previously known results

Theorem 2.4. *For any quasi-order (Q, \preceq) , the following are equivalent.*

1. (Q, \preceq) is a well quasi-order.
2. For all $f : \mathbb{N} \rightarrow Q$ there is an infinite set A such that for all $i, j \in A$, $i < j \rightarrow f(i) \preceq f(j)$.
3. (Q, \preceq) has no infinite descending chains and no infinite antichains.
4. Every linear extension of (Q, \preceq) is well ordered.
5. (Q, \preceq) has the finite basis property.

Our goal is to examine how difficult it is to prove these equivalences from the point of view of reverse mathematics. We use wqo to denote our official definition of well quasi-order in RCA_0 . We let wqo(set) denote the condition in 2, wqo(anti) denote the condition in 3, wqo(ext) denote the condition in 4, and wqo(fbp) denote the condition in 5. That is, we say Q satisfies wqo(ext) to mean that every linear extension of Q is well ordered, and we write $\text{wqo(ext)} \rightarrow \text{wqo(anti)}$ to denote the statement: for all quasi-orders Q , if every linear extension of Q is well ordered, then Q has no infinite descending chains and no infinite antichains.

A number of facts about the relationships between these notions are already known. For example, it follows trivially from the definitions that $\text{RCA}_0 \vdash \text{wqo(set)} \rightarrow \text{wqo}$. The following theorem summarizes the results of Lemmas 4.2, 4.4 and 4.8 in Marcone [7] and Lemma 3.2 in Simpson [10]. (Lemma 3.2 in [10] is stated in terms of partial orders rather than quasi-orders, but the proof translates quite easily.)

Theorem 2.5 (Marcone, Simpson). 1. $\text{RCA}_0 \vdash \text{wqo} \rightarrow \text{wqo(anti)}$.

2. $\text{RCA}_0 \vdash (\text{wqo} \rightarrow \text{wqo(set)}) \rightarrow \text{RT}_{<\infty}^1$.
3. $\text{RCA}_0 \vdash \text{wqo} \leftrightarrow \text{wqo(fbp)}$.

Because $\text{WKL}_0 \not\vdash \text{RT}_{<\infty}^1$, statement 2 in Theorem 2.5 implies that $\text{WKL}_0 \not\vdash \text{wqo} \rightarrow \text{wqo(set)}$. We can draw a number of other simple consequences from these results. For example, since $\text{RCA}_0 \vdash \text{wqo(set)} \rightarrow \text{wqo}$ and $\text{RCA}_0 \vdash \text{wqo} \rightarrow \text{wqo(anti)}$, we have $\text{RCA}_0 \vdash \text{wqo(set)} \rightarrow \text{wqo(anti)}$. For a second example, fix any model \mathcal{M} of RCA_0 and $\text{wqo(anti)} \rightarrow \text{wqo(set)}$. Since $\text{RCA}_0 \vdash$

\rightarrow	wqo	wqo(set)	wqo(anti)	wqo(ext)
wqo	$\text{RCA}_0 \vdash$	$\rightarrow \text{RT}_{<\infty}^1$ $\text{WKL}_0 \not\vdash$ $\text{CAC} \vdash$ (3.4) $\text{REC} \not\equiv$ (3.7)	$\text{RCA}_0 \vdash$	$\text{RCA}_0 \vdash$ (3.12)
wqo(set)	$\text{RCA}_0 \vdash$	$\text{RCA}_0 \vdash$	$\text{RCA}_0 \vdash$	$\text{RCA}_0 \vdash$ (3.13)
wqo(anti)	$\text{REC} \not\equiv$ (3.9) $\text{CAC} \vdash$ (3.5) $\text{WKL}_0 \not\vdash$ (3.11)	$\leftrightarrow \text{CAC}$ (3.3) $\text{WKL}_0 \not\vdash$ (3.6)	$\text{RCA}_0 \vdash$	$\text{REC} \not\equiv$ (3.10) $\text{CAC} \vdash$ (3.5) $\text{WKL}_0 \not\vdash$ (3.19)
wqo(ext)	$\text{REC} \not\equiv$ (3.21) $\text{WKL}_0 \vdash$ (3.17)	$\rightarrow \text{RT}_{<\infty}^1$ (3.20) $\text{WKL}_0 \not\vdash$ (3.20) $\text{REC} \not\equiv$ (3.24) $\text{WKL}_0 + \text{CAC} \vdash$ (3.4+3.17)	$\text{REC} \not\equiv$ (3.23) $\text{WKL}_0 \vdash$ (3.18)	$\text{RCA}_0 \vdash$

Table 2: Equivalent definitions for well quasi-orders

$\text{wqo} \rightarrow \text{wqo(anti)}$, we have that $\mathcal{M} \models \text{wqo} \rightarrow \text{wqo(set)}$ and hence $\mathcal{M} \models \text{RT}_{<\infty}^1$. Therefore, $\text{RCA}_0 \vdash (\text{wqo(anti)} \rightarrow \text{wqo(set)}) \rightarrow \text{RT}_{<\infty}^1$. We summarize the results in Theorem 2.5 and their simple consequences in Table 1. To read this table, consider the entries in square marked (*). The $\rightarrow \text{RT}_{<\infty}^1$ indicates that (over RCA_0) $\text{wqo} \rightarrow \text{wqo(set)}$ implies $\text{RT}_{<\infty}^1$ and the $\text{WKL}_0 \not\vdash$ indicates that $\text{WKL}_0 \not\vdash \text{wqo} \rightarrow \text{wqo(set)}$. In Table 1, we did not put a row or column for the condition wqo(fbp) because by statement 3 in Theorem 2.5, the row or column for wqo(fbp) would be equivalent to the corresponding row or column for wqo .

Notice that while statement 2 in Theorem 2.5 implies that $\text{WKL}_0 \not\vdash \text{wqo} \rightarrow \text{wqo(set)}$, it does not rule out the possibility that this implication is true in all ω -models of RCA_0 . In particular, it leaves open the question of whether REC , the ω -model consisting of the computable sets, is a model for this implication.

In Section 3, we improve the results in Theorem 2.5 to give all the results contained in Table 2. This table is read exactly as Table 1 and the information which is not contained in Table 1 will be proved in Section 3. As a reference to the reader, for all the results in Table 2 which do not appear in Table 1, we indicate in parentheses the number of the theorem, lemma or corollary in which we prove the particular statement.




Diagram 2: the exact implications in RCA_0 , in REC and in $\text{RCA}_0 + \text{RT}_{<\infty}^1$.




Diagram 3: the exact implications in WKL_0 .

The reverse mathematics content of Table 2 can be more easily visualized and summarized as follows. The system $\text{WKL}_0 + \text{CAC}$ is strong enough to prove the equivalence between all of the definitions for well quasi-order considered in this paper. Over RCA_0 , the only implications between the various definitions are those shown in Diagram 2 plus the fact that $\text{RCA}_0 \vdash \text{wqo} \leftrightarrow \text{wqo}(\text{fbp})$. Because we give computable counterexamples to implications which do not hold and because REC is a model for $\text{RCA}_0 + \text{RT}_{<\infty}^1$, Diagram 2 also illustrates which implications hold in the ω -model REC and in the subsystem $\text{RCA}_0 + \text{RT}_{<\infty}^1$. Over WKL_0 , the implications which hold are exactly those shown in Diagram 3.

Over $\text{RCA}_0 + \text{CAC}$, the known implications are those shown in Diagram 4. Unlike Diagrams 2 and 3, it is possible that the lone missing implication (from $\text{wqo}(\text{ext})$ to the other three conditions) in Diagram 4 does hold. However, notice that since $\text{WKL}_0 \vdash \text{wqo}(\text{ext}) \rightarrow \text{wqo}$, any proof that $\text{RCA}_0 + \text{CAC} \not\vdash \text{wqo}(\text{ext}) \rightarrow \text{wqo}$ would yield a proof that $\text{RCA}_0 + \text{CAC} \not\vdash \text{WKL}_0$ and hence $\text{RCA}_0 + \text{RT}_2^2 \not\vdash \text{WKL}_0$. Such a proof would answer one of the major open questions in reverse mathematics.

Finally, we note that there are many other combinatorial principles which sit below $\text{WKL}_0 + \text{RT}_2^2$




Diagram 4: the implications in $\text{RCA}_0 + \text{CAC}$. It remains open whether the missing arrows hold.

in reverse mathematics and which could be compared to the implications examined in this paper. For example, Hirschfeldt and Shore [5] have examined variations on the statement that every infinite linear order has either an infinite ascending sequence or an infinite descending sequence (denote ASDS). It is easy to show that RCA_0 suffices to prove that $\text{wqo} \rightarrow \text{wqo}(\text{set})$ implies ASDS. To see this claim, we assume that $\text{wqo} \rightarrow \text{wqo}(\text{set})$ and we work in RCA_0 . Fix a linear order L and assume that L has no infinite descending sequences. Then L obviously satisfies wqo . Let f be a one-to-one enumeration of L and let A be the infinite set obtained from the statement of $\text{wqo}(\text{set})$. For any $i, j \in A$, if $i <_{\mathbb{N}} j$, then $f(i) <_L f(j)$ and so A gives us the ascending chain required for ASDS. It follows from the work in [5] that $\text{COH} \not\vdash \text{wqo} \rightarrow \text{wqo}(\text{set})$ where COH is the combinatorial statement that every sequence of sets $X_i, i \in \mathbb{N}$, has an infinite cohesive set C . That is, for all i , either C is contained in X_i except for finitely many elements or X_i is contained in C for finitely many elements. We will not explore these connections any further in this paper, but we wanted to alert the reader that the situation concerning combinatorial principles below RT_2^2 is diverse and complicated.

3 Proofs for well quasi-order results

We begin this section by showing that we can replace the use of RT_2^2 by CAC in the proofs of several of our implications. Since it is not known whether CAC implies RT_2^2 , it is possible that this replacement is not actually an improvement on the known results. First, we show two consequences of $\text{RCA}_0 + \text{CAC}$. (We assume that Lemma 3.1 is already widely known and Lemma 3.2 is due to Hirschfeldt and Shore [5]. We include proofs only for the sake of completeness.)

Lemma 3.1. $\text{RCA}_0 + \text{CAC} \vdash \text{RT}_{<\infty}^1$.

Proof. Fix $k \in \mathbb{N}$ and let $f : \mathbb{N} \rightarrow k$. We need to find an infinite set on which f is constant. Define a partial order $<_f$ on \mathbb{N} by $n <_f m$ if and only if $n <_{\mathbb{N}} m$ and $f(n) = f(m)$. It is clear that for any n and m , if $f(n) = f(m)$, then either $n \leq_f m$ or $m \leq_f n$. Therefore, any antichain in our partial order has size at most k . Applying CAC , we obtain an infinite chain in this partial order. However, f must be constant on any chain so we have our desired infinite set. \square

Lemma 3.2. $\text{RCA}_0 + \text{CAC}$ suffices to prove that for every partial order P and for every infinite chain $C \subset P$, either C contains an infinite ascending sequence or C contains an infinite descending sequence.

Proof. Fix a partial order P and an infinite chain $C \subseteq P$. We define a new partial order \leq' on C by setting $n \leq' m$ if and only if $n \leq_P m$ and $n \leq_{\mathbb{N}} m$. Applying CAC , there is either an infinite chain or an infinite antichain in \leq' . It is not hard to check that RCA_0 suffices to prove that an infinite chain in \leq' corresponds to an infinite increasing sequence in \leq_P and an infinite antichain in \leq' corresponds to an infinite descending sequence in \leq_P . (To see these correspondences, just list the chain or antichain in $\leq_{\mathbb{N}}$ -increasing order.) \square

Lemma 3.3. (RCA_0) The following are equivalent:

1. CAC

2. $wqo(anti) \rightarrow wqo(set)$

Proof. We first show that CAC is strong enough to prove $wqo(anti) \rightarrow wqo(set)$. Fix any quasi-order (Q, \prec) which has no infinite descending chains or infinite antichains and fix any function $f : \mathbb{N} \rightarrow Q$. Consider the equivalence relation \sim on Q defined by $x \sim y$ if and only if $x \preceq y$ and $y \preceq x$. If the range of f intersects only finitely many equivalence classes under \sim , then we can apply $RT_{<\infty}^1$ to obtain an infinite set A such that f restricted to A maps into a single equivalence class of \sim . For any $i, j \in A$, we have $f(i) \preceq f(j)$, so A satisfies $wqo(set)$ for the map f . Therefore, by Lemma 3.1, we can assume that f maps into infinitely many equivalence classes under \sim .

We now look at the partial order Q / \sim and we view f as mapping into this partial order. By our assumption the range of f when viewed as mapping into Q / \sim is infinite. We abuse notation slightly and write $x \preceq y$ to denote the order relation in Q / \sim . We can assume without loss of generality that f is one-to-one, since f can be replaced by a one-to-one function h with the same range and such that for all n and k , if $h(n) = k$, then there is an $m \geq n$ such that $f(m) = k$. We need to find an infinite set A such that for all $i, j \in A$, if $i < j$, then $f(i) \preceq f(j)$. We define an auxiliary partial order P with domain \mathbb{N} such that $n <_P m$ if and only if $f(n) \prec f(m)$. Notice that any antichain in \preceq_P corresponds to an antichain in Q / \sim , which in turn corresponds to an antichain in Q . Since there are no infinite antichains in Q , there cannot be an infinite antichain in P .

Since P is an infinite partial order, we can apply CAC to it. Furthermore, since P has no infinite antichains, we must obtain an infinite chain when we do this. However, by Lemma 3.2, CAC can prove that this infinite chain contains either an infinite descending sequence or an infinite ascending sequence. Since an infinite descending sequence in P would yield an infinite descending sequence in Q , there cannot be such a sequence. Therefore, we obtain an infinite ascending sequence $b_0 <_P b_1 <_P \dots$ in P . We define an infinite set A by enumerating it in increasing order $a_0 <_{\mathbb{N}} a_1 <_{\mathbb{N}} \dots$ as follows. Set $a_0 = b_0$. If $a_i = b_j$, then let $a_{i+1} = b_k$ where k is the least index such that $k > j$ and $b_k >_{\mathbb{N}} b_j$. Notice that $a_i <_{\mathbb{N}} a_j$ if and only if $i <_{\mathbb{N}} j$ and that $a_i <_{\mathbb{N}} a_j$ implies $a_i <_P a_j$ which implies $f(a_i) \prec f(a_j)$. Therefore, A is the desired set for the function f in the statement of $wqo(set)$.

We next show that $wqo(anti) \rightarrow wqo(set)$ implies CAC. Fix an infinite partial order P . We need to show that P has either an infinite chain or an infinite antichain. Assume that P does not have an infinite descending chain or an infinite antichain. Then P satisfies $wqo(anti)$ and by $wqo(anti) \rightarrow wqo(set)$ we have that P satisfies $wqo(set)$. Let f be a one-to-one map from \mathbb{N} onto P . The infinite set A from the statement of $wqo(set)$ gives the desired infinite chain. \square

Corollary 3.4. $RCA_0 + CAC \vdash wqo \rightarrow wqo(set)$.

Proof. This corollary follows from Lemma 3.3 since $RCA_0 \vdash wqo \rightarrow wqo(anti)$ (from Theorem 2.5). \square

Corollary 3.5. $RCA_0 + CAC$ suffices to prove $wqo(anti) \rightarrow wqo$ and $wqo(anti) \rightarrow wqo(ext)$.

Proof. These two statements follow from Lemma 3.3 plus $RCA_0 \vdash wqo(set) \rightarrow wqo$ (trivial from the definitions) and $RCA_0 \vdash wqo(set) \rightarrow wqo(ext)$ (to be proved in Lemma 3.12). \square

Corollary 3.6. $WKL_0 \not\vdash wqo(anti) \rightarrow wqo(set)$

Proof. This corollary follows from Lemma 3.3 because $\text{WKL}_0 \not\vdash \text{CAC}$. \square

Next we show that several of the implications between the classical definitions for a well quasi-order are not true in REC. Recall that REC is the ω model consisting of the computable sets and that it is the minimal ω model of RCA_0 .

Lemma 3.7. $\text{REC} \not\vdash \text{wqo} \rightarrow \text{wqo}(\text{set})$.

Proof. Let L be a computable linear order which has order type $\omega + \omega^*$ but which does not have any infinite computable ascending or descending chains. (The existence of such orders was proved independently by Denisov and Tennenbaum and a proof can be found in [2].) The fact that L has no infinite descending chains means that it satisfies wqo in REC. However, the fact that L has no infinite computable ascending chains means that L does not satisfy wqo(set) in REC. \square

Our use of an order which is not really a wqo but only looks like a wqo in REC is necessary in the proof of Lemma 3.7. To see why this is so, consider the following result from Herrmann [4]. We have stated the result in a slightly different form than in [4], but an examination of Herrmann's proof shows that it also yields this statement.

Theorem 3.8 (Herrmann). *Every computable partial order has either a computable infinite ascending sequence or a Δ_2^0 infinite descending sequence or a Π_2^0 infinite antichain.*

Let (P, \leq_P) be a computable partial order which is a real wqo so that P has neither an infinite antichain nor an infinite descending sequence. We claim that P satisfies wqo(set) in REC. Fix any map $f : \mathbb{N} \rightarrow P$. Since $\text{REC} \models \text{RT}_{<\infty}^1$, we can assume as in the proof of Lemma 3.3 that the range of f is infinite and that f is one-to-one. We define an auxiliary partial order \leq_f with domain \mathbb{N} such that $n <_f m$ if and only if $f(n) <_P f(m)$. By Herrmann's result, \leq_f has a computable infinite increasing sequence $n_0 <_f n_1 <_f \dots$. As in the proof of Lemma 3.3, this infinite increasing sequence yields an infinite set A which satisfies wqo(set) for the map f .

Theorem 3.9. $\text{REC} \not\vdash \text{wqo}(\text{anti}) \rightarrow \text{wqo}$ and hence $\text{RCA}_0 \not\vdash \text{wqo}(\text{anti}) \rightarrow \text{wqo}$.

Proof. We present the proof of this theorem in considerable detail and trust the reader to fill in similar details on subsequent proofs. We build a computable partial order (P, \leq_P) in stages. At stage s , we specify a finite set P_s and a partial order on P_s . Once the partial order on P_s is defined at stage s , we cannot change it later in the construction. At the end of the construction, the domain of P (which is $\bigcup P_s$) will be equal to \mathbb{N} , and hence (P, \leq_P) will be a computable partial order. As previously mentioned, we let $n \perp m$ denote that n and m are incomparable in P . P will have the stronger property that $n \perp m$ if and only if n and m are incompatible: $\neg \exists z (z \leq_P n \wedge z \leq_P m)$.

To make sure that (P, \leq_P) has no computable infinite descending chains and no computable infinite antichains, we meet the following requirements.

$$\begin{aligned} \mathcal{R}_e : \varphi_e \text{ total} &\rightarrow \exists n <_{\mathbb{N}} m (\varphi_e(n) \leq_P \varphi_e(m) \vee \varphi_e(n) \perp \varphi_e(m)) \\ \mathcal{S}_e : \varphi_e \text{ total} &\rightarrow \exists n \neq m (\varphi_e(n) \leq_P \varphi_e(m)). \end{aligned}$$

The priority order on these requirements is $\mathcal{R}_0, \mathcal{S}_0, \mathcal{R}_1, \mathcal{S}_1, \dots$. To make sure that (P, \leq_P) is not a wqo in REC, we need to construct a computable function f such that for all $i < j$, either $f(i) \perp$

$f(j)$ or $f(j) <_P f(i)$. Our function f is the simplest possible: $f(n) = n$. That is, if we just list the element of P in $\leq_{\mathbb{N}}$ increasing order, we get a computable witness that (P, \leq_P) is not a wqo.

The basic module for \mathcal{R}_e is as follows.

1. Put down two incomparable points q_0^e and q_1^e . Force all lower priority requirements to place new points in P below q_0^e .
2. Wait for stage s and a number x such that $\varphi_{e,s}(x) = y$ for some y with $y \leq_P q_0^e$.
3. For all stages $t > s$, force lower priority requirements to place new points in P below q_1^e instead of below q_0^e . That is, \mathcal{R}_e forbids putting any points below q_0^e after the current stage.

This module will succeed, at least in isolation from the other strategies. If there is never convergence as in 2, then φ_e enumerates only points not below q_0^e . Since there are only finitely many such points, φ_e cannot be an infinite descending chain. If there is convergence as in 2, then φ_e enumerates at least one point below q_0^e . By 3, the number of points below q_0^e is finite, so φ_e cannot be an infinite descending chain.

The basic module for \mathcal{S}_e is as follows.

1. Put down a single point q^e . Force all lower priority requirements to place new points in P below q^e .
2. Wait for stage s and a number x such that $\varphi_{e,s}(x) = y$ for some y with $y \leq_P q^e$.
3. Pick a point $z \leq_P y$ which is currently minimal in P . For all stages $t > s$, force lower priority requirements to place new points in P below z . That is, \mathcal{S}_e requires lower priority requirements to work in the cone below z after the current stage. (The only reason for choosing the point z is to make the order relation easier to describe in the general construction.)

Again, we explain why this module succeeds in isolation from the other strategies. If there is no convergence as in 2, then φ_e enumerates only points which are not below q^e . Since there are only finitely many such points, φ_e cannot enumerate an infinite antichain. If there is convergence as in 2, then φ_e enumerates at least one point y above z . Any element incomparable with this point y cannot be below z . In this case, by 3 there are only finitely many points not below z , so φ_e cannot enumerate an infinite antichain.

There is an obvious source of conflict between these strategies, but it can easily be resolved with a finite injury construction. We present the formal construction using a tree of strategies, although such a tree is not necessary. However, it will be useful for the proof of Corollary 3.10 below to have the notation given by this tree. The tree of strategies T is the full binary branching tree on the symbols w and d with the d outcome to the left of the w outcome. More formally, we define T inductively. The empty string λ is in T and is an \mathcal{R}_0 node. If α is an \mathcal{R}_e node in T , then $\alpha * d$ and $\alpha * w$ are both \mathcal{S}_e nodes in T with $\alpha * d$ to the left of $\alpha * w$. If α is an \mathcal{S}_e node in T , then both $\alpha * d$ and $\alpha * w$ are \mathcal{R}_{e+1} nodes in T with $\alpha * d$ to the left of $\alpha * w$. For $\alpha \neq \beta \in T$, we say that α has higher priority than β if $\alpha \sqsubset \beta$ or $\alpha(i) = d$ and $\beta(i) = w$ where i is the least value for which $\alpha(i) \neq \beta(i)$. If $\alpha \in T$ and $\alpha \neq \lambda$, then α^- denotes the node in T formed by removing the last element of α .

We can now present the formal construction. At stage 0, the current true path δ_0 in T consists only of the node λ . The λ strategy defines $q_0^\lambda = 0$ and $q_1^\lambda = 1$, adds these two points to P and makes them incomparable in \leq_P . Also, λ defines $u_\lambda = q_0^\lambda$. The parameter u_λ is used to tell lower priority requirements which cones they have to place their points in. We end stage 0 and say that the strategy λ was eligible to act at stage 0.

At stage $t > 0$, we define the current true path δ_t in T such that $|\delta_t| = t$. We define δ_t by induction using a sequence of approximations $\delta_{t,s}$, for $s \leq t$, such that $|\delta_{t,s}| = s$ and $\delta_{t,s} \sqsubset \delta_{t,s+1}$. We set $\delta_t = \delta_{t,t}$. We begin with $\delta_{t,0} = \lambda$. To proceed by induction, assume that $\delta_{t,s} = \alpha$. We describe the action taken by α and how to define $\delta_{t,s+1}$ (if $s < t$).

If α is an \mathcal{R}_e strategy, then proceed as follows. The parameter u_{α^-} denotes the element of P below which α must place its new points, if this is the first stage at which α is eligible to act. Notice that step 1 will never apply to λ since λ was eligible to act at stage 0. Therefore, we do not need to worry that λ^- is undefined.

1. If α has been on the current true path before, then skip to step 2. Otherwise, let q_0^α and q_1^α be the $\leq_{\mathbb{N}}$ -least numbers not in P yet. Make them incomparable with each other in \leq_P , below u_{α^-} , and incomparable with all points which are not $\geq_P u_{\alpha^-}$. Set $u_\alpha = q_0^\alpha$. If $s = t$, then end the stage, and otherwise, go to step 2.
2. Check if there is an $x \leq t$ such that $\varphi_{e,t}(x)$ converges to a node $\leq_P q_0^\alpha$. If not, then leave $u_\alpha = q_0^\alpha$ and let $\delta_{t,s+1} = \alpha * w$. If there is such an x , then redefine $u_\alpha = q_1^\alpha$ and let $\delta_{t,s+1} = \alpha * d$. (Of course, once we have seen this convergence and redefined u_α , then we do not redefine it again at future stages.)

If α is an \mathcal{S}_e node, then proceed as follows.

1. If α has been on the current true path before, then skip to step 2. Otherwise, let q^α be the $\leq_{\mathbb{N}}$ -least number not in P yet. Place q^α below u_{α^-} and incomparable with all points which are not $\geq_P u_{\alpha^-}$. Let $u_\alpha = q^\alpha$. If $s = t$, then end the stage, and otherwise, go to step 2.
2. Check if there is an $x \leq t$ such that $\varphi_{e,t}(x)$ converges to an element $y \leq_P q^\alpha$. If not, then leave $u_\alpha = q^\alpha$ and let $\delta_{t,s+1} = \alpha * w$.
3. If there is such an x and $\alpha * d$ was on the current true path at stage $t - 1$ (that is, $\alpha * d \sqsubseteq \delta_{t-1}$), then let u_α be the same as it was at stage $t - 1$ and let $\delta_{t,s+1} = \alpha * d$. (This case corresponds to the situation in which we have already diagonalized and just want to continue the action described at stage $t - 1$.)
4. If there is such an x and $\alpha * d$ was not on the current true path at stage $t - 1$ (that is, $\alpha * d \not\sqsubseteq \delta_{t-1}$), then consider the least such x and let $\varphi_e(x) = y$. Let z be any point such that $z \leq_P y$ and there are currently no points in P below z . Redefine $u_\alpha = z$ and let $\delta_{t,s+1} = \alpha * d$.

This completes the description of the construction of P . We sketch the verification that the construction succeeds. First, we claim that if α is first eligible to act at stage t , then at stage t , there are no points below u_{α^-} . This is proved by induction on t and on the nodes α on δ_t . If α is on the current true path for the first time at stage t , then it is for one of three reasons: α^- is first on the

current true path at stage t , $\alpha = \alpha^- * d$ and α^- has always taken the $\alpha^- * w$ outcome before, or $|\alpha| = t$. It is straightforward to verify in each case that u_{α^-} has no nodes below it before α acts.

Second, the current true path can only move left between consecutive stages of the construction. Therefore, the initialization of lower priority strategies is automatically taken care of by moving to strategies which have never acted before. Furthermore, this property means that we can define the true path as the pointwise limit of the current true paths visited during the construction. If α is on the true path, then there is a stage s such that $\alpha \sqsubseteq \delta_t$ (that is, α is on the current true path) for every stage $t \geq s$. The module for a strategy α on the true path succeeds exactly as in the informal description of the modules working in isolation.

Third, notice that during stage t of the construction, we never place a new point above a point which is already in P . Since we place points in P in $\leq_{\mathbb{N}}$ increasing order, this means that the function $f(n) = n$ is a computable witness that P is not a wqo. \square

Corollary 3.10. $\text{REC} \not\models \text{wqo}(anti) \rightarrow \text{wqo}(ext)$ and $\text{RCA}_0 \not\models \text{wqo}(anti) \rightarrow \text{wqo}(ext)$.

Proof. Let \leq_P be the computable partial order from the proof of Theorem 3.9. We prove that there is a computable linear extension \leq_L of \leq_P and an infinite descending chain in \leq_L given by a computable function f . Before describing the linear extension, we note three properties of the construction of P . First, if $x <_P q_0^\alpha$ for some \mathcal{R}_e strategy α , then x is placed in the partial order by a strategy β such that $\alpha * w \sqsubseteq \beta$. Second, if $x <_P q_1^\alpha$ for some \mathcal{R}_e strategy α , then x is placed in the partial order by a strategy β such that $\alpha * d \sqsubseteq \beta$. Third, if $x <_P q^\alpha$ for some \mathcal{S}_e strategy α , then x is placed in the partial order by a strategy β such that $\alpha \sqsubset \beta$. These properties are all easily verified by induction on the stages of the construction.

We next assign each point x in P to a unique node in the tree of strategies from the proof of Theorem 3.9. If $x = q_0^\alpha$ for some \mathcal{R}_e strategy α , then x is assigned to $\alpha * w$. If $x = q_1^\alpha$ for some \mathcal{R}_e strategy α , then x is assigned to $\alpha * d$. If $x = q^\alpha$ for some \mathcal{S}_e strategy α , then x is assigned to $\alpha * w$. We let p_α denote the unique element (if any) of P assigned to α . Notice that every point in P is assigned to some α . The linear order \leq_L is defined by $p_\beta \leq_L p_\gamma$ if and only if $\gamma \sqsubseteq \beta$ or β is to the left of γ . Since \leq_L just copies the Kleene-Brouwer order on the tree of strategies, it is clearly a linear order on P .

We need to see that \leq_L is a linear extension of \leq_P . We show by induction on t that \leq_L is a linear extension of \leq_P on the points of P at the end of stage t . At the end of stage 0, P contains only the incomparable points q_0^λ and q_1^λ . Therefore, our induction holds at stage 0. We assume the induction holds at stage t and we show it holds at stage $t + 1$ by induction on the strategies in the current true stage. That is, we examine how points are added during stage $t + 1$ and show that \leq_L extends \leq_P after each addition. Suppose a strategy α adds points to P at stage $t + 1$ and that before α adds its points, \leq_L is an extension of \leq_P . We split into cases depending on what type of strategy α is and whether it is below a d or w outcome.

Consider the case when α is an \mathcal{R}_e strategy and $\alpha = \alpha^- * w$. Then, α^- is an \mathcal{S}_{e-1} strategy and in \leq_P , we add q_0^α and q_1^α as incomparable points immediately below q^{α^-} . Therefore, it suffices to show that q_0^α and q_1^α are $<_L q^{\alpha^-}$. In our notation of assigning elements of P to strategies, we have $q^{\alpha^-} = p_\alpha$, $q_0^\alpha = p_{\alpha * w}$ and $q_1^\alpha = p_{\alpha * d}$. Since $\alpha \sqsubset \alpha * w$ and $\alpha \sqsubset \alpha * d$, we have that $q_0^\alpha <_L q^{\alpha^-}$ and $q_1^\alpha <_L q^{\alpha^-}$ as required. A similar proof works in the cases when α is an \mathcal{S}_e strategy and either $\alpha = \alpha^- * w$ or $\alpha = \alpha^- * d$.

It remains to examine the case when α is an \mathcal{R}_e strategy and $\alpha = \alpha^- * d$. In this case, α adds the points q_0^α and q_1^α to P immediately below some point $z \leq_P q^{\alpha^-}$. Therefore, it suffices to show that q_0^α and q_1^α are $<_L z$. By the properties listed in the first paragraph of this proof, z is added to P by some strategy β with $\alpha^- \sqsubseteq \beta$. Since $\alpha = \alpha^- * d$ has never been eligible to act before this stage, we cannot have $\alpha \sqsubseteq \beta$. Therefore, either $\beta = \alpha^-$ or $\alpha^- * w \sqsubseteq \beta$. If $\beta = \alpha^-$, then $z = q^{\alpha^-}$ and so $z = p_{\alpha^- * w}$. Since $\alpha = \alpha^- * d$ is to the left of $\alpha^- * w$, we have that $\alpha * w$ and $\alpha * d$ are to the left of $\alpha^- * w$. Therefore, $q_0^\alpha = p_{\alpha * w}$ and $q_1^\alpha = p_{\alpha * d}$ are $<_L q^{\alpha^-} = p_{\alpha^- * w}$ as required. The last remaining case is when $\alpha^- * w \sqsubseteq \beta$. But, then α is to the left of β and a similar argument shows q_0^α and q_1^α are $<_L z$ as required.

We still need to define a computable function $f : \mathbb{N} \rightarrow P$ such that for all t , $f(t+1) <_L f(t)$. Let t be any stage of the construction. There is a last strategy α_t which is eligible to act at this stage and since $|\alpha_t| = t$, t is the first stage at which α_t is eligible to act. (Technically, α_t is equal to δ_t , but we use the notation α_t to distinguish the strategy from the current true path.) Therefore, α_t adds at least one point to P and defines u_{α_t} to be one of the new points it has added to P . Furthermore, in the notation introduced above, u_{α_t} is always equal to $p_{\alpha_t * w}$. We define $f(t) = u_{\alpha_t} = p_{\alpha_t * w}$.

It is clear that f is computable. To verify that $f(t+1) <_L f(t)$, we split into two cases. First, if $\alpha_{t+1}^- \neq \alpha_t$, then α_{t+1} is to the left of α_t in the tree of strategies. (This follows since the current true path in the construction only moves to the left.) Therefore, $\alpha_{t+1} * w$ is to the left of $\alpha_t * w$ and so $f(t+1) = p_{\alpha_{t+1} * w} <_L p_{\alpha_t * w} = f(t)$. Second, assume that $\alpha_{t+1}^- = \alpha_t$. If $\alpha_{t+1} = \alpha_t * w$, then $\alpha_t * w \sqsubset \alpha_{t+1} * w$ and so $f(t+1) = p_{\alpha_{t+1} * w} <_L p_{\alpha_t * w} = f(t)$. Otherwise, we have $\alpha_{t+1} = \alpha_t * d$. In this case, we first assume that α_t is an \mathcal{S}_e strategy. We know $f(t)$ is set to the value of u_{α_t} at the end of stage t . Thus, $f(t) = q^{\alpha_t}$. However, α_t was the last strategy to act at stage t and so no strategy has placed any points below q^{α_t} . Therefore, when α_t goes to the $\alpha_t * d$ outcome at stage $t+1$, the only choice for u_{α_t} at stage $t+1$ is again q^{α_t} . Therefore, α_{t+1} defines $f(t+1) = u_{\alpha_{t+1}}$ to be a point $<_P q^{\alpha_t} = f(t)$. Since \leq_L extends \leq_P , we have $f(t+1) <_L f(t)$. The last remaining case is when $\alpha_{t+1} = \alpha_t * d$ and α_t is an \mathcal{R}_e strategy. Then $f(t) = q_0^{\alpha_t}$ and α_{t+1} places its new points $<_P q_1^{\alpha_t}$. Again, because \leq_L extends \leq_P , we have $f(t+1) <_L q_1^{\alpha_t}$. However, we set $q_1^{\alpha_t} <_L q_0^{\alpha_t}$ since $q_1^{\alpha_t} = p_{\alpha * d}$, $q_0^{\alpha_t} = p_{\alpha * w}$ and $\alpha * d$ is to the left of $\alpha * w$. Therefore, $f(t+1) <_L f(t)$ as required. This completes the proof that f gives a computable infinite descending chain in \leq_L . \square

The proof of Theorem 3.9 can be strengthened to show that WKL_0 does not suffice to prove $\text{wqo}(\text{anti}) \rightarrow \text{wqo}$. The technique used to prove this stronger result is similar to the methods used in [3] and the reader is referred there for more applications of this method.

Theorem 3.11. $\text{WKL}_0 \not\vdash \text{wqo}(\text{anti}) \rightarrow \text{wqo}$.

Proof. To prove this theorem, we show that a particular model (or rather a particular type of model) of WKL_0 cannot be a model of the implication $\text{wqo}(\text{anti}) \rightarrow \text{wqo}$. Formally, we show that if X_0, X_1, \dots is a sequence of uniformly Δ_2^0 , uniformly low sets, then we can construct a computable partial order \leq_P which meets the requirements:

$$\begin{aligned} \mathcal{R}_{\langle e, i \rangle} : \varphi_e^{X_i} \text{ total} &\rightarrow \exists n <_{\mathbb{N}} m (\varphi_e^{X_i}(n) \leq_P \varphi_e^{X_i}(m) \vee \varphi_e^{X_i}(n) \perp \varphi_e^{X_i}(m)) \\ \mathcal{S}_{\langle e, i \rangle} : \varphi_e^{X_i} \text{ total} &\rightarrow \exists n \neq m (\varphi_e^{X_i}(n) \leq_P \varphi_e^{X_i}(m)). \end{aligned}$$

The motivation for meeting these requirements is that there is such a sequence of sets X_i for which the ω -model \mathcal{A} with second order part $\{Y \mid \exists i (Y \leq_T X_i)\}$ is a model of WKL_0 . Since \leq_P is a

computable partial order, it is an element of \mathcal{A} . Furthermore, the $\mathcal{R}_{\langle e, i \rangle}$ requirements show that \leq_P has no infinite descending sequences in \mathcal{A} and the $\mathcal{S}_{\langle e, i \rangle}$ requirements show that \leq_P has no infinite antichains in \mathcal{A} . Therefore, \leq_P satisfies wqo(anti) in \mathcal{A} . To make sure that \leq_P does not satisfy wqo in \mathcal{A} , we construct a computable function f such that for all $i < j$, either $f(i) \perp f(j)$ or $f(j) <_P f(i)$, so we even have a computable witness that \leq_P does not satisfy wqo. As above, our function is $f(n) = n$.

This argument is almost identical to the argument in Theorem 3.9 except that we may finitely often switch between the outcomes w and d rather than just moving from w to d and never returning to w . The argument is still finite injury, but it may take more than one (but still finitely many) attempts to successfully diagonalize. Although this argument can be given using a tree of strategies as in Theorem 3.9, it is probably easiest to formalize in a more straightforward manner. The main point of using the tree of strategies in the proof of Theorem 3.9 was to make the proof of Corollary 3.10 easier. We present the basic strategies for the $\mathcal{R}_{\langle e, i \rangle}$ and $\mathcal{S}_{\langle e, i \rangle}$ requirements and after a brief discussion of their interaction, leave the formal details to the reader. The requirements can be given any priority ordering of order type ω , for example giving $\mathcal{R}_{\langle e, i \rangle}$ the $2\langle e, i \rangle$ priority and giving $\mathcal{S}_{\langle e, i \rangle}$ the $2\langle e, i \rangle + 1$ priority.

We begin our discussion with the action taken for the $\mathcal{S}_{\langle e, i \rangle}$ requirement. As before, when $\mathcal{S}_{\langle e, i \rangle}$ is first eligible to act (say at stage s), it picks a new point $q^{\langle e, i \rangle}$ and places it in the partial order below a point dictated by the higher priority requirements. Let n denote the number of points in P at stage s . $\mathcal{S}_{\langle e, i \rangle}$ forces all lower priority requirements to work below $q^{\langle e, i \rangle}$ until a stage $t > s$ such that all of the computations $\varphi_e^{X_i}(0), \dots, \varphi_e^{X_i}(n)$ appear to converge at stage t . (Notice that we are now approximating the Δ_2^0 set X_i , so we are really looking at computations of the form $\varphi_{e,t}^{X_i}(x)$. For simplicity of notation, we drop explicit mention of the stage from these computations.) At stage t , there are two possibilities: either these computations all converge to points which are not below $q^{\langle e, i \rangle}$, in which case at least two of these computations must converge to the same value and so $\mathcal{S}_{\langle e, i \rangle}$ is currently satisfied, or there is a least $x \leq_{\mathbb{N}} n$ such that $\varphi_e^{X_i}(x)$ appears to converge to a point $y \leq_P q^{\langle e, i \rangle}$. In the former case, $\mathcal{S}_{\langle e, i \rangle}$ takes no action and continues to force lower priority requirements to build below $q^{\langle e, i \rangle}$. In the latter case, $\mathcal{S}_{\langle e, i \rangle}$ picks a currently \leq_P minimal point z below y and requires all lower priority requirements to work below z at stages $\geq t$.

The main difference between this construction and the one in the proof of Theorem 3.9 is that our approximation of X_i may change on small values and hence our apparent computations for $\varphi_e^{X_i}(0), \dots, \varphi_e^{X_i}(n)$ may diverge again at a stage $\geq t$. If these computations change, then $\mathcal{S}_{\langle e, i \rangle}$ initializes all requirements of lower priority and starts over again with the same witness $q^{\langle e, i \rangle}$. That is, it waits for the computations $\varphi_e^{X_i}(0), \dots, \varphi_e^{X_i}(n)$ to converge to new values and either takes no action (if it appears that $\mathcal{S}_{\langle e, i \rangle}$ is satisfied) or it picks a new value of z and forces lower priority requirements to work below this new value of z . The obvious worry about this strategy is that it might repeat itself infinitely many times. However, this is the point at which we can appeal to the fact that X_i is low. Because X_i is low, O' knows whether $\varphi_e^{X_i}(0), \dots, \varphi_e^{X_i}(n)$ really converge or not. Therefore, we have a Δ_2^0 approximation to the question of whether they converge. We alter the above argument by making $\mathcal{S}_{\langle e, i \rangle}$ wait for a stage $t > s$ such that the computations $\varphi_e^{X_i}(0), \dots, \varphi_e^{X_i}(n)$ converge and that our Δ_2^0 approximation to whether these computations really converge says that they do in fact converge. Now we are in the situation that either these computations really do converge, in which case at some finite stage we see their real values and these values never change,

or they do not all converge, in which case there is a finite stage after which we never believe that they all converge and so $\mathcal{S}_{\langle e,i \rangle}$ takes no action after that stage. This description makes it clear that the action of $\mathcal{S}_{\langle e,i \rangle}$ is finitary, and hence (assuming that $\mathcal{S}_{\langle e,i \rangle}$ is only initialized finitely often) it only initialized the lower priority requirements finitely often.

The basic strategy for the $\mathcal{R}_{\langle e,i \rangle}$ requirement is similarly modified from the proof of Theorem 3.9. When $\mathcal{R}_{\langle e,i \rangle}$ is first eligible to act (or at the first stage at which it is eligible to act after being initialized), it picks two new points $q_0^{\langle e,i \rangle}$ and $q_1^{\langle e,i \rangle}$ and forces lower priority requirements to work below $q_0^{\langle e,i \rangle}$. Let n denote the number of points in P at stage s . $\mathcal{R}_{\langle e,i \rangle}$ waits for a stage $t > s$ at which all the computations $\varphi_e^{X_i}(0), \dots, \varphi_e^{X_i}(n)$ appear to converge. There are two cases: either none of these computations converge to a point below either $q_0^{\langle e,i \rangle}$ or $q_1^{\langle e,i \rangle}$ (in which case at least two of the computations must converge to the same point so $\mathcal{R}_{\langle e,i \rangle}$ currently is satisfied) or there is a least $x \leq_{\mathbb{N}} n$ for which $\varphi_e^{X_i}(x)$ appears to converge to a point below one of $q_0^{\langle e,i \rangle}$ or $q_1^{\langle e,i \rangle}$. In the former case, $\mathcal{R}_{\langle e,i \rangle}$ takes no new action. In the latter case, assume that $\varphi_e^{X_i}(x) = y$ and $y \leq_P q_j^{\langle e,i \rangle}$. $\mathcal{R}_{\langle e,i \rangle}$ now forces all lower priority requirements to work below $q_{1-j}^{\langle e,i \rangle}$.

Just as with the $\mathcal{S}_{\langle e,i \rangle}$ strategy, the apparent computations $\varphi_e^{X_i}(0), \dots, \varphi_e^{X_i}(n)$ may later diverge or converge to different values. However, the lowness of X_i guarantees that there is a finite stage at which either these computations settle down or there is a finite stage after which we never again believe that these computations really converge. Therefore, \mathcal{R}_e^i may finitely often change its mind about whether lower priority requirements have to build below $q_0^{\langle e,i \rangle}$ or $q_1^{\langle e,i \rangle}$, but eventually, if $\varphi_e^{X_i}$ is total, then \mathcal{R}_e^i eventually finds true computations and either has to perform no action to diagonalize (because $\varphi_e^{X_i}$ is not one-to-one) or sees $\varphi_e^{X_i}(x) = y \leq_P q_j^{\langle e,i \rangle}$ for a $\leq_{\mathbb{N}}$ least value of x and forces all lower priority requirements to build below $q_{1-j}^{\langle e,i \rangle}$ forever.

As in the proof of Theorem 3.9, there is an obvious source of conflict between these requirements. However, here we have described the solution to this conflict in terms of initializing lower priority strategies. Since the action of each \mathcal{S}_e^i and \mathcal{R}_e^i strategy is finitary, each requirement eventually reached a stage after which it is never initialized and henceforth it behaves exactly as in the description above. This completes the description of the construction and the proof of Theorem 3.11. \square

We next examine how wqo and wqo(ext) are related. RCA_0 suffices to prove the implications $\text{wqo} \rightarrow \text{wqo}(\text{ext})$ and $\text{wqo}(\text{set}) \rightarrow \text{wqo}(\text{ext})$.

Lemma 3.12. $\text{RCA}_0 \vdash \text{wqo} \rightarrow \text{wqo}(\text{ext})$.

Proof. Fix a wqo (Q, \leq) and assume that Q has a linear extension (L, \leq_L) which is not well founded. Recall that this means L is a linear extension of Q / \sim (where $x \sim y$ if and only if $x \leq y$ and $y \leq x$). Fix a function $f : \mathbb{N} \rightarrow L$ such that for all $i < j$, $f(j) <_L f(i)$. Because \leq_L is an extension of \leq , $f(j) <_L f(i)$ implies that either $f(j) < f(i)$ or $f(j) \perp f(i)$. Therefore, viewing f as a map from \mathbb{N} into Q , we see that f contradicts the fact that Q is a wqo. \square

Corollary 3.13. $\text{RCA}_0 \vdash \text{wqo}(\text{set}) \rightarrow \text{wqo}(\text{ext})$.

Proof. This corollary follows from Lemma 3.12 together with the fact that $\text{RCA}_0 \vdash \text{wqo}(\text{set}) \rightarrow \text{wqo}$. \square

We also want to examine the implication $\text{wqo}(\text{ext}) \rightarrow \text{wqo}$. One method for trying to prove this implication for partial orders is as follows. (The general case for quasi-orders follows from this special case.) Let \leq_P be a partial order with domain \mathbb{N} which does not satisfy wqo and let $f : \mathbb{N} \rightarrow P$ be a function which witnesses that P does not satisfy wqo . That is, if $n <_{\mathbb{N}} m$, then $f(n) \not\leq_P f(m)$. We would like to define a linear extension of the partial order $<_P$ under which f enumerates an infinite descending chain. Such a linear extension would show that P does not satisfy $\text{wqo}(\text{ext})$ and hence would finish our proof that $\text{wqo}(\text{ext}) \rightarrow \text{wqo}$. We begin to construct such a linear extension by defining a binary relation R on P : $x R y$ holds if and only if either $x <_P y$ or there are numbers $n <_{\mathbb{N}} m$ such that $x = f(m)$ and $y = f(n)$. Let $<'_P$ be the transitive closure of the relation R . Because R is an acyclic relation (defined below), $<'_P$ is a partial order extending $<_P$. Because we made $f(n+1) R f(n)$ hold for all n , f enumerates an infinite descending chain in $<'_P$. Let $<_L$ be any linear extension of $<'_P$. Then $<_L$ is a linear extension of $<_P$ which is not well founded.

This proof sketch has three parts: defining R , taking the transitive closure, and extending a partial order to a linear order. The definition of R appears to require ACA_0 , but by substituting an appropriate subsequence of f in place of f we can make sure that the new R is definable in RCA_0 . Moreover RCA_0 is strong enough to prove that every partial order has a linear extension (see for example [2]). However, as we show below, taking the transitive closure of an arbitrary relation requires ACA_0 . Before presenting the formal result, we introduce some definitions.

Definition 3.14. (RCA_0) Let A be a set and let $R \subseteq A \times A$ be a relation on A . A *chain* in R is a finite sequence $\langle x_0, \dots, x_k \rangle$ of elements of A such that $x_0 R x_1 R \dots R x_k$. R is called *acyclic* if there is no chain $\langle x_0, \dots, x_k \rangle$ in R such that $x_k R x_0$. The *transitive closure* of R is the relation R' defined by $i R' j$ if and only if there is a chain $\langle x_0, \dots, x_k \rangle$ in R such that $x_0 = i$ and $x_k = j$.

Lemma 3.15. (RCA_0) *The following are equivalent:*

1. ACA_0 ;
2. *the transitive closure of every acyclic relation exists.*

Proof. The implication from 1 to 2 follows because the transitive closure is arithmetically definable. To prove the implication from 2 to 1, it suffices to show that the range of every one-to-one function exists. We reason within RCA_0 . Fix a one-to-one function f and consider the set $\{b\} \cup \{x_n | n \in \mathbb{N}\} \cup \{y_n | n \in \mathbb{N}\}$. Define an acyclic relation R on this set by $R = \{\langle b, x_n \rangle | n \in \mathbb{N}\} \cup \{\langle x_n, y_{f(n)} \rangle | n \in \mathbb{N}\}$. By 2, we let R' be the transitive closure of R . Then, $m \in \text{range}(f)$ if and only if $\langle b, y_m \rangle \in R'$. \square

Because of Lemma 3.15, we cannot naively form the transitive closure in the proof sketched above. However, notice that we do not necessarily need the transitive closure of R , but rather we only need some extension of R which is a partial order. The ability to extend any acyclic relation to a partial order turns out to be equivalent to WKL_0 .

Lemma 3.16. (RCA_0) *The following are equivalent:*

1. WKL_0 ;
2. *every acyclic relation can be extended to a partial order.*

Proof. First, we show that WKL_0 suffices to prove that every acyclic relation can be extended to a partial order. Let R be an acyclic relation on A . To simplify our notation, we assume that $A = \mathbb{N}$. We define a tree $T \subseteq 2^{<\omega}$ such that every path through T codes a partial order extending R . We view all sequences $s \in 2^{<\mathbb{N}}$ as sequences of pairs of elements from \mathbb{N} and we think of s as giving a finite approximation $<_s$ to a partial order extending R . That is, we interpret $s(\langle n, m \rangle) = 1$ as specifying $n <_s m$ and $s(\langle n, m \rangle) = 0$ as specifying $n \not<_s m$. We put $s \in T$ if and only if

1. $m R n$ and $\langle m, n \rangle \in \text{dom}(s)$ imply $s(\langle m, n \rangle) = 1$;
2. $s(\langle l, m \rangle) = 1$, $s(\langle m, n \rangle) = 1$ and $\langle l, n \rangle \in \text{dom}(s)$ imply $s(\langle l, n \rangle) = 1$;
3. $\langle n, m \rangle, \langle m, n \rangle \in \text{dom}(s)$ implies $s(\langle m, n \rangle) + s(\langle n, m \rangle) \leq 1$;
4. $\langle n, n \rangle \in \text{dom}(s)$ implies $s(\langle n, n \rangle) = 0$.

It is clear that T is a tree, and we verify that it is infinite. Fix $N \in \mathbb{N}$ and for each pair $\langle n, m \rangle < N$, define $s(\langle n, m \rangle) = 1$ if and only if there is a chain $\langle x_0, \dots, x_k \rangle$ in R such that each $x_i < N$ and $x_0 = n$ and $x_k = m$. It is straightforward to verify, using the fact that R is acyclic, that $s \in T$. Therefore, T is infinite and by WKL_0 it has a path. If h is any path in T , then define $<_h$ by $n <_h m$ if and only if $h(\langle n, m \rangle) = 1$. By our definition of T , \leq_h is a partial order extending R .

Second, we show that 2 implies 1 in the statement of the lemma. To show WKL_0 , it suffices to consider an arbitrary pair of one-to-one functions $f, g : \mathbb{N} \rightarrow \mathbb{N}$ with disjoint ranges and show that there is a set X such that for all n , $f(n) \in X$ and $g(n) \notin X$. We define our relation on the set $\{b\} \cup \{x_n | n \in \mathbb{N}\} \cup \{y_n | n \in \mathbb{N}\} \cup \{z_n | n \in \mathbb{N}\}$ by

$$R = \{\langle x_n, b \rangle | n \in \mathbb{N}\} \cup \{\langle b, y_n \rangle | n \in \mathbb{N}\} \cup \{\langle z_{f(n)}, x_n \rangle | n \in \mathbb{N}\} \cup \{\langle y_n, z_{g(n)} \rangle | n \in \mathbb{N}\}.$$

Because f and g have disjoint ranges, this relation is acyclic. By 1, there is a partial order $<_P$ which extends R . Let $X = \{m | z_m <_P b\}$ and notice that X is our desired separating set. \square

Theorem 3.17. $\text{WKL}_0 \vdash \text{wqo}(\text{ext}) \rightarrow \text{wqo}$.

Proof. We reason in WKL_0 . Fix a quasi-order Q which does not satisfy wqo and let P be the partial order Q / \sim . Since Q does not satisfy wqo , neither does P . Therefore, we can fix a function $f : \mathbb{N} \rightarrow P$ such that for all $n < m$, $f(n) \not\leq_P f(m)$. In particular, this means that f is one-to-one. Furthermore, we can define a subsequence $g : \mathbb{N} \rightarrow P$ of f as follows. Set $g(0) = f(0)$. If $g(n) = f(i)$, then let $g(n+1) = f(j)$ where j is the $\leq_{\mathbb{N}}$ least number such that $i <_{\mathbb{N}} j$ and $g(n) <_{\mathbb{N}} f(j)$. Such a j exists because f is one-to-one. It is clear from our definition of g that for all $n < m$, $g(n) \not\leq_P g(m)$ and for all n , $g(n) \geq_{\mathbb{N}} n$. To show that Q does not satisfy $\text{wqo}(\text{ext})$, it suffices to produce a linear extension \leq_L of \leq_P which is not well ordered. We use the function g to give an infinite descending sequence in \leq_L by making $g(n+1) <_L g(n)$ for all n .

Define an acyclic relation R on P by $\langle i, j \rangle \in R$ if and only if $i <_P j$ or there are $n \leq_{\mathbb{N}} j$ and $m \leq_{\mathbb{N}} i$ such that $g(m) = i$, $g(n) = j$ and $n <_{\mathbb{N}} m$. It follows from our hypothesis on g that R is an acyclic relation. Therefore, by Lemma 3.16, there is a partial order $<'_P$ extending R . It is clear that $<'_P$ extends $<_P$ and that g enumerates an infinite descending sequence in $<'_P$. Let $<_L$ be any linear extension of $<'_P$ and notice that $<_L$ is a linear extension of $<_P$ such that $<_L$ has an infinite descending chain given by g . \square

Corollary 3.18. $\text{WKL}_0 \vdash \text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{anti})$.

Proof. This result follows from Theorem 3.17 together with the fact that $\text{RCA}_0 \vdash \text{wqo} \rightarrow \text{wqo}(\text{anti})$ (from Theorem 2.5). \square

Corollary 3.19. $\text{WKL}_0 \not\vdash \text{wqo}(\text{anti}) \rightarrow \text{wqo}(\text{ext})$.

Proof. By Theorem 3.17, $\text{WKL}_0 \vdash \text{wqo}(\text{ext}) \rightarrow \text{wqo}$, and by Theorem 3.11, $\text{WKL}_0 \not\vdash \text{wqo}(\text{anti}) \rightarrow \text{wqo}$. \square

Notice that while WKL_0 suffices to prove most of the implications from $\text{wqo}(\text{ext})$, it does not suffice to prove $\text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{set})$. (Recall that WKL_0 is not strong enough to prove $\text{RT}_{<\infty}^1$.)

Lemma 3.20. $\text{RCA}_0 \vdash (\text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{set})) \rightarrow \text{RT}_{<\infty}^1$ and hence $\text{WKL}_0 \not\vdash \text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{set})$.

Proof. Let k denote the trivial partial order on $\{0, \dots, k-1\}$ in which all elements are incomparable. RCA_0 is strong enough to prove this partial order is a wqo , and hence by Lemma 3.12 this partial order satisfies $\text{wqo}(\text{ext})$. Consider any map $f : \mathbb{N} \rightarrow k$. Since k satisfies $\text{wqo}(\text{ext})$, the implication $\text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{set})$ gives an infinite set A such that for all $i < j \in A$, $f(i) \leq f(j)$ in the order on k . However, since all elements of k are incomparable, $f(i) \leq f(j)$ implies that $f(i) = f(j)$ and hence A is the desired homogeneous set. \square

Lemma 3.20 does not rule out the possibility that $\text{REC} \models \text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{set})$ since $\text{REC} \models \text{RT}_{<\infty}^1$. However, a corollary of the next theorem will eliminate this possibility.

Theorem 3.21. $\text{REC} \not\models \text{wqo}(\text{ext}) \rightarrow \text{wqo}$ and hence $\text{RCA}_0 \not\vdash \text{wqo}(\text{ext}) \rightarrow \text{wqo}$.

Proof. We build a computable partial order (P, \leq_P) with $P = \mathbb{N}$ such that every computable linear extension of \leq_P is well ordered with respect to REC . Formally, for all $e, i \in \mathbb{N}$, we meet the requirement $\mathcal{R}_{\langle e, i \rangle}$ which says that if φ_e and φ_i are both total and φ_e is a linear extension of \leq_P , then there exist $n < m$ such that $\varphi_i(n) \leq_{\varphi_e} \varphi_i(m)$. That is, if φ_e is a linear extension of \leq_P , then φ_i is not an infinite descending chain in the order given by φ_e .

Furthermore, we need to make sure that \leq_P is not a well quasi-order with respect to REC by defining a computable function f such that for all $i < j$, either $f(i) \perp f(j)$ or $f(j) <_P f(i)$. Our function f will in fact have the stronger property that it defines a computable antichain: for all $i \neq j$, $f(i) \perp f(j)$.

We first describe the module for a requirement $\mathcal{R}_{\langle e, i \rangle}$ working in isolation. (In the general construction, $\mathcal{R}_{\langle e, i \rangle}$ will be working inside an interval given by higher priority requirements. If $\mathcal{R}_{\langle e, i \rangle}$ is the highest priority requirement, then this interval is defined by the least and the greatest point in our partial order. Because these points play no role in the action of $\mathcal{R}_{\langle e, i \rangle}$ other than to dictate where it takes place, we ignore them for the moment.) In order to construct our computable function f , we begin to define an antichain of points $f(0), f(1), \dots$, placing one new point in this antichain at each stage of the construction. $\mathcal{R}_{\langle e, i \rangle}$ picks two new points (which are kept out of the range of f) $x_{\langle e, i \rangle}$ and $y_{\langle e, i \rangle}$ and makes them incomparable in \leq_P with each other and with all the points in the range of f . Until φ_e gives us an order on these two auxiliary points, $\mathcal{R}_{\langle e, i \rangle}$ does not believe that φ_e is total and so does nothing to diagonalize. Assume that φ_e eventually does give us

an order on our auxiliary points and that it makes $x_{\langle e,i \rangle} <_{\varphi_e} y_{\langle e,i \rangle}$. After this stage, we begin to place all the new points of P below $x_{\langle e,i \rangle}$. That is, we continue to build an antichain for the range of f , but we place these points below $x_{\langle e,i \rangle}$. (In the general construction, we also force all lower priority requirements to place their new points below $x_{\langle e,i \rangle}$.) $\mathcal{R}_{\langle e,i \rangle}$ now waits for φ_i to enumerate a point below $x_{\langle e,i \rangle}$. Since there are only finitely many points in P which are not below $x_{\langle e,i \rangle}$, we win $\mathcal{R}_{\langle e,i \rangle}$ unless φ_i picks one of these points. Once we see some n for which $\varphi_i(n)$ converges to a point $\leq_P x_{\langle e,i \rangle}$, we switch our tactics again and begin to build the antichain given by f above $y_{\langle e,i \rangle}$. At this point, $\mathcal{R}_{\langle e,i \rangle}$ promises to continue the tactic of allowing new points only above $y_{\langle e,i \rangle}$ for the rest of the construction. Since there are only finite many points which are not above $y_{\langle e,i \rangle}$, we have that either φ_i is not total, or it has finite range, or $\varphi_i(m)$ eventually converges for some $m >_{\mathbb{N}} n$ with $y_{\langle e,i \rangle} \leq_P \varphi_i(m)$. If φ_e really is a linear extension of \leq_P , then the numbers n and m are our witnesses that $\mathcal{R}_{\langle e,i \rangle}$ succeeds since $n <_{\mathbb{N}} m$ and $\varphi_i(n) \leq_P x_{\langle e,i \rangle} <_{\varphi_e} y_{\langle e,i \rangle} \leq_P \varphi_i(m)$, so that $\varphi_i(n) \leq_{\varphi_e} \varphi_i(m)$.

The general construction just iterates the above procedure. The tree of strategies is the full ternary branching tree on the letters a, b, w with a to the left of b and b to the left of w . The nodes α at level $\langle e, i \rangle$ of this tree are the strategies working for $\mathcal{R}_{\langle e,i \rangle}$. A strategy α keeps four parameters: two witnesses x_α and y_α which are kept out of the range of f , and two points $u_\alpha <_P v_\alpha$ which tell the requirements of lower priority to work in the interval between u_α and v_α . The first time a strategy α is eligible to act, it picks witnesses x_α and y_α , adds these point to P , makes them incomparable in \leq_P and in the appropriate interval as dictated by higher priority requirements, and keeps them out of the range of f . While α is waiting for φ_e to give an order to x_α and y_α , it takes outcome $\alpha * w$ and sets its parameters u_α and v_α to define the same interval that α was given by the higher priority requirements. Once an order between these nodes is established (and without loss of generality we assume $x_\alpha <_{\varphi_e} y_\alpha$), it takes outcome $\alpha * b$ and forces future points in the range of f and witnesses for lower priority requirements to be placed below x_α . To pass this information to the lower priority requirements, it resets its parameter $v_\alpha = x_\alpha$ and leaves the parameter u_α unchanged. If φ_i ever gives a computation $\varphi_i(n)$ with value $\leq_P x_\alpha$, then α switches to outcome $\alpha * a$ and forces future points in the range of f and witnesses for lower priority requirements to be placed above y_α . It passes this information to lower priority requirements by resetting its parameters so that $u_\alpha = y_\alpha$ and v_α is the upper bound passed down from the higher priority requirements. That is, v_α returns to the original value that α was given before it saw that $x_\alpha <_{\varphi_e} y_\alpha$.

The construction proceeds as follows. At stage 0, we begin by placing 0 as the least point in \leq_P and 1 as the greatest point in \leq_P . We define $f(0) = 2$ and let \mathcal{R}_λ set $x_\lambda = 3$ and $y_\lambda = 4$. These three points are pairwise incomparable in \leq_P and between the greatest and least points in \leq_P . \mathcal{R}_λ defines $u_\lambda = 0$ and $v_\lambda = 1$.

At stage $t > 0$, we assume that we have already defined $f(0), \dots, f(t-1)$ to be a finite antichain. The module for a strategy α that is working for $\mathcal{R}_{\langle e,i \rangle}$ and that is eligible to act is as follows. (As in the previous constructions, α^- denotes the strategy of immediately higher priority than α . Notice that for stages $t > 0$, the highest priority strategy λ will not act as in step 1 below, so it does not matter that λ^- is not defined.)

1. If this is the first stage at which α is eligible to act, then we let x_α and y_α be the $\leq_{\mathbb{N}}$ -least numbers not in P yet. α has been passed parameters u_{α^-} and v_{α^-} from α^- . The points x_α and y_α are placed in \leq_P as follows: they are incomparable with each other; if $z \leq_P u_{\alpha^-}$, then

$z <_P x_\alpha, y_\alpha$; if $v_{\alpha^-} \leq_P z$, then $x_\alpha, y_\alpha <_P z$; and they are incomparable with all other points. Let $\alpha * w$ be the next strategy eligible to act. Set $u_\alpha = u_{\alpha^-}$, set $v_\alpha = v_{\alpha^-}$ and pass these parameters to $\alpha * w$.

2. If α has been eligible to act before, then check if φ_e defines an order on x_α and y_α yet. If not, then let $\alpha * w$ be the next to act and leave the parameters u_α or v_α as they are currently defined. If so, then without loss of generality, assume that $x_\alpha <_{\varphi_e} y_\alpha$. Set $v_\alpha = x_\alpha$ and leave u_α as it is currently defined. Proceed to step 3.
3. Check if $\varphi_{i,t}(n)$ converges for some $n \leq_{\mathbb{N}} t$ such that $\varphi_{i,t}(n) \leq_P x_\alpha$. If not, then let $\alpha * b$ be the next strategy eligible to act and pass the parameters u_α and v_α to $\alpha * b$. Otherwise, if there is such an n , set $u_\alpha = y_\alpha$ and set $v_\alpha = v_{\alpha^-}$. Let $\alpha * a$ the next strategy eligible to act and pass the parameters u_α and v_α to $\alpha * a$.

Once we have defined the current true path δ_t at this stage, we need to define $f(t)$ to be the next element of our antichain. Let $f(t)$ be the $\leq_{\mathbb{N}}$ -least number not in P yet and let $\alpha = \delta_t$ be the last strategy that was eligible to act. Place $f(t)$ in \leq_P as follows: if $z \leq_P u_\alpha$, then $z <_P f(t)$; if $v_\alpha \leq_P z$, then $f(t) <_P z$; and $f(t)$ is incomparable with all other points in \leq_P at this stage. This completes the description of the construction at stage t .

We sketch the verification that this construction succeeds. The construction is finite injury and the current true path can only move left between stages of the construction. As before, the initialization of lower priority strategies is taken care of by the movement of the current true path and we can define the true path to be the pointwise limit of the current true paths. The individual modules for requirements along the true path are satisfied just as in the informal description.

We need to see that the range of f is an antichain in P . We prove this fact by a sequence of claims, each of which can be verified by induction on the stages of the construction. First, once a point in the range of f is placed in P , nothing is subsequently placed above it or below it. Second, any point marked as a parameter u_α may have points from the range of f above it in \leq_P , but it will never have any points from the range of f below it in \leq_P . Similarly, a point marked v_α may have points from the range of f below it, but it will never have any points from the range of f above it. Given these claims, it is clear from the definition of $f(t)$ that when it is placed in P , there are no points from the range of f either above it or below it. This property is then maintained for the rest of the construction. \square

We mention one corollary which is stated in the terminology of effective algebra rather than reverse mathematics. While similar corollaries could be stated for several of the earlier results, we mention this one in particular because it seems interesting as a statement independent from the general project of this paper.

Corollary 3.22. *There is a computable partial order (P, \leq_P) such that P has a computable infinite antichain and yet every computable linear extension of P is computably well ordered.*

Proof. This corollary follows directly from the proof of Theorem 3.21 since our computable function f is an infinite antichain. \square

Corollary 3.23. $\text{REC} \not\models \text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{anti})$ and hence $\text{RCA}_0 \not\models \text{wqo}(\text{ext}) \rightarrow \text{wqo}(\text{anti})$.

Proof. Let P be a computable partial order as in Corollary 3.22. REC is a model for the fact that P satisfies $wqo(\text{ext})$ but is not a model for P satisfying $wqo(\text{anti})$. \square

Corollary 3.24. $\text{REC} \not\models wqo(\text{ext}) \rightarrow wqo(\text{set})$.

Proof. Suppose that $\text{REC} \models wqo(\text{ext}) \rightarrow wqo(\text{set})$. Since $\text{RCA}_0 \vdash wqo(\text{set}) \rightarrow wqo$, we have that $\text{REC} \models wqo(\text{ext}) \rightarrow wqo$ which contradicts Theorem 3.21. \square

4 Closure properties for well quasi-orders

If (Q_1, \preceq_1) and (Q_2, \preceq_2) are quasi-orders, then the product quasi-order on $Q_1 \times Q_2$ is given by $\langle q_1, q_2 \rangle \preceq \langle r_1, r_2 \rangle$ if and only if $q_1 \preceq_1 r_1$ and $q_2 \preceq_2 r_2$. In this section, we examine how difficult it is to show that the various properties defining well quasi-orders are closed under intersection and products. That is, we consider each pair of properties P_1, P_2 from $wqo, wqo(\text{set}), wqo(\text{anti})$ and $wqo(\text{ext})$ and examine how difficult it is to show that if \preceq_1 and \preceq_2 are quasi-orders satisfying P_1 on Q , then $\preceq_1 \cap \preceq_2$ satisfies P_2 on Q , and to show that if (Q_1, \preceq_1) and (Q_2, \preceq_2) are quasi-orders satisfying P_1 , then the product order on $Q_1 \times Q_2$ satisfies P_2 . We begin by examining the closure properties under intersection and then note that by Lemma 5.20 in [7], these results hold for products as well.

Lemma 4.1. RCA_0 suffices to prove that if \preceq_1 and \preceq_2 satisfy $wqo(\text{set})$, then $\preceq_1 \cap \preceq_2$ satisfies $wqo(\text{set})$.

Proof. Assume that \preceq_1 and \preceq_2 satisfy $wqo(\text{set})$ and fix a map $f : \mathbb{N} \rightarrow Q$. Viewing f as a map from \mathbb{N} into the domain of \preceq_1 , there is an infinite set A_1 such that for all $i <_{\mathbb{N}} j \in A_1$, $f(i) \preceq_1 f(j)$. Now consider the restriction of f to A_1 . There is an infinite set $A_2 \subseteq A_1$ such that for all $i <_{\mathbb{N}} j \in A_2$, $f(i) \preceq_2 f(j)$. Because $A_2 \subseteq A_1$, we have that for all $i <_{\mathbb{N}} j \in A_2$, $f(i) \preceq_1 f(j)$ and hence $f(i)$ is below $f(j)$ in $\preceq_1 \cap \preceq_2$. (More formally, to define A_2 , we enumerate A_1 as $a_0 <_{\mathbb{N}} a_1 <_{\mathbb{N}} \dots$ and define a map $f' : \mathbb{N} \rightarrow Q$ by $f'(n) = f(a_n)$. We then apply $wqo(\text{set})$ for \preceq_2 to the map f' .) \square

Corollary 4.2. Let P be any of the properties $wqo, wqo(\text{ext}), wqo(\text{anti})$ or $wqo(\text{set})$. RCA_0 is strong enough to prove that if \preceq_1 and \preceq_2 satisfy $wqo(\text{set})$, then $\preceq_1 \cap \preceq_2$ satisfies P .

Proof. This corollary follows since RCA_0 is strong enough to prove that $wqo(\text{set})$ implies each of the other properties defining well quasi-orders. \square

We next show that WKL_0 is not strong enough to prove the closure under intersections for any of the remaining properties. Let X_0, X_1, \dots be a sequence of uniformly Δ_2^0 and uniformly low sets such that the ω -model \mathcal{A} with second order part $\{Y \mid \exists i(Y \leq_T X_i)\}$ is a model of WKL_0 . The following theorem shows that \mathcal{A} is not a model for the closure of quasi-orders satisfying wqo under intersection.

Theorem 4.3. There are computable partial orders \leq_0 and \leq_1 such that both \leq_0 and \leq_1 satisfy wqo in \mathcal{A} but $\leq_0 \cap \leq_1$ is an infinite antichain.

Proof. We build \leq_0 and \leq_1 in stages so that each is a partial order on \mathbb{N} . We guarantee that each satisfies wqo in \mathcal{A} by meeting the following requirements for $e, i \in \mathbb{N}$ and $j \in \{0, 1\}$.

$$\mathcal{R}_{\langle e, i, j \rangle} : \varphi_e^{X_i} \text{ total and one-to-one} \rightarrow \exists k < l (\varphi_e(k) \leq_j \varphi_e(l))$$

In fact, we will have the additional property that each \leq_j has no infinite descending chains at all (that is even outside of \mathcal{A}) since $n \leq_j m$ will imply $n \leq_{\mathbb{N}} m$. We make $\leq_0 \cap \leq_1$ an antichain by guaranteeing that if $n <_j m$, then $n \not\leq_{1-j} m$. We refer to this requirement as our antichain requirement. We order the priority for our $\mathcal{R}_{\langle e, i, j \rangle}$ requirements in type ω by ordering the triples $\langle e, i, j \rangle$ in type ω .

Consider first the basic module for the requirement $\mathcal{R}_{\langle 0, 0, 0 \rangle}$. $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ picks two points q^0 and q^1 and makes them incomparable in both the \leq_0 and \leq_1 orders. It forces the \leq_0 partial order to place all its points above q^0 and the \leq_1 order to place all its points above q^1 . This guarantees that $q^j \leq_j x$ for all $x \neq q^{1-j}$, while $q^j \not\leq_{1-j} x$ for all x . We therefore are meeting our requirements for making $\leq_0 \cap \leq_1$ an antichain with respect to the points q^0 and q^1 . Currently, there is exactly one point (the point q^1) which is not $\geq_0 q^0$ in the \leq_0 order. We wait for the computations $\varphi_0^{X_0}(0)$ and $\varphi_0^{X_0}(1)$ to appear to converge. If one of these computations never appears to converge, then $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ is easily won since $\varphi_0^{X_0}$ is not total. Therefore, assume that these computations do appear to converge at some stage s . We have two cases to consider. Either both computations appear to converge to q^1 , in which case $\varphi_0^{X_0}$ does not appear to be one-to-one and so $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ takes no additional action, or one of these computations converges to a point y such that $q^0 \leq_0 y$. At this stage, $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ initializes all lower priority requirements (that is makes them begin again with new witnesses) and forces them to work above y in the \leq_0 order. Since at stage s there are only finitely many points so far not above y in the \leq_0 order, this guarantees that if $\varphi_0^{X_0}$ is total and one-to-one then $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ will be met.

When $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ begins to force lower priority requirements to work above y in \leq_0 , we also need to consider where to make lower priority requirements work in the \leq_1 order. Because each lower priority requirement works in a similar manner to $\mathcal{R}_{\langle 0, 0, 0 \rangle}$, points are always placed in \leq_0 and \leq_1 in pairs. That is, the point y is one point in a pair of points z^0, z^1 which were placed in the partial orders as witnesses for a lower priority requirement. Without loss of generality, we suppose that $y = z^1$. The points z^0, z^1 were made incomparable in both partial orders and so when $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ forces lower priority requirements to work above $y = z^1$ in the \leq_0 order, it also forces all lower priority requirements to work above the other element of the pair, in this case z^0 , in the \leq_1 order. In this fashion we maintain our antichain requirements.

As in the proof of Theorem 3.11, the computations $\varphi_0^{X_0}(0)$ and $\varphi_0^{X_0}(1)$ may change at a later stage. Each time these computations change values, $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ initializes the lower priority requirements and picks a new value of $y \geq_0 q^0$ to force lower priority requirements to work above in \leq_0 and uses the corresponding element to force requirements to work above in \leq_1 . The fact that X_0 is Δ_2^0 and low means that after finitely many such changes, we reach a stage at which we see the true value of the computations or after which we never believe both computations converge. Therefore, the action of $\mathcal{R}_{\langle 0, 0, 0 \rangle}$ is finitary.

The general construction just iterates this strategy for one requirement. Because the action of a single $\mathcal{R}_{\langle e, i, j \rangle}$ strategy is finitary, the argument is finite injury. It can be formalized with a tree of strategies as in Theorem 3.9, although it is easier to formalize without the extra machinery of the

tree of strategies. We describe a formal module for $\mathcal{R}_{\langle e, i, j \rangle}$ below and leave the details of combining these strategies in a finite injury argument to the reader.

Let α be a triple $\langle e, i, j \rangle$. The requirement \mathcal{R}_α keeps five pieces of information: two points q_α^0 and q_α^1 which it picks and adds to the partial orders when it is first eligible to act (and repicks when it is initialized), two parameters u_α^0 and u_α^1 which it uses to tell requirements of lower priority where to place their points, and a number n_α which gives the length of convergence of $\varphi_e^{X_i}$ that \mathcal{R}_α is waiting for. It requires that lower priority requirements place points in the \leq_k order in the cone about u_α^k . The module for \mathcal{R}_α is as follows. (We let \mathcal{R}_{α^-} denote the requirement of immediately higher priority than \mathcal{R}_α .)

1. The first time α is eligible to act or the first time it is eligible to act after having been initialized, it takes the two $\leq_{\mathbb{N}}$ -least numbers not yet in the domain of our partial orders and uses them as the witnesses q_α^0 and q_α^1 . In the \leq_0 order, it places them immediately above $u_{\alpha^-}^0$, incomparable with each other and incomparable with all nodes which are not below $u_{\alpha^-}^0$. In the \leq_1 order, it places them immediately above $u_{\alpha^-}^1$, incomparable with each other and incomparable with all nodes which are not below $u_{\alpha^-}^1$. Set $u_\alpha^k = q_\alpha^k$ and let n_α denote the number of points which are currently in the partial orders.
2. Wait for a stage t such that $\varphi_e^{X_i}(x)$ appears to converge for all $x \leq_{\mathbb{N}} n_\alpha$. If $\varphi_e^{X_i}$ is not one-to-one on these points, then \mathcal{R}_α takes no further action at this stage. Otherwise, let x be the $\leq_{\mathbb{N}}$ -least point $\leq_{\mathbb{N}} n_\alpha$ such that $\varphi_e^{X_i}(x)$ appears to converge to a point y such that $q_\alpha^j \leq_j y$.
3. When this convergence occurs, initialize all lower priority requirements and set $u_\alpha^j = y$. The point y is q_k^β for some lower priority strategy \mathcal{R}_β and some $k \in \{0, 1\}$. Set $u_\alpha^{1-j} = q_{1-k}^\beta$. If some computation for $\varphi_e^{X_i}(z)$ with $z \leq_{\mathbb{N}} n_\alpha$ changes, return to step 2. Otherwise, the values of u_α^0 and u_α^1 remain fixed.

We sketch the verification that this construction succeeds. The construction is finite injury, so each requirement \mathcal{R}_α eventually stops being initialized and defines final values for q_α^0 , q_α^1 and n_α . After finitely many more stages, the computations $\varphi_e^{X_i}(z)$ for $z \leq_{\mathbb{N}} n_\alpha$ settle down and the values of u_α^0 and u_α^1 stabilize. At this point, \mathcal{R}_α is met exactly as in the informal description of $\mathcal{R}_{\langle 0, 0, 0 \rangle}$. Therefore, both \leq_0 and \leq_1 satisfy wqo in \mathcal{A} .

Second, we check the antichain requirements. It is clear by induction on the stages of the construction that at all stages and for all triples $\beta = \langle e, i, j \rangle$, the points u_β^0 and u_β^1 are incomparable in both the \leq_0 and \leq_1 orders. Furthermore, in both \leq_0 and \leq_1 , the cones above u_β^0 and u_β^1 are disjoint. For a contradiction, assume that there are points x and y such that $x <_0 y$ and $x <_1 y$. In this case, we must have $x <_{\mathbb{N}} y$ and hence y is placed in the partial orders after x . Let s be the stage at which y is placed in the partial orders and let β denote the triple such that \mathcal{R}_β places y in the partial orders. Since $x <_0 y$, \mathcal{R}_β must be forced to work in the cone above x by a higher priority requirement. That is, there is a triple α such that \mathcal{R}_α has higher priority than \mathcal{R}_β and $x = u_\alpha^0$ at stage s . However, in this situation, \mathcal{R}_β is forced to make $u_\alpha^1 <_1 y$ and hence to make $x = u_\alpha^0 \not<_1 y$, giving us the desired contradiction. \square

Corollary 4.4. *Let P_1 be any property from among wqo , $wqo(ext)$ and $wqo(anti)$. Let P_2 be any property from among wqo , $wqo(set)$, $wqo(ext)$ and $wqo(anti)$. WKL_0 is not strong enough to prove that if \leq_0 and \leq_1 satisfy P_1 then $\leq_0 \cap \leq_1$ satisfies P_2 .*

Proof. Fix P_1 and P_2 from the lists of properties in the statement of the corollary and consider the partial orders \leq_0 and \leq_1 and the model \mathcal{A} of WKL_0 from Theorem 4.3. Since RCA_0 suffices to prove $wqo \rightarrow wqo(ext)$ and $wqo \rightarrow wqo(anti)$, we have that \leq_0 and \leq_1 satisfy P_1 in \mathcal{A} . The fact that $\leq_0 \cap \leq_1$ is an infinite antichain means that it does not satisfy P_2 . (If P_2 is $wqo(ext)$, then the linear order defined by $n \leq' m$ if and only if $n \geq_{\mathbb{N}} m$ is a non well-founded linear extension of $\leq_0 \cap \leq_1$.) \square

The following theorem allows us to transfer our results on intersection to products. The proof for the case when P is the property wqo is contained in Lemma 5.20 of [7] and essentially the same proof works for each of the other properties P .

Lemma 4.5 (Marcone). *Let P be any of the properties wqo , $wqo(set)$, $wqo(ext)$ or $wqo(anti)$. Over RCA_0 , the following statements are equivalent.*

1. *If Q_1 and Q_2 satisfy P , then $Q_1 \times Q_2$ with the product order satisfies P .*
2. *If Q satisfies P with respect to \leq_1 and \leq_2 , then Q satisfies P with respect to $\leq_1 \cap \leq_2$.*

Corollary 4.6. *Let P be any of the properties wqo , $wqo(ext)$, $wqo(anti)$ or $wqo(set)$. RCA_0 is strong enough to prove that if (Q_1, \leq_1) and (Q_2, \leq_2) satisfy $wqo(set)$, then $Q_1 \times Q_2$ under the product order satisfies P .*

Corollary 4.7. *Let P_1 be any property from among wqo , $wqo(ext)$ and $wqo(anti)$. Let P_2 be any property from among wqo , $wqo(set)$, $wqo(ext)$ and $wqo(anti)$. WKL_0 is not strong enough to prove that if (Q_1, \leq_1) and (Q_2, \leq_2) satisfy P_1 then $Q_1 \times Q_2$ under the product order satisfies P_2 .*

5 Equivalent definitions for better quasi-orders

We turn our attention to the notion of a better quasi-order. In this section, all strings are finite increasing sequences in $[\mathbb{N}]^{<\omega}$. We think of each string as enumerating a finite set in strictly increasing order. Similarly, we equate each infinite set X with the \mathbb{N} -sequence of the elements of X in strictly increasing order. For strings s and t , we continue to write $s \sqsubseteq t$ if s is an initial segment of t , but we also write $s \subseteq t$ (or $s \subset t$) if the finite set represented by s is a subset (or strict subset respectively) of the finite set represented by t . Similarly, we write $n \in s$ if n is an element of the finite set represented by s .

Given a set $B \subseteq \mathbb{N}^{<\omega}$, we define $\text{base}(B)$ to be $\bigcup B$, or more formally

$$\text{base}(B) = \{n \mid \exists s \in B (n \in s)\}.$$

It is easy to show that the existence of $\text{base}(B)$ for arbitrary B is equivalent to ACA_0 (see Lemma 1.4 in [7]). Therefore, when we make statements like “ $\text{base}(B)$ is infinite” in a subsystem weaker than ACA_0 , we mean the formal statement $\forall m \exists n > m \exists s \in B (n \in s)$. Similarly, when we say X is a subset of $\text{base}(B)$, we mean $\forall x \in X \exists s \in B (x \in s)$.

Definition 5.1. (RCA_0) $B \subseteq [\mathbb{N}]^{<\omega}$ is a *block* if

1. $\text{base}(B)$ is infinite;
2. $\forall X \in [\text{base}(B)]^\omega \exists s \in B (s \sqsubset X)$;
3. $\forall s, t \in B (s \not\sqsubset t)$.

B is a *barrier* if it satisfies the first two conditions above and the (stronger) third condition that $\forall s, t \in B (s \not\subset t)$. Notice that every barrier is a block.

Definition 5.2. (RCA_0) Let s and t be strings. We write $s \triangleleft t$ if there are strings u, v such that $u = u(0) * v$ (that is, v is obtained by deleting the first element of u), $s \sqsubseteq u$ and $t \sqsubseteq v$.

Definition 5.3. (RCA_0) A quasi-order (Q, \preceq) is a *better quasi-order* (or *bqo*) if for every barrier B and every function $f : B \rightarrow Q$, there exist $s, t \in B$ such that $s \triangleleft t$ and $f(s) \preceq f(t)$.

Classically, there are a number of equivalent definitions for a bqo. First, we can restrict the definition by only considering barriers B such that $\text{base}(B) = \mathbb{N}$. Lemma 1.6 in [7] shows that RCA_0 suffices to prove that $\text{base}(B)$ exists for a barrier B . Therefore, in RCA_0 , the definition of a bqo in terms of barriers with $\text{base } \mathbb{N}$ is equivalent to the one given above.

Second, we can replace the barriers in the definition of a bqo by blocks and we can restrict the use of blocks to those blocks B with $\text{base}(B) = \mathbb{N}$. Below, we prove that RCA_0 suffices to prove that $\text{base}(B)$ exists for any block B . This result shows that, over RCA_0 , it is equivalent to define a bqo in terms of blocks and in terms of blocks with $\text{base } \mathbb{N}$. Finally, we present a method for passing from blocks to barriers which was first used by Marcone in [6] and we prove that this method works in WKL_0 . Together, these results show that in WKL_0 the definition of bqo in terms of barriers is equivalent to the definition of bqo in terms of blocks.

Third, one could consider the condition $\text{bqo}(\text{set})$, which is defined analogously to $\text{wqo}(\text{set})$. If B is a barrier and Y is an infinite subset of $\text{base}(B)$, then there is a unique barrier $B' \subseteq B$ such that $\text{base}(B') = Y$: $B' = \{s \in B \mid s \subset Y\}$. In this situation, we call B' a *subbarrier* of B . A quasi-order (Q, \preceq) satisfies $\text{bqo}(\text{set})$ if for all barriers B and all functions $f : B \rightarrow Q$, there exists a subbarrier $B' \subseteq B$ such that for all $s, t \in B'$ if $s \triangleleft t$, then $f(s) \preceq f(t)$. Clearly, $\text{RCA}_0 \vdash \text{bqo}(\text{set}) \rightarrow \text{bqo}$, and in Theorem 4.9 of [7], it is shown that the implication $\text{bqo} \rightarrow \text{bqo}(\text{set})$ is equivalent (over RCA_0) to ATR_0 . (ATR_0 is the subsystem formed by extending ACA_0 to allow definitions by the transfinite recursion of arithmetic predicates over well orders. It is significantly stronger than any of the systems considered in this paper.)

In the following lemma, we state a property of blocks which will enable us to prove that the base of any block exists in RCA_0 . In the statement of the lemma, we do not want to assume that the base of the block B exists. Therefore, the requirement that $s \subset \text{base}(B)$ is shorthand for the formal statement that for all $n < |s|$, there is a $u \in B$ such that $s(n) \in u$.

Lemma 5.4. (RCA_0) *If B is a block, then there is a string s and a number k such that $s \subset \text{base}(B)$, $\forall u \sqsubseteq s (u \notin B)$, and $\forall t \in B (s \sqsubset t \rightarrow |t| \leq |s| + k)$.*

Proof. For a contradiction, assume that there is no such string s . That is, if $s \subset \text{base}(B)$ and for all $r \sqsubseteq s$, $r \notin B$, then it is not the case that there is a k for which all $t \in B$ with $s \sqsubset t$ satisfy $|t| \leq |s| + k$. We define a set $Z \in [\text{base}(B)]^\omega$ such that Z has no initial segment B . This set contradicts the fact that B is a block and completes the proof.

We begin to define Z by applying our assumed hypothesis with s equal to the empty string. Our assumption tells us that there must be an $s_0 \in B$ with $|s_0| \geq 2$. Take the $<_{\mathbb{N}}$ -least such string and let $n_0 = s_0(0)$ be the least element of Z . We know that $\langle n_0 \rangle \notin B$ (since $\langle n_0 \rangle$ is a proper initial segment of s_0) so by our assumption there must be a string $s_1 \in B$ with $s_1(0) = n_0$ and $|s_1| \geq 3$. Take the $<_{\mathbb{N}}$ -least such string and let $n_1 = s_1(1)$ be the second smallest element of Z . Notice that neither $\langle n_0 \rangle$ nor $\langle n_0, n_1 \rangle$ is in B .

In general, assume that we have defined $n_0 < n_1 < \dots < n_k$ to be an initial segment of Z and that $\langle n_0, \dots, n_i \rangle \notin B$ for all $i \leq k$. By our assumption, there is an element $s_{k+1} \in B$ with $\langle n_0, \dots, n_k \rangle \sqsubset s_{k+1}$ and $|s_{k+1}| \geq k + 3$. Fix the $<_{\mathbb{N}}$ -least such string and let $n_{k+1} = s_{k+1}(k + 1)$ be the next smallest element in Z . Notice that the induction hypothesis is maintained so the recursion can continue.

Formally, we have defined a function f such that $f(k) = n_k$ and we let Z be the range of f . Z exists as a set since $k \in Z$ if and only if $\exists i \leq k (k = n_i)$. Also, by our construction, the nonempty initial segments of Z are precisely $\langle n_0, \dots, n_q \rangle$ for $q \in \mathbb{N}$ and none of these sequences is in B . This contradicts the fact that B is a block and finishes the proof. \square

Lemma 5.5. (RCA₀) *If $B \subset [\mathbb{N}]^{<\omega}$ is a block, then $\text{base}(B)$ is a set.*

Proof. It suffices for us to give a Π_1^0 definition of the base of B . By Lemma 5.4, we can fix a string s and a number k such that $s \subset \text{base}(B)$, $\forall u \sqsubseteq s (u \notin B)$ and for all $t \in B$ such that $s \sqsubset t$ we have $|t| \leq |s| + k$. Let n be the maximum value in s . We prove that $\text{base}(B)$ consists of a finite subset of $\{0, \dots, n\}$ plus the set X consisting of all $m > n$ for which

$$\forall m_1 < \dots < m_{k-1} ([m < m_1 \wedge (m_1, \dots, m_{k-1} \in \text{base}(B))] \rightarrow \exists i < k (s * \langle m, m_1, \dots, m_i \rangle \in B)).$$

Notice that the definition of X is Π_1^0 since the predicate “ $\in \text{base}(B)$ ” is Σ_1^0 . The fact that $X \subseteq \text{base}(B)$ is immediate from the definition of X . To see that $(\text{base}(B) \setminus \{0, \dots, n\}) \subseteq X$, fix $m_1, \dots, m_{k-1} \in \text{base}(B)$ such that $m < m_1 < \dots < m_{k-1}$ and let Y be an infinite set such that $s * \langle m, m_1, \dots, m_{k-1} \rangle \sqsubset Y$. Such a Y does exist because $\text{base}(B)$ is infinite and hence RCA₀ proves that it contains an infinite subset. Y must have an initial segment in B and we know that for all $r \sqsubseteq s$, $r \notin B$. Therefore, some extension of t of s with $|t| \leq |s| + k$ and $t \sqsubset Y$ must be in B . Therefore, there is an $i < k$ such that $s * \langle m, m_1, \dots, m_i \rangle \in B$.

In the previous paragraph, we defined X using a Π_1^0 formula and then showed that X is equal to the set of all $m > n$ such that $m \in \text{base}(B)$. Therefore, X also has a Σ_1^0 definition and hence it exists as a set. Furthermore, by bounded Σ_1^0 comprehension (which holds in RCA₀, see [11, Theorem II.3.9]), the set Z of all $m \leq n$ such that $m \in \text{base}(B)$ exists. Therefore, $\text{base}(B) = X \cup Z$ exists. \square

By Lemma 5.5, we can assume without loss of generality in RCA₀ that our blocks have base \mathbb{N} . This assumption is convenient since it allows us to avoid having to repeatedly state that we only work with strings whose elements come from $\text{base}(B)$.

Definition 5.6. (RCA_0) For strings s and t , we write $t \ll s$ if and only if $|t| = |s|$ and $\forall i < |t| (t(i) \leq s(i))$.

Given a block B with base \mathbb{N} , we define a tree $T(B)$ such that B is exactly the set of leaves of $T(B)$. We define a supertree $T^*(B)$ of $T(B)$ and let B^* be the set of leaves of $T^*(B)$. The set B^* will be a barrier. (A tree in this context means a nonempty subset of $[\mathbb{N}]^{<\omega}$ which is closed under initial segments and for which every string is strictly increasing.)

Definition 5.7. (RCA_0) For a block B with base \mathbb{N} , we define

$$\begin{aligned} T(B) &= \{s \mid \forall t \sqsubset s (t \notin B)\} \\ T^*(B) &= \{t \mid \exists s \ll t (s \in T(B))\} \\ B^* &= \{t \in T^*(B) \mid \forall u (t \sqsubset u \rightarrow u \notin T^*(B))\}. \end{aligned}$$

Notice that $T(B)$ and $T^*(B)$ can be defined in RCA_0 (because the quantifications $\forall t \sqsubset s$ and $\exists s \ll t$ are bounded, since $s \ll t$ implies $|s| = |t|$). At first, it appears that B^* requires ACA_0 to be defined since our trees are infinitely branching. We show below that this is not the case, but first we state the following obvious lemma.

Lemma 5.8. (RCA_0) $T(B) \subseteq T^*(B)$.

Proof. Obvious from the definition since $s \ll s$ for any string s . □

We prove the following characterization of B^* : for any $t \in T^*(B)$

$$t \in B^* \Leftrightarrow \forall s \ll t (s \in T(B) \rightarrow s \in B).$$

This characterization shows that RCA_0 is sufficient to define B^* . To see that (\Rightarrow) holds, suppose there is an $s \ll t$ such that $s \in T(B)$ and $s \notin B$. Then, for some $n > s(|s| - 1)$, the sequence $s * \langle n \rangle$ is in $T(B)$. Hence t has infinitely many extensions in $T^*(B)$ and therefore is not in B^* . To see that (\Leftarrow) holds, suppose that $t \notin B^*$ and fix $u \in T^*(B)$ such that $t \sqsubset u$. Because $u \in T^*(B)$, there is a v such that $v \ll u$ and $v \in T(B)$. Let s be v restricted to $|t|$. Notice that s is an element of $T(B)$, $s \ll t$ and $s \sqsubset v$. Therefore, s is not a leaf in $T(B)$ and hence is not an element of B . This establishes the equivalence.

Lemma 5.9. (WKL_0) If $T(B)$ has no infinite path, then $T^*(B)$ has no infinite path.

Proof. Suppose that $T^*(B)$ has an infinite path $X = \{x_0 < x_1 < \dots\}$. For each n , let $\sigma_n = \langle x_0, \dots, x_{n-1} \rangle$. Since $\sigma_n \in T^*(B)$, there must be an element $s \in T(B)$ such that $s \ll \sigma_n$. Let $S = \{s \in T(B) \mid s \ll \sigma_{|s|}\}$. S is a finitely branching subtree of $T(B)$ and the branching is bounded at level n by the function $f(n) = x_n$. Furthermore, since X is a path through $T^*(B)$, S is infinite. Applying Bounded König's Lemma (which is available in WKL_0 , see [11, Lemma IV.1.4]) to S , S must have an infinite path (which is of course also a path in $T(B)$). □

Restating Lemma 5.9 in purely computability theoretic language yields the following corollary.

Corollary 5.10. If $T(B)$ has no low path, then $T^*(B)$ has no computable path.

Proof. If X is a computable path through $T^*(B)$, then the bounding function f from the proof of Lemma 5.9 is computable. Hence, S is a computably bounded Π_1^0 class and by the Low Basis Theorem must have a low path. \square

Lemma 5.11. (WKL₀) *If B is a block with base \mathbb{N} , then B^* is a barrier.*

Proof. We check the properties required of a barrier. First, notice that because B is a block, every infinite set Y has some initial segment $s \sqsubset Y$ with $s \in B$. Furthermore, s cannot have any extension in $T(B)$. Therefore, $T(B)$ has no infinite path. By Lemma 5.9, $T^*(B)$ has no infinite path, which implies that every Y has a longest initial segment $t \sqsubset Y$ with $t \in T^*(B)$. Let $k = |t|$. We claim that $t \in B^*$. Towards a contradiction suppose that $t * n \in T^*(B)$ for some n . Then for some $s \ll t$ and $m \leq n$ we have $s * m \in T(B)$, so that $s \notin B$. Since B is a block with $\text{base}(B) = \mathbb{N}$, $s * i \in T(B)$ for every $i \in \mathbb{N}$ and, in particular, $s * Y(k) \in T(B)$. Since $s * Y(k) \ll t * Y(k)$ we have $t * Y(k) = Y|(k+1) \in T^*(B)$. Contradiction. Therefore, B^* contains initial segments of every set. Second, because $T^*(B)$ has no infinite paths, Lemma 5.8 shows that every $s \in B$ has some extension $t \sqsupseteq s$ such that t is a leaf in $T^*(B)$. Therefore, $\text{base}(B) \subseteq \text{base}(B^*)$, so $\text{base}(B^*) = \mathbb{N}$. Third, fix distinct leaves s and t in $T^*(B)$ and assume for a contradiction that $s \subset t$. Notice that $s \subset t$ implies that $|s| < |t|$ and if $|s| = n$ then $t|n \ll s$. Since $t \in T^*(B)$, there must be a $\sigma \in T(B)$ such that $|\sigma| = |t|$ and $\sigma \ll t$. Restricting to an initial segment gives $\sigma|n \ll t|n \ll s$. Hence, there is a string u properly extending s such that $|u| = |\sigma|$ and $\sigma \ll u$. Therefore, $u \in T^*(B)$. However, this contradicts the fact that s is a leaf on $T^*(B)$. \square

Theorem 5.12. (WKL₀) *A quasi-order (Q, \preceq) is a bqo if and only if for every block B with base \mathbb{N} and every function $f : B \rightarrow Q$, there exist $s, s' \in B$ such that $s \triangleleft s'$ and $f(s) \preceq f(s')$.*

Proof. The (\Leftarrow) direction is clear since every barrier is a block. For the (\Rightarrow) direction, let (Q, \preceq) be a bqo and fix both a block B and a function $f : B \rightarrow Q$. We have to find the corresponding s and s' . Let B^* be the barrier associated to B as above. Define $g : B^* \rightarrow B$ by setting $g(t)$ to be the unique $s \in B$ such that $s \sqsubseteq t$. (It is not hard to see that such s must exist and be unique since B is a block and $T(B) \subseteq T^*(B)$.) Define $f^* : B^* \rightarrow Q$ by composing f and g . Since Q is a bqo, there are $t, t' \in B^*$ such that $t \triangleleft t'$ and $f^*(t) \preceq f^*(t')$. Let $s = g(t)$ and $s' = g(t')$. It is clear that $s, s' \in B$ and $f(s) = f^*(t) \preceq f^*(t') = f(s')$. Because $t \triangleleft t'$, we can fix strings u and v such that $u = u(0) * v$, $t \sqsubseteq u$ and $t' \sqsubseteq v$. Since $s \sqsubseteq t$ and $s' \sqsubseteq t'$, the strings u and v also witness that $s \triangleleft s'$. \square

References

- [1] Peter Cholak, Carl Jockusch & Ted Slaman, The strength of Ramsey's Theorem for pairs, *Journal of Symbolic Logic*, 66(1) March 2001, 1-55. [1](#)
- [2] Rodney G. Downey, Computability theory and linear orderings, in Y.L. Ershov, S.S. Goncharov, A. Nerode & J.B. Remmel editors, *Handbook of Recursive Mathematics: Volume 2*, 823-976, North Holland, Amsterdam, 1998. [3](#), [3](#)

- [3] Rodney G. Downey, Denis R. Hirschfeldt, Steffen Lempp & Reed Solomon, Computability-theoretic and proof-theoretic aspects of partial and linear orderings, to appear in *Israel Journal of Mathematics*. 3
- [4] Eberhard Herrmann, Infinite chains and antichains in computable partial orderings, *Journal of Symbolic Logic*, 66(2) June 2001, 923-934. 3
- [5] Denis R. Hirschfeldt & Richard A. Shore, Some facts about linear orderings provable in RT_2^2 , to appear. 2, 3
- [6] Alberto Marcone, Foundations of bqo theory, *Transactions of the American Mathematical Society*, 345(2), October 1994, 641-660. 5
- [7] Alberto Marcone, Wqo and bqo theory in subsystems of second order arithmetic, to appear in S.G. Simpson editor, *Reverse mathematics 2001*. 1, 2, 4, 4, 5, 5
- [8] Eric C. Milner, Basic wqo- and bqo-theory, in Ivan Rival, editor, *Graphs and order*, 487-502, D. Reidel, Dordrecht, 1985. 1
- [9] Stephen G. Simpson, Bqo theory and Fraïssé's Conjecture, in R. Mansfield & G. Weitekamp, *Recursive aspects of descriptive set theory*, 124-138, Oxford University Press, Oxford 1985. 1
- [10] Stephen G. Simpson, Ordinal numbers and the Hilbert Basis Theorem, *Journal of Symbolic Logic*, 53(3) September 1988, 961-974. 2, 2
- [11] Stephen G. Simpson, *Subsystems of second order arithmetic*, Springer-Verlag, 1998. 1, 1, 5, 5