

### Multiple Choice

1.(5pts) Find the reduced echelon form of the matrix  $\begin{bmatrix} 3 & 2 & 4 \\ 2 & 2 & 2 \\ 3 & 4 & 2 \end{bmatrix}$ .

(a)  $\begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$

(b)  $\begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$

(c)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(d)  $\begin{bmatrix} 1 & -2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$

(e)  $\begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

2.(5pts) Determine by inspection which one of the following sets of vectors is linearly independent.

(a)  $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} \right\}$

(b)  $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ -4 \end{bmatrix} \right\}$

(c)  $\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix} \right\}$

(d)  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 0 \\ -2 \end{bmatrix} \right\}$

(e)  $\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \right\}$

3.(5pts) For which value of  $h$  is the vector  $\begin{bmatrix} 1 \\ h \\ 2 \end{bmatrix}$  in the span of the vectors  $\begin{bmatrix} 1 \\ -3 \\ 4 \end{bmatrix}$  and  $\begin{bmatrix} 2 \\ 3 \\ 2 \end{bmatrix}$ ?

- (a)  $h = 0$             (b)  $h = 1$             (c)  $h = 2$             (d)  $h = 3$             (e)  $h = 4$

4.(5pts) Let  $A$  be a  $3 \times 5$  matrix  $A$  and  $\mathbf{b}$  in  $\mathbb{R}^3$ . Which of the following statements about the matrix equation  $A\mathbf{x} = \mathbf{b}$  for  $\mathbf{x} \in \mathbb{R}^5$ , and the corresponding homogeneous equation  $A\mathbf{x} = \mathbf{0}$ , *could be true*?

- (a)  $A\mathbf{x} = \mathbf{b}$  has infinitely many solutions.  
(b)  $A\mathbf{x} = \mathbf{0}$  is inconsistent.  
(c)  $A\mathbf{x} = \mathbf{0}$  has exactly two solutions.  
(d)  $A\mathbf{x} = \mathbf{0}$  has a unique solution and  $A\mathbf{x} = \mathbf{b}$  has infinitely many solutions.  
(e)  $A\mathbf{x} = \mathbf{b}$  has a unique solution and  $A\mathbf{x} = \mathbf{0}$  has infinitely many solutions.

5.(5pts) Recall that an  $m \times n$  matrix has  $m$  rows and  $n$  columns. Let  $T : \mathbb{R}^6 \rightarrow \mathbb{R}^8$  be a linear transformation. What is the size of the standard matrix  $A$  for  $T$ ?

- (a)  $8 \times 6$
- (b)  $6 \times 6$
- (c)  $8 \times 8$
- (d)  $6 \times 8$
- (e) There is not enough information to determine the answer.

6.(5pts) Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation such that  $T \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right) = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$  and  $T \left( \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) = \begin{bmatrix} 7 \\ -2 \end{bmatrix}$ . What is the standard matrix  $A$  for  $T$ ?

- (a)  $A = \begin{bmatrix} 1 & 6 \\ 2 & -4 \end{bmatrix}$
- (b)  $A = \begin{bmatrix} 1 & 2 \\ 7 & -2 \end{bmatrix}$
- (c)  $A = \begin{bmatrix} 1 & 2 \\ 6 & -4 \end{bmatrix}$
- (d)  $A = \begin{bmatrix} 1 & 6 \\ 2 & 5 \end{bmatrix}$
- (e) Since we do not know  $T \left( \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right)$ , there is not enough information to determine the answer.

7.(5pts) Which of the following is a subspace of  $\mathbb{R}^3$ ?

(1) The set of all vectors,  $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$ , where  $a, b, c$  are positive.

(2) The set of all vectors,  $\begin{bmatrix} a \\ b \\ 1 \end{bmatrix}$ , where  $a, b$  are any numbers.

(3) The set of all vectors,  $\begin{bmatrix} a \\ 0 \\ c \end{bmatrix}$ , where  $a, c$  are any numbers.

(4) The set of all vectors,  $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$ , where  $a, b, c$  are integers.

(5) The set of all vectors,  $\begin{bmatrix} a \\ 2a \\ 3a \end{bmatrix}$ , where  $a$  is a real number.

- (a) (3) and (5)      (b) Only (3)      (c) Only (5)      (d) (4) and (5)      (e) (1) and (2)

8.(5pts) Which matrix below is invertible?

(a)  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 3 \\ 0 & 2 & 0 \end{bmatrix}$

(b)  $\begin{bmatrix} 2 & 3 & 0 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix}$

(c)  $\begin{bmatrix} 2 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$

(d)  $\begin{bmatrix} 1 & 1 & 4 & 2 \\ 8 & 0 & 6 & 3 \\ -1 & 2 & 7 & 1 \end{bmatrix}$

(e)  $\begin{bmatrix} 1 & 3 & -1 \\ -4 & -8 & 2 \\ 2 & 2 & 0 \end{bmatrix}$

9.(5pts) Let  $A$  be a  $3 \times 2$ ,  $B$  be a  $3 \times 5$  matrix, and  $C$  be a  $2 \times 5$  matrix. Which of the following expressions make sense?

(1)  $AB$

(2)  $BA$

(3)  $A + B$

(4)  $A^T B + C$

(5)  $A(B^T) + C$

(a) (4) only

(b) (2) and (3)

(c) (1) and (4)

(d) (5) only

(e) (1) and (5)

**Partial Credit**

**10.**(12pts) Express the solution set of the homogeneous linear system

$$\begin{aligned}x_1 + x_2 - x_3 + x_4 + 5x_5 &= 0 \\2x_1 + x_2 - 2x_3 + 4x_4 &= 0 \\3x_1 + 2x_2 - 3x_3 + 5x_4 + 6x_5 &= 0\end{aligned}$$

in *parametric vector form*.

11.(12pts) Find the inverse of  $A = \begin{bmatrix} 1 & 1 & 5 \\ 1 & 0 & -1 \\ 0 & 0 & 1 \end{bmatrix}$ .

12.(12pts) (a) Find a basis for the column space of  $A = \begin{bmatrix} 1 & 0 & -3 & 1 \\ -2 & -2 & 4 & 0 \\ -1 & -1 & -1 & 1 \end{bmatrix}$ .

(b) What is the rank of  $A$ ?

# Solutions

1.  $\begin{bmatrix} 3 & 2 & 4 \\ 2 & 2 & 2 \\ 3 & 4 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 \\ 2 & 2 & 2 \\ 3 & 4 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & -2 \\ 0 & 4 & -4 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -1 \\ 0 & 4 & -4 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$  where  $\sim$  denotes row equivalence.
2. First vector in (a) is non-zero and second is not a scalar multiple of it; so they are linearly independent.  
 (b) Not linearly independent;  $3\mathbf{v}_2 - 2\mathbf{v}_1 = \mathbf{0}$ .  
 (c) Four vectors in  $\mathbb{R}^3$  must be linearly dependent.  
 (d) Not linearly independent;  $\mathbf{v}_3 = 3\mathbf{v}_1 - 2\mathbf{v}_2$   
 (e) Not linearly independent; contains the zero vector.
3. The first vector is in the span of the other two exactly when the linear system with augmented matrix  $\begin{bmatrix} 1 & 2 & | & 1 \\ -3 & 3 & | & h \\ 4 & 2 & | & 2 \end{bmatrix}$  is consistent. Row reduce:  $\begin{bmatrix} 1 & 2 & | & 1 \\ -3 & 3 & | & h \\ 4 & 2 & | & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & | & 1 \\ 0 & 9 & | & h+3 \\ 0 & -6 & | & -2 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & | & 1 \\ 0 & 1 & | & (h+3)/9 \\ 0 & -6 & | & -2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & | & 1/3 - 2h/9 \\ 0 & 1 & | & h/9 + 1/3 \\ 0 & 0 & | & 2h/3 \end{bmatrix}$ . The system is consistent exactly when  $2h/3 = 0$ ; that is, when  $h = 0$ .
4.  $A\mathbf{x} = \mathbf{b}$  has infinitely many solutions if it is consistent and the reduced echelon form of  $A$  has at least one free variable, so (a) could be true. A linear system can have (only) 0, 1 or infinitely many solutions and a homogeneous system  $A\mathbf{x} = \mathbf{0}$  is always consistent (with solution  $\mathbf{x} = \mathbf{0}$ ) so (c) and (b) are false. If  $A\mathbf{x} = \mathbf{b}$  is consistent, all its solutions are obtained by adding a solutions of the homogeneous system to a particular solution of  $A\mathbf{x} = \mathbf{b}$ , so  $A\mathbf{x} = \mathbf{b}$  and  $A\mathbf{x} = \mathbf{0}$  have exactly the same number of solutions in that case; (d) and (e) are therefore also false.
5. Since  $T$  sends vectors in  $\mathbb{R}^6$  to vectors in  $\mathbb{R}^8$ , the standard matrix  $A$  must be  $8 \times 6$ .
6. The standard matrix for the linear transformation is the matrix  $[T(\mathbf{e}_1) \ T(\mathbf{e}_2)]$ , where  $\mathbf{e}_1$  is the first column of the  $2 \times 2$  identity matrix and  $\mathbf{e}_2$  is the second column of the  $2 \times 2$  identity matrix. By linearity, we have that  $T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) - T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 6 \\ -4 \end{bmatrix}$ , thus the standard matrix  $A = \begin{bmatrix} 1 & 6 \\ 2 & -4 \end{bmatrix}$ .
7. (1) is not a subspace since it is not closed under multiplication by scalars. For instance,  $-2 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \\ -2 \end{bmatrix}$  is not in the set. (2) is not a subspace since the zero vector,  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ , is not in the set. (4) is not a subspace since it is not closed under multiplication by scalars. For instance,  $.5 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} .5 \\ .5 \\ .5 \end{bmatrix}$  is not in the set. (3) and (5) satisfy all of the properties of a subspace.



8.  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 3 \\ 0 & 2 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 2 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  so all columns are pivots so invertible.

$\begin{bmatrix} 2 & 3 & 0 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix}$  columns 1 and 2 are dependent so not invertible.

$\begin{bmatrix} 2 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 1 & 4 & 2 \\ 8 & 0 & 6 & 3 \\ -1 & 2 & 7 & 1 \end{bmatrix}$  are not square matrices and hence not invertible.

$\begin{bmatrix} 1 & 3 & -1 \\ -4 & -8 & 2 \\ 2 & 2 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 3 & -1 \\ 0 & 4 & -2 \\ 0 & -4 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 3 & -1 \\ 0 & 4 & -2 \\ 0 & 0 & 0 \end{bmatrix}$  so not invertible.

9.  $AB$  does not make sense since the number of columns of  $A$  is not equal to the number of rows of  $B$ . For the same reason,  $BA$  also does not make sense.  $A+B$  does not make sense because we cannot sum matrices of different sizes.  $A^T B$  is a  $2 \times 5$  matrix, which can be added to  $C$ , since  $C$  is a  $2 \times 5$  matrix. So  $A^T B + C$  makes sense.  $A(B^T) + C$  does not make sense since the number of columns of  $A$  is not the same as the number of rows of  $B^T$ . So  $A(B^T) + C$  does not make sense.

10. Row reduce the coefficient matrix:  $\begin{bmatrix} 1 & 1 & -1 & 1 & 5 \\ 2 & 1 & -2 & 4 & 0 \\ 3 & 2 & -3 & 5 & 6 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & -1 & 1 & 5 \\ 0 & -1 & 0 & 2 & -10 \\ 0 & -1 & 0 & 2 & -9 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & -1 & 1 & 5 \\ 0 & 1 & 0 & -2 & 10 \\ 0 & -1 & 0 & 2 & -9 \end{bmatrix} \sim$

$\begin{bmatrix} 1 & 1 & -1 & 1 & 5 \\ 0 & 1 & 0 & -2 & 10 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & -1 & 1 & 0 \\ 0 & 1 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -1 & 3 & 0 \\ 0 & 1 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$ . The system is equivalent

to

$$\begin{aligned} x_1 - x_3 + 3x_4 &= 0 \\ x_2 - 2x_4 &= 0 \\ x_5 &= 0 \end{aligned}$$

The bound variables are  $x_1, x_2$  and  $x_5$  (corresponding to pivot columns) and the free variables  $x_3, x_4$  can take arbitrary values. Rewriting with free variables on the right,

$$\begin{aligned} x_1 &= x_3 - 3x_4 \\ x_2 &= 2x_4 \\ x_3 &= x_3 \\ x_4 &= x_4 \\ x_5 &= 0 \end{aligned}$$

(we include the equation  $x_i = x_i$ , for  $i = 3$  or  $4$ , to indicate that the free variable  $x_i$  can take arbitrary values). In parametric form

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_3 \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -3 \\ 2 \\ 0 \\ 1 \\ 0 \end{bmatrix}$$

or writing  $x_3 = r, x_4 = s$ ,

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = r \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -3 \\ 2 \\ 0 \\ 1 \\ 0 \end{bmatrix}$$

11. Row reduce the augmented matrix  $[A \ I]$ :  $\begin{bmatrix} 1 & 1 & 5 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 5 & 1 & 0 & 0 \\ 0 & -1 & -6 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} \sim$   
 $\begin{bmatrix} 1 & 1 & 0 & 1 & 0 & -5 \\ 0 & -1 & 0 & -1 & 1 & 6 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & -1 & 0 & -1 & 1 & 6 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & -1 & -6 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix}$ . So  $A^{-1} =$   
 $\begin{bmatrix} 0 & 1 & 1 \\ 1 & -1 & -6 \\ 0 & 0 & 1 \end{bmatrix}$ .

12. Row-reduce  $\begin{bmatrix} 1 & 0 & -3 & 1 \\ -2 & -2 & 4 & 0 \\ -1 & -1 & -1 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -3 & 1 \\ 0 & -2 & -2 & 2 \\ -1 & -1 & -1 & 1 \end{bmatrix} \sim$   
 $\begin{bmatrix} 1 & 0 & -3 & 1 \\ 0 & -2 & -2 & 2 \\ 0 & -1 & -4 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -3 & 1 \\ 0 & 1 & 1 & -1 \\ 0 & -1 & -4 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -3 & 1 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & -3 & 1 \end{bmatrix}$ . From this we see that the pivot  
columns of  $A$  are the first three columns and thus a basis for  $\text{Col}A$  is  
 $\left\{ \begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ -1 \end{bmatrix}, \begin{bmatrix} -3 \\ 4 \\ -1 \end{bmatrix} \right\}$ . Alternatively, since  $\text{Col}A = \mathbb{R}^3$ , any basis of  $\mathbb{R}^3$  would do. The rank of  
 $A$  is the dimension of  $\text{Col}A$ , which is 3.