

Continuous uniform probability model

Math 30530, Fall 2013

August 31, 2013

Continuous uniform probability model

- 1 **Experiment:** Have a region R in the plane, and select a point from it at random (e.g. by throwing a dart at the region), with no bias towards or against any particular part of the region

Continuous uniform probability model

- 1 **Experiment:** Have a region R in the plane, and select a point from it at random (e.g. by throwing a dart at the region), with no bias towards or against any particular part of the region
- 2 **Sample space:** $\Omega = \{(x, y) \mid (x, y) \text{ is a point in region } R\}$

Continuous uniform probability model

- 1 **Experiment:** Have a region R in the plane, and select a point from it at random (e.g. by throwing a dart at the region), with no bias towards or against any particular part of the region
- 2 **Sample space:** $\Omega = \{(x, y) \mid (x, y) \text{ is a point in region } R\}$
- 3 **Probabilities:** For any event E ,

$$\Pr(E) = \frac{\text{Area}(E)}{\text{Area}(R)}$$

i.e. the probability of the dart landing inside E is proportional to the area of E

Continuous uniform probability model

- 1 **Experiment:** Have a region R in the plane, and select a point from it at random (e.g. by throwing a dart at the region), with no bias towards or against any particular part of the region
- 2 **Sample space:** $\Omega = \{(x, y) | (x, y) \text{ is a point in region } R\}$
- 3 **Probabilities:** For any event E ,

$$\Pr(E) = \frac{\text{Area}(E)}{\text{Area}(R)}$$

i.e. the probability of the dart landing inside E is proportional to the area of E

- 4 **Consequence:** If p is a particular point in R , then (homework!)

$$\Pr(p) = 0$$

This is an example of an event that could occur, but has probability 0 of occurring