


1. Dimensional analysis in cooking a turkey: If you've ever looked at the cooking timetables for large things like turkeys or roasts, you may have noticed that they are pretty complex: so much per pound if the weight is in one range, and different length in another range, etc. In general, the cooking time per pound goes down the larger the bird. If we assume that 1) all turkeys are geometrically similar, 2) the physical properties such as thermal diffusivity, density, etc., are all constant, and 3) the cooking time depends only on the mass, the density, and the thermal diffusivity (for idealized turkeys, anyway), use dimensional analysis to show how cooking time varies with the mass of the bird. Graphically compare your model (determining the constant empirically) to the values obtained from any source you wish. (If you don't have a cookbook, a web search on "turkey cooking time" is quite useful!)

2. Consider the damped pendulum depicted below. The pendulum consists of a ball of mass  $m$  and radius  $a$  suspended by a fine wire of length  $L$  below a pivot. The ball is moving through a fluid of viscosity  $\mu$  (the fluid is assumed massless for convenience) which retards its motion. It is assumed that the ball moves sufficiently slowly that Stokes Law applies. Under these conditions, the equation describing the motion of the sphere is given by:



$$ML \frac{d^2 \theta}{dt^2} = -Mg \sin(\theta) - 6\pi \mu a L \frac{d\theta}{dt}$$


$$\theta_{t=0} = \theta_0 ; \quad \frac{d\theta}{dt} \Big|_{t=0} = 0$$

a. By choosing an appropriate reference time scale, render the equation and initial condition dimensionless. Show that the dimensionless solution depends on only two dimensionless groups. What are the physical meanings of these parameters?

b. You are assigned the problem of adjusting the fluid viscosity so as to bring the sphere to rest as rapidly as possible. Recognizing that if  $\mu = 0$  the pendulum will oscillate forever, and if  $\mu$  is very large the sphere will take a long time to get to the equilibrium ( $\theta = 0$ ) position, estimate the correct value of the viscosity for critical damping. Estimate from dimensional analysis how long it will take the sphere to approach to within, say, 25% of the equilibrium position.

c. Given that  $\theta_0$  is very small, we may make the approximation  $\sin(\theta) \approx \theta$ . Solve the resulting differential equation, and compare the exact solution to that estimated by dimensional analysis above.

3. A disk of radius  $R$  is falling face on toward a plane with velocity  $V$  through a fluid of viscosity  $\mu$  as depicted below. Using the lubrication approximations (e.g., assuming that the flow in the narrow gap of size  $h$  between the disk and the plane is a quasi-parallel channel flow in the radial direction, and that inertial effects are negligible), calculate the force on the disk due to the squeezing flow out of the gap.



4. By taking the divergence of the equations of motion and applying the equation of continuity, prove that  $\nabla^2 p = 0$  for an incompressible fluid undergoing flow at zero Reynolds number. Use index notation only, and note that the  $\nabla$  and  $\nabla^2$  operators commute.