

Real solutions to Alt-Burmester problems

Jonathan D. Hauenstein

The family of Alt-Burmester problems [3] consist of computing four-bar linkages which attain M poses and pass through N points, e.g., Figure 1 is a solution for $(M, N) = (0, 9)$ which is the classical Alt problem [1]. The classical Burmester problem [4] corresponds with $(M, N) = (5, 0)$.




FIGURE 1. Four-bar linkage whose coupler curve passes through 9 given points

Since the initial pose can be trivially set, the classical Alt problem with $(M, N) = (0, 9)$ is the same as $(M, N) = (1, 8)$. The following table from [3] lists the Alt-Burmester problems which generically have finitely many solutions along to the corresponding polynomial system.

(M, N)	generic # of solutions
$(5, 0)$	$16 = 4 \cdot 4$
$(4, 2)$	60
$(3, 4)$	402
$(2, 6)$	2224
$(1, 8) = (0, 9)$	$8652 = 1442 \cdot 6$

This table highlights that the solutions to both the classical Burmester problem $(M, N) = (5, 0)$ and Alt problem $(M, N) = (1, 8)$ are actually in natural families of size 4 and 6, respectively.

Problem 1. *What is the maximum number of real solutions to each generically zero-dimensional Alt-Burmester problem?*

The classical Burmester problem can have all solutions real, while the other cases are open.

REFERENCES

- [1] H. Alt. Über die Erzeugung gegebener ebener Kurven mit Hilfe des Gelenkvierecks, *ZAMM*, 3(1), 13–19, 1923.
- [2] D.J. Bates, W. Decker, J.D. Hauenstein, C. Peterson, G. Pfister, F.-O. Schreyer, A.J. Sommese, and C.W. Wampler. Comparison of probabilistic algorithms for analyzing the components of an affine algebraic variety. *Appl. Math. Comput.*, 231, 619–633, 2014.
- [3] D.A. Brake, J.D. Hauenstein, A.P. Murray, D.H. Myszka, and C.W. Wampler. The complete solution of Alt-Burmester synthesis problems for four-bar linkages. *J. Mech. Robot.*, 8(4), 041018, 2016.
- [4] L. Burmester, *Lehrbuch der Kinematik*, Verlag Von Arthur Felix, Leipzig, Germany, 1886.
- [5] J.D. Hauenstein and J.I. Rodriguez. Multiprojective witness sets and a trace test. Available at www.nd.edu/~jhauenst/preprints/hrMultiHom.pdf.
- [6] J.D. Hauenstein, A.J. Sommese, and C.W. Wampler. Regeneration homotopies for solving systems of polynomials. *Math. Comput.*, 80, 345–377, 2011.
- [7] C.W. Wampler, A.P. Morgan, and A.J. Sommese. Complete solution of the nine-point path synthesis problem for four-bar linkages. *ASME J. Mech. Des.*, 114(1), 153–159, 1992.