

# Equivariant Witness Sets

Jonathan D. Hauenstein

Many varieties are naturally invariant under a finite group action. For example,

$$(1) \quad f = q_{400}q_{040}q_{004} - q_{400}q_{022}^2 - q_{040}q_{202}^2 - q_{004}q_{220}^2 - 2q_{220}q_{202}q_{022}$$

from [4] defines a hypersurface  $\mathcal{H} \subset \mathbb{P}^5$  that is invariant under the group  $G = \langle \sigma \rangle$  where

$$\sigma(q_{ijk}) = q_{kij}.$$

This hypersurface  $\mathcal{H}$  is related to the Luroth hypersurface [5] which also invariant under the same group  $G$ . Other examples include secant varieties of Segre varieties.

**Problem 1.** *Suppose that  $V$  is an irreducible variety and  $G$  is a finite group that acts invariantly on  $V$ , i.e.,  $g(v) \in V$  for every  $v \in V$  and  $g \in G$ . Can one utilize  $G$  to create an “equivariant” witness set for  $V$ . Such a witness set should be able to be used, e.g., for testing membership in  $V$  by tracking less paths than a standard witness set for  $V$ .*

Some special cases were used in [1, 2, 3].

Returning to  $f$  as in (1) with  $[q_{400}, q_{040}, q_{004}, q_{220}, q_{022}, q_{202}] \in \mathbb{P}^5$ , the system

$$\begin{bmatrix} f \\ (q_{400} + q_{040} + q_{004}) + 3(q_{220} + q_{202} + q_{022}) \\ 2q_{400} + 3q_{040} - q_{004} - 2q_{220} + 5q_{022} - 3q_{202} \\ 2q_{040} + 3q_{004} - q_{400} - 2q_{022} + 5q_{202} - 3q_{220} \\ 2q_{004} + 3q_{400} - q_{040} - 2q_{202} + 5q_{220} - 3q_{022} \end{bmatrix} = 0$$

has 3 solutions which are invariant under  $G$ , approximately

$$\begin{aligned} &[1.0000, -0.0964, -0.9036, -0.2268, -0.2985, 0.5253], \\ &[-0.9036, 1.0000, -0.0964, 0.5253, -0.2268, -0.2985], \\ &[-0.0964, -0.9036, 1.0000, -0.2985, 0.5253, -0.2268]. \end{aligned}$$

Hence, due to this invariance, only one point is needed rather than  $3 = \deg f$ .

Problem 1 aims to generalize this construction. Remark 4.1 of [2] shows that some caution is needed in this generalization.

## REFERENCES

- [1] C. Amendola and J.I. Rodriguez. Solving parameterized polynomial system with decomposable projections. Preprint available at [arxiv.org/pdf/1612.08807.pdf](https://arxiv.org/pdf/1612.08807.pdf).
- [2] J.D. Hauenstein, L. Oeding, G. Ottaviani, and A.J. Sommese. Homotopy techniques for tensor decomposition and perfect identifiability. To appear in *J. Reine Angew. Math.*
- [3] J.D. Hauenstein and J.I. Rodriguez. Multiprojective witness sets and a trace test. Preprint available at [www.nd.edu/~jhauenst/preprints/hrMultiHom.pdf](http://www.nd.edu/~jhauenst/preprints/hrMultiHom.pdf).
- [4] J.D. Hauenstein and F. Sottile. Newton polytopes and witness sets. *Math. Comp. Sci.*, 8(2), 235–251, 2014.
- [5] J. L roth. Einige Eigenschaften einer gewissen Gattung von Curven vierten Ordnung. *Math. Ann* 1, 37–53, 1869.