

Math 30710

Exam 1

March 1, 2023

Name _____

This is a 50-minute exam. Books and notes are not allowed. Make sure that your work is legible, and make sure that it is clearly marked where your answers are. There's a blank page at the end in case you need it either to continue an answer or for scrap paper, but mark your answers clearly!!

Show all work!

If a problem calls for a proof or explanation, you will not get full credit for a correct answer if you don't supply the proof or explanation. If you have some ideas for solving a problem but can't figure out how to finish it, be sure to show me what you do know!! If something isn't clear, ask me! If you need more space, there is a blank sheet at the back.

The Honor Code is in effect for this examination, including keeping your own exam under cover. Good luck!!

1. Short answer. For each of the following statements, if it's true then give a proof. If it's not true then explain why not (possibly by giving a counterexample, unless an explanation is more appropriate).

(a) (7 points) If G is a non-abelian group then every non-trivial subgroup of G is also non-abelian.

(b) (7 points) If G is a multiplicative group then the identity element for G is the only element equal to its own square.

(c) (7 points) If G is a group then the order of the identity element, e , is not well-defined because $e^k = e$ for all integers k .

(d) (7 points) D_4 is the smallest non-trivial subgroup of S_4 .

2. Let $G = (\mathbb{Z}_{10}, +)$.

(a) (5 points) Write down the subgroup diagram for G .

(b) (7 points) Write the Cayley digraph for G using the generators 2 and 5. Be sure to indicate very carefully what each arrow represents. (Note that $5 + 5 = 0$ in \mathbb{Z}_{10} .)

3. Let $G = (\mathbb{C}^*, \cdot)$ be the multiplicative group of non-zero complex numbers.

(a) (7 points) Let H be a subgroup of G and assume that H is finite. Prove that every element of H has to lie on the unit circle.

(b) (8 points) If $z = e^{i\theta}$ is a point on the unit circle, prove that the order of z under multiplication is finite if and only if some integer multiple of θ is a multiple of 2π , i.e. there is some positive integer n so that $n\theta$ is either 2π , 4π , etc.

4. Let $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 \\ 3 & 11 & 2 & 7 & 10 & 9 & 8 & 1 & 6 & 4 & 12 & 5 \end{pmatrix}$.

(a) (5 points) Write σ as a product of disjoint cycles.

(b) (5 points) Write σ as a product of transpositions and say if it is even or odd.

5. Let

$$\sigma_1 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 1 & 4 & 3 & 6 & 5 \end{pmatrix} \qquad \sigma_2 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 1 & 6 & 5 & 4 & 3 \end{pmatrix}$$

(a) (5 points) Find $\sigma_1\sigma_2$ and write your answer here:

$$\sigma_1\sigma_2 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ & & & & & \end{pmatrix}$$


You do not have to explain how you got your answer.

(b) (5 points) Find σ_1^{-1} and write your answer here:

$$\sigma_1^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ & & & & & \end{pmatrix}$$

You do not have to explain how you got your answer.

(c) (5 points) Suppose we number the corners of a regular hexagon in the usual (consecutive) way numbered clockwise:



Of the given permutations σ_1 and σ_2 in this problem, one is in D_6 and one is not. Which is in D_6 ? Describe it in terms of rotations and/or reflections. (It would probably help if you use the above picture, or one like it, in your answer.)

6. (10 points) Let $G = \{e, a, b, c, d, f\}$ be a group and let $*$ be the binary operation on G . Assume e is the identity element for G . **Assume further that G is abelian.** Suppose the following is the group table for G .

$*$	e	a	b	c	d	f
e						
a		f	c	d	b	
b			f	e	a	
c				a	f	
d					e	
f						

Fill in the blank spaces so that you complete the group table. You'll have to use what you know about the group table of an abelian group. You do not need to explain your answer in this problem.

7. (10 points) Prove that if $\phi : G \rightarrow G'$ is an isomorphism, and if G is abelian, then G' is abelian (i.e. the property of a group being abelian is preserved under isomorphism).

(Blank page.)