

Many-Body Methods Applied to Parity Nonconserving Transitions in Atoms: The Weak Charge and Anapole Moment of ^{133}Cs

Walter Johnson

University of Notre Dame, Notre Dame, IN (USA)

<http://www.nd.edu/~johnson>

I. Atomic Theory Considerations for PNC Calculations

- a) Correlation effects in heavy alkali-metals atoms are large.
- b) Relativistic effects in heavy alkali-metal atoms are important.
- c) Precise *ab-initio* theory is essential in PNC studies.

Outline

- (Quick) Review of Relativistic MBPT
- 2nd- and 3rd-Order Energies
- 2nd- and 3rd-Order Matrix Elements
- All-order **SDCC** and **SD** Equations.
- SD Energies & Fine-Structure Intervals.
- SD Dipole Matrix Elements & Hyperfine Constants.
- Polarizabilities.

Dirac Hamiltonian

One-electron atoms:

$$h_0 = c \boldsymbol{\alpha} \cdot \boldsymbol{p} + \beta mc^2 + U(r) + V_{\text{nuc}} .$$

$$h_0 \phi_i = \epsilon_i \phi_i$$

The spectrum of h_0 consists of electron scattering states ($\epsilon_i > mc^2$), electron bound states ($mc^2 > \epsilon_i > 0$) and negative energy (positron) states ($-mc^2 > \epsilon_i$).

Many-electron atoms:

The *no-pair* Hamiltonian

is a many-electron generalization obtained from the field-theoretic Hamiltonian of QED by performing a contact transformation to eliminate the electron - photon interaction to order e^2 .

The *no-pair* Hamiltonian¹ can be written

$$\begin{aligned} H &= H_0 + V \\ H_0 &= \sum_i \epsilon_i a_i^\dagger a_i \\ V &= \frac{1}{2} \sum_{ijkl} v_{ijkl} a_i^\dagger a_j^\dagger a_l a_k - \sum_{ij} U_{ij} a_i^\dagger a_j, \end{aligned}$$

where,

1. sums are restricted to electron states only,
2. v_{ijkl} is a two-particle matrix element of Coulomb + Breit interactions,
3. U_{ij} compensates for including $U(r)$ in h_0 .

¹ BROWN, G.E. & RAVENHALL, D.G. 1951 *Proc. R. Soc. London, Ser. A* **208**, 552-559.

MBPT (One Valence Electron)

Choose $U = V_{\text{DHF}}$, then $\Psi^{(0)} = \Psi_{\text{DHF}}$

$$\Psi = \Psi^{(0)} + \Psi^{(1)} + \dots$$


$$E = E^{(0)} + E^{(1)} + \dots$$

$$E_v^{(0)} = \epsilon_v$$

$$E_v^{(2)} = - \sum_{bm n} \frac{v_{m n v b} \tilde{v}_{v b m n}}{\epsilon_m + \epsilon_n - \epsilon_v - \epsilon_b} + \sum_{a b n} \frac{v_{v n a b} \tilde{v}_{a b v n}}{\epsilon_v + \epsilon_n - \epsilon_a - \epsilon_b}$$

$$E_v^{(3)} = \sum_{b m n r s} \frac{\tilde{v}_{v b m n} v_{m n r s} v_{r s v b}}{(\epsilon_v + \epsilon_b - \epsilon_m - \epsilon_n)(\epsilon_v + \epsilon_b - \epsilon_r - \epsilon_s)}$$

+ ... (11 more lines)



Example


2nd- & 3rd-Order Energies

Valence State Energies for Alkali-Metal Atoms (a.u.).


Term	Li 2s	Na 3s	K 4s	Rb 5s	Cs 6s	Fr 7s
DHF	-0.19632	-0.18203	-0.14749	-0.13929	-0.12737	-0.13108
$E^{(2)}$	-0.00165	-0.00593	-0.01246	-0.01507	-0.01782	-0.02174
$E^{(3)}$	-0.00012	-0.00032	0.00205	0.00309	0.00570	0.00770
Sum	-0.19809	-0.18829	-0.15790	-0.15127	-0.13949	-0.14512
NIST	-0.19814	-0.18886	-0.15952	-0.15351	-0.14310	-0.14967
$\Delta\%$	0.03	0.3	1.0	1.5	2.8	3.0

Lesson: It is necessary to go beyond 3rd-order MBPT to obtain energies accurate to better than 1% for heavy alkali-metal atoms.

1st-, 2nd-, & 3rd-order Matrix Elements



Dipole Matrix Elements for Alkali-Metal Atoms



Lesson: Third-order transition moments for heavy alkali-metal atoms are accurate to about 1%

Dipole Matrix Elements (Gauge)

Term	L	V	L	V
Cs	$6s_{1/2} - 6p_{1/2}$		$6s_{1/2} - 6p_{3/2}$	
1st	5.2777	5.0371	7.4265	7.0662
2nd	4.9747	4.9747	7.0137	7.0137
3rd	4.5402	4.5400	6.3892	6.3891
Fr	$7s_{1/2} - 7p_{1/2}$		$7s_{1/2} - 7p_{3/2}$	
1st	5.1438	4.8402	7.0903	6.6424
2nd	4.7741	4.7741	6.6268	6.6268
3rd	4.3236	4.3234	5.9450	5.9448

Lesson: “Dressed” third-order dipole matrix element are independent of gauge.

Hyperfine Matrix Elements

Cesium Hyperfine Constants (MHz)

Term	$6s_{1/2}$	$6p_{1/2}$	$6p_{3/2}$
A (1)	1423.77	160.92	23.92
A (2)	292.18	40.67	18.85
BO(3)	718.60	84.73	16.12
SR(3)	-7.97	5.43	-7.49
No(3)	-22.80	-1.20	-0.23
A (3)	687.83	88.96	8.40
Sum	2403.78	290.55	51.17
Expt.	2298.16	291.89	50.28
$\Delta\%$	4.6%	0.5%	1.8%

Lesson: (once again) One must go beyond 3rd-order to obtain matrix elements accurate to better than 1% for heavy alkali-metal atoms.

Coupled-Cluster Expansion

$$\begin{aligned}\Psi_v = & \exp \{1 + S_1 + S_2 + \dots + S_N\} \Psi_{\text{DHF}} \\ \approx & \left\{ 1 + S_1 + S_2 + \frac{1}{2}S_1^2 + S_1S_2 + \frac{1}{2}S_2^2 + \frac{1}{6}S_1^3 + \frac{1}{2}S_1^2S_2 + \frac{1}{24}S_1^4 \right\} \Psi_{\text{DHF}}\end{aligned}$$

$$\begin{aligned}S_1 = & S_{1c} + S_{1v} = \sum_{ma} \rho_{ma} a_m^\dagger a_a + \sum_{m \neq v} \rho_{mv} a_m^\dagger a_v \\ S_2 = & S_{2c} + S_{2v} = \frac{1}{2} \sum_{mnab} \rho_{mnab} a_m^\dagger a_n^\dagger a_b a_a + \sum_{mna} \rho_{mnvb} a_m^\dagger a_n^\dagger a_b a_v\end{aligned}$$

Nonlinear terms account for about 10% of correlation energy in Cs and about 4% in Na.
These nonlinear corrections are partially canceled by triples!

All-Order SD Approximation

$$\Psi_v = \Psi_{\text{DHF}} + \delta\Psi$$

$$\begin{aligned} \delta\Psi = & \left\{ \sum_{am} \rho_{ma} a_m^\dagger a_a + \frac{1}{2} \sum_{abmn} \rho_{mnab} a_m^\dagger a_n^\dagger a_b a_a \right. \\ & \left. + \sum_{m \neq v} \rho_{mv} a_m^\dagger a_v + \sum_{bm} \rho_{mnbv} a_m^\dagger a_n^\dagger a_b a_v \right\} \Psi_{\text{DHF}} \end{aligned}$$

$$E_c = E_c^{\text{DHF}} + \delta E_c$$


$$E_v = E_v^{\text{DHF}} + \delta E_v$$

Later, we will discuss triples!



Core Excitation Equations

$$\begin{aligned}(\epsilon_a - \epsilon_m)\rho_{ma} &= \sum_{bn} \tilde{v}_{mban}\rho_{nb} + \sum_{bnr} v_{mbnr}\tilde{\rho}_{nrab} - \sum_{bcn} v_{bcn}\tilde{\rho}_{mnbc} \\(\epsilon_a + \epsilon_b - \epsilon_m - \epsilon_n)\rho_{mnab} &= v_{mnab} + \sum_{cd} v_{cdab}\rho_{mncd} + \sum_{rs} v_{mnrs}\rho_{rsab} \\&+ \left[\sum_r v_{mnrb}\rho_{ra} - \sum_c v_{cnab}\rho_{mc} + \sum_{rc} \tilde{v}_{cnrb}\tilde{\rho}_{mrac} \right] + [\quad a \leftrightarrow b \quad m \leftrightarrow n \quad] \\ \delta E_c &= \frac{1}{2} \sum_{abmn} v_{abmn}\tilde{\rho}_{mnab}\end{aligned}$$

15,000,000 ρ_{mnab} coefficients for Cs ($\ell = 6$).



+ exchange terms



+ exchange terms

Brueckner-Goldstone Diagrams for the core SD equations.

Valence Equations

$$\begin{aligned}
 (\epsilon_v - \epsilon_m + \delta E_v) \rho_{mv} &= \sum_{bn} \tilde{v}_{mbvn} \rho_{nb} + \sum_{bnr} v_{mbnr} \tilde{\rho}_{nrvb} - \sum_{bcn} v_{bcvn} \tilde{\rho}_{mnbc} \\
 (\epsilon_v + \epsilon_b - \epsilon_m - \epsilon_n + \delta E_v) \rho_{mnvb} &= v_{mnvb} + \sum_{cd} v_{cdvb} \rho_{mncd} + \sum_{rs} v_{mnrs} \rho_{rsvb} \\
 &\quad + \left[\sum_r v_{mnrb} \rho_{rv} - \sum_c v_{cnvb} \rho_{mc} + \sum_{rc} \tilde{v}_{cnrb} \tilde{\rho}_{mrvc} \right] + \left[\begin{array}{c} v \leftrightarrow b \\ m \leftrightarrow n \end{array} \right] \\
 \delta E_v &= \sum_{ma} \tilde{v}_{vavm} \rho_{ma} + \sum_{mab} v_{abvm} \tilde{\rho}_{mvab} + \sum_{mna} v_{vbmn} \tilde{\rho}_{mnvb}
 \end{aligned}$$

1,000,000 ρ_{mnvb} coefficients for each state (Cs)

Perturbation Expansion of SD Energy

- δE_c agrees with MBPT through third-order.
- $\delta E_v^{(3)}$ disagrees with $E_v^{(3)}$ from MBPT.

Add a limited class of triple excitations to the SD wave function

$$\frac{1}{6} \sum_{abmn} \rho_{mnrvab} a_m^\dagger a_n^\dagger a_r^\dagger a_v a_b a_a \Psi_{\text{DHF}}$$

$$\delta E_{\text{extra}} = \frac{1}{2} \sum_{mnab} \tilde{v}_{abmn} \rho_{mnvvab}$$

- $\delta E_v + \delta E_{\text{extra}}$ includes the entire third-order MBPT valence correlation energy

Lithium Energy Levels

Summary for lithium $2s$ and $2p$ states (a.u.)


Term	$2s_{1/2}$	$2p_{1/2}$	$2p_{3/2}$
DHF	-0.196320	-0.128638	-0.128636
δE_{SD}	-0.001853	-0.001606	-0.001606
δE_{extra}	0.000011	0.000010	0.000010
Breit + RM + MP	0.000019	0.000009	0.000008
Total	-0.198143	-0.130226	-0.130225
Expt.	-0.198142	-0.130236	-0.130235

Sodium Energy Levels

Summary for sodium $3s$ and $3p$ states (cm^{-1})

Term	$3s$	$3p_{1/2}$	$3p_{3/2}$
DHF	-39951.6	-24030.4	-24014.1
δE_{SD}	-1488.8	-463.9	-461.6
δE_{extra}	-9.2	-1.5	-1.6
Breit	1.2	1.4	0.1
RM+MP	1.0	0.5	0.5
Theory	-41447.3	-24493.9	-24476.7
Expt.	-41449.4	-24493.3	-24476.1

Energy Comparisons for Alkali-Metal Atoms



SD Energies for Heavy Atoms

Energies for cesium and francium cm^{-1}

Cs	$6s$	$7s$	$8s$	$9s$
Theory	31262	12801	7060	4479
Expt.	31407	12871	7089	4496
Cs	$6p_{1/2}$	$7p_{1/2}$	$8p_{1/2}$	$9p_{1/2}$
Theory	20204	9621	5687	3760
Expt.	20228	9641	5698	3769
Fr	$7s$	$8s$	$9s$	$10s$
Theory	32735	13051	7148	4522
Expt.	32849	13106	7168	4538
Fr	$7p_{1/2}$	$8p_{1/2}$	$9p_{1/2}$	$10p_{1/2}$
Theory	20583	9712	5724	3782
Expt.	20612	9736	5738	3795

SD Fine Structure Intervals

		Theory	Expt.
Rb	$5p_{3/2} - 5p_{1/2}$	236.5	237.6
	$6p_{3/2} - 6p_{1/2}$	76.5	77.5
	$7p_{3/2} - 7p_{1/2}$	34.8	35.1
	$8p_{3/2} - 8p_{1/2}$	18.6	18.9
Cs	$6p_{3/2} - 6p_{1/2}$	552.2	554.1
	$7p_{3/2} - 7p_{1/2}$	178.6	181.0
	$8p_{3/2} - 8p_{1/2}$	81.4	82.6
	$9p_{3/2} - 9p_{1/2}$	43.9	44.7
Fr	$7p_{3/2} - 7p_{1/2}$	1676	1687
	$8p_{3/2} - 8p_{1/2}$	536	545
	$9p_{3/2} - 9p_{1/2}$	244	250
	$10p_{3/2} - 10p_{1/2}$	132	136

SD Matrix Elements

$$\begin{aligned}
 \langle w | \left[\sum_{ij} z_{ij} a_i^\dagger a_j \right] | v \rangle &= \left(z_{wv} + \sum_{i=a}^t Z_{wv}^{(i)} \right) / (N_w N_v) \\
 Z_{wv}^{(a)} &= \sum_{am} z_{am} \tilde{\rho}_{wmva} + \text{c.c.} \quad (\text{RPA}) \\
 Z_{wv}^{(b)} &= - \sum_a z_{av} \rho_{wa} + \text{c.c.} \\
 Z_{wv}^{(c)} &= \sum_m z_{wm} \rho_{mv} + \text{c.c.} \quad (\text{BO}) \\
 &\dots \quad (16 \text{ similar terms}) \\
 Z_{wv}^{(t)} &= - \sum_{abmn} \rho_{mnba}^* z_{mv} \tilde{\rho}_{nwab} + \text{c.c.}
 \end{aligned}$$

Matrix elements are complete through 3rd order!

Dipole Transitions

Matrix Elements for heavy alkali metals (a.u.)

	K	Rb	Cs	Fr
$np_{1/2}-ns_{1/2}$	$n = 4$	$n = 5$	$n = 6$	$n = 7$
SD	4.098	4.221	4.478	4.256
Expt.	4.102(5)	4.231(3)	4.489(7)	4.277(8)
$np_{3/2}-ns_{1/2}$	$n = 4$	$n = 5$	$n = 6$	$n = 7$
SD	5.794	5.956	6.298	5.851
Expt.	5.800(8)	5.977(4)	6.324(7)	5.898(15)

SD Hyperfine Matrix Elements

^{23}Na hyperfine constants A (MHz)

	$3s_{1/2}$	$3p_{1/2}$	$3p_{3/2}$
DHF	623.5	63.39	12.59
MBPT	860.9	91.40	19.80
SD	888.1	94.99	18.84
Expt.	885.81	94.44(13)	18.534(12)

Polarizabilities

For a valence s state:

$$\begin{aligned}\alpha_v &= \frac{1}{3} \sum_m \left(\frac{|\langle v || z || mp_{1/2} \rangle|^2}{E_{mp_{1/2}} - E_v} + \frac{|\langle v || z || mp_{3/2} \rangle|^2}{E_{mp_{3/2}} - E_v} \right) \\ \alpha_c &= \frac{2}{3} \sum_{ma} \frac{|\langle a || z || m \rangle|^2}{E_m - E_a} \\ \alpha_{vc} &= \frac{1}{3} \sum_a \frac{|\langle a || z || v \rangle|^2}{E_a - E_v}\end{aligned}$$

SD Results for Polarizabilities

Static polarizabilities (a.u.) of alkali-metal atoms.

	Na	K	Rb	Cs	Fr
α_v^{main}	162.06	284.70	308.43	383.8	294.0
α_v^{tail}	0.08	0.07	0.14	0.2	1.4
α_c	0.95	5.46	9.08	15.8	20.4
α_{vc}	-0.02	-0.13	-0.26	-0.5	-0.9
α^{SD}	163.07	290.10	317.39	399.3	314.9
Recom.	162.6(0.3)	290.2(0.8)	318.6(0.6)	399.9(1.9)	317.8(2.4)
Expt.	162.7(0.8)	293.6(6.1)	319.9(6.1)	403.6(8.1)	

Summary

The SD theory gives:

- Removal energies: $1\text{--}100 \text{ cm}^{-1}$
- Fine-structure intervals: 0.5%
- Transition Amplitudes: 0.3 – 1%
- Hyperfine constants: 0.1 – 1%
- Polarizabilities: 0.1 – 1%
- Applications to PNC in Cs and Fr (Next Lecture)