## Math 10250 Activity 13: The Derivative of a Function (Section 3.2)

**GOAL:** To understand that the slope of the graph of a function f(x) is dependent on x. The function that gives the slope of the graph of f(x) is called the derivative of f(x). We will also learn about some basic properties of the derivative, as well as the derivatives of power functions and polynomial.


**Example 1** For the function y = f(x) whose graph is shown, compute or estimate the following values:

Slope of the tangent line to the graph of f(x) at x=2 is:

Slope of the graph of f(x) at x = 4 is:

Instantaneous rate of change of f(x) at x=6 is:

Rate of change of f(x) at x = -1 is:



**Remark:** The slope of the graph of f(x) or rate of change of f(x) varies according to x. This gives us a new function called the **derivative** of f(x). We denote the derivative of f(x) by f'(x).

Find the following values for the function in Example 1:

$$f'(1) \stackrel{?}{=}$$

$$f'(2) \stackrel{?}{=}$$


$$f'(4) \stackrel{?}{=}$$


$$f'(6) \stackrel{?}{=}$$


$$f'(6) \stackrel{?}{=} \qquad f'(-1) \stackrel{?}{=}$$

## ▶ Difference Quotient and Leibniz's notation

Recalling the limit definition of the rate of change of f(x), we have:








## ▶ Other Notations for the Derivative

For each fixed value a in the domain of f(x), we can also write:

f'(a) =



**Example 2** Suppose that f(x) is a function whose graph goes through the point (1,5) and whose tangent line at that point has the equation 2x + y = 7. Without computing, find each of the following limits:

(a) 
$$\lim_{h\to 0} \frac{f(1+h)-5}{h} \stackrel{?}{=}$$

(a) 
$$\lim_{h \to 0} \frac{f(1+h) - 5}{h} \stackrel{?}{=}$$
 (b)  $\lim_{\Delta x \to 0} \frac{f(1+\Delta x) - 5}{\Delta x} \stackrel{?}{=}$  (c)  $\lim_{x \to 1} \frac{f(x) - 5}{x - 1} \stackrel{?}{=}$ 

(c) 
$$\lim_{x \to 1} \frac{f(x) - 5}{x - 1} \stackrel{?}{=}$$

**Example 3** Use the definition of derivative and no other formula to find f'(x) where  $f(x) = \frac{1}{x} = x^{-1}$ .

- ▶ Rules for finding derivatives: (Notations:  $f'(x) = (f(x))' = \frac{d}{dx}[f(x)]$ )
  - $(c)' \stackrel{?}{=}$  ;e.g.,  $(8)' \stackrel{?}{=}$  or  $(\sqrt{2})' \stackrel{?}{=}$  or  $(e)' \stackrel{?}{=}$ 0. The derivative of a constant is zero:
  - 1. The Power Rule  $(x^m)' \stackrel{?}{=}$  ;e.g.,  $(x^5)' \stackrel{?}{=}$  or  $(x^{-0.8})' \stackrel{?}{=}$
  - 2. The Constant Multiple Rule  $\frac{d}{dx}[cf(x)] \stackrel{?}{=} \qquad \qquad ; e.g., (3x^5)' \stackrel{?}{=}$ 3. The Sum Rule  $\frac{d}{dx}[f(x) + g(x)] \stackrel{?}{=} \qquad \qquad ; e.g., (x^2 5x + 4)' \stackrel{?}{=}$

**Q2:** Explain why the Sum Rule is true.

**A2**:

Example 4 
$$\frac{d}{dx} \left[ 2x^4 + 3x^{-3} - \frac{\pi}{e} \right] \stackrel{?}{=}$$

**Example 5** Find the equation of the line tangent to the graph  $y = x^3 - 2x$  at x = 2.