

**Math 10360 – Example Set 05A**  
**Section 7.1 - Integration by Parts**

1. A 10m chain with non-uniform linear mass density  $\rho(y) = e^y$  kg/m for  $0 \leq y \leq 10$  is coiled on the ground is lifted straight up from its heavier end (labelled A) so that End A is 10m above the ground and the rest of the chain dangles free below. Find the work done in lifting the chain. You may take the acceleration due to gravity as  $g = 10\text{m/s}^2$ .

### Integration by Parts

**IDEA:** Recall that Integration by Substitution “reverses” chain rule. Today we learn another technique, called *integration by parts*, which “reverses” the product rule.

Let  $u(x)$  and  $v(x)$  be two differentiable functions. Applying product rule, we have:

$$\frac{d}{dx}(u(x)v(x)) = u(x)v'(x) + u'(x)v(x)$$

By definition of anti-derivative:

$$u(x)v(x) = \underline{\hspace{10em}} = \int u(x)v'(x) dx + \int u'(x)v(x) dx.$$

Rearranging terms, we have:

$$\int u(x)v'(x) dx = u(x)v(x) - \int v(x)u'(x) dx$$

Note  $\frac{du}{dx} = u'(x) \Rightarrow du = \underline{\hspace{2em}}$ . Also  $\frac{dv}{dx} = v'(x) \Rightarrow dv = \underline{\hspace{2em}}$ .

Suppressing variable  $x$ , we get:

$$\boxed{\int u dv = \underline{\hspace{10em}}}. \rightarrow \text{Integration by Parts Formula}$$

As a definite integral, we have:

$$\boxed{\int_a^b u dv = \underline{\hspace{10em}}}$$

2. Evaluate the following integrals:

(a)  $\int_0^{10} ye^y dy$

(b)  $\int x^3 \ln x dx$

(c)  $\int xe^{x^2} dx$

(d)  $\int \arctan x dx$

2e.  $\int \sin^4 x \cos x dx \stackrel{?}{=}$

**Math 10360 – Example Set 05B**  
**Section 7.2 Trigonometric Integrals**

1. Use the following identities to complete the blanks below:

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\sin A \cos B = \underline{\hspace{10cm}}$$

$$\cos A \cos B = \underline{\hspace{10cm}}$$

$$\sin A \sin B = \underline{\hspace{10cm}}$$

$$\sin(2A) = \underline{\hspace{10cm}}$$

The Pythagorean Identities and one of the given identities above, write the following in terms of  $\cos(2A)$ :

$$\cos^2 A = \underline{\hspace{10cm}}$$

$$\sin^2 A = \underline{\hspace{10cm}}$$

2. Using appropriate identities, evaluate the following definite integrals:

a.  $\int \sin(2x) \cos(3x) dx$

b.  $\int \cos^2 2z dz$

c.  $\int_0^{\pi/3} \sin^5 x dx$

d.  $\int_0^{\pi} \sin^4(x) dx$

e.  $\int_0^{\pi/2} \cos(5x) \cos(x) dx$