

Brief Article

The Author

October 29, 2004

=2.5cm=usual =0 October 29, 1997 **Math 108, Exam 2**

This exam is worth a total of 100 points. There are 25 multiple choice problems and 3 partial credit. Each multiple choice problem is assigned next to the partial credit problems. The partial credit section of the test inside the test booklet. The multiple choice section by putting a in the appropriate box. Start at 9:20am to complete the exam. Good luck!

Sign your name

6=2.5in =0.8cm =1cm =0.4cm=1 Find the constant solutions of the differential equation $y' = y^3 + 5y^2 - 24y$.

$y = 0, y = 3, y = -8$
 $y = 0, y = 2, y = -12$
 $y = 0, y = -6, y = 4$
 $y = 3, y = -8$
 $y = -6, y = 4$

Determine the relationship of the two lines:

$$x - 3y = 6, 2x + y = -1.$$

Cannot be determined. The lines are parallel. They are the same line. There are infinitely many solutions. There is a unique solution.

The function $y = t^2 + 3t + 7$ is a solution to which of the following differential equations?

$y' - y = -t^2 - 4$
 $(y')^2 - 4y = -19$
 $(y')^2 - y = 2$
 $y' - y = t^2 + 6$
 $y' = y^2 + 14$

If $\vec{x} = (1, 3, -2, 0)$ and $\vec{y} = (-1, 2, 4, -6)$ are two vectors in \mathbb{R}^4 , determine $(2\vec{x}) \cdot (-3\vec{y})$.

(6, -36, 48, 0) 18 54 (5, 0, -16, 9) -42

An experimenter reports that a certain strain of bacteria grows at a rate proportional to the square of the size of the population. Set up a differential equation which describes the growth of the population and has a solution $y = f(t)$ where $f(t)$ is the size of the population at any time t .

$$(y')^2 = k \quad y' = k^2 y \quad y' = ky \quad y' = ky^2 \quad y' = kt^2$$

Determine the solution of the differential equation $y' = 6t - 5t^2$.

$6 - 10t$
 $6t^2 - 5t^3 + C$
 $3t^2 - 5t^3 + C$
 $3t^2 - \frac{5}{3}t^3 + C$
 None of the above

What is the length of the vector $\vec{x} = (-2, 4, 2, 5)$?

7 49 $\sqrt{41}$ 9 41

Given the differential equation $y' = (y-2)(y+3)$, determine which of the following is true for the constant solutions $y = 2$ and $y = -3$.

$y = -3$ is unstable and $y = 2$ is stable.
 $y = 2$ is stable and $y = -3$ is stable.
 $y = 2$ is unstable and $y = -3$ is stable.
 $y = -3$ is unstable and $y = 2$ is unstable.
 None of the above.

Solve the following separable differential equation with the given initial value. Recall that $e^{xy} = (e^y)^x$ and that $e^{(\dots)+C} = Ae^{(\dots)}$.

(12 points)

$$y' = \frac{y^2 + 6}{yt}, \quad y(1) = 3$$

b) Sketch some solutions of this differential equation.

c) Given that the person initially deposits \$3000, solve the differential equation to determine $M(t)$ - the amount of money in the account at any time t .

Evaluate the following system of linear equations using Gaussian elimination. Your answer should be of the form $x_i = ?$. (12 points)

$$x_1 - 3x_2 + x_3 - 2x_4 = 13x_1 - 6x_2 + 12x_3 - 6x_4 = 34x_1 - 9x_2$$

(20 points)

A certain individual decides to open an Individual Retirement Account (IRA). This person makes continuous deposits of \$2000 each year. The interest rate is 5%.

a) Find the differential equation whose solution is given by the function $M(t)$ where $M(t)$ is the amount of money in the account at any time t .

