

**Math. 125 Quiz #1**      September 3, 2002

If

$$(f \circ g)(x) = f(g(x)) = (\sin x)^2 + 3 \sin x + 7$$

what is a possibility for  $f$  and  $g$ ?

If  $h(u) = \cos u$  and  $m(s) = 2s + 3$  what is  $(h \circ m)(y) = h(m(y))$ ?

If the domain of  $f$  is the interval  $(3, 7)$ , what is the domain of  $f(x + 2)$ ?

.....

$$g(x) = \sin x \text{ and } f(x) = x^2 + 3x + 7.$$
$$(h \circ m)(y) = h(m(y)) = h(2y + 3) = \cos(2y + 3).$$

The domain of  $f(x + 2)$  is in the interval  $(1, 5)$ .

**Math. 125 Quiz #2**      September 10, 2002

1. What is  $\lim_{x \rightarrow 3} \frac{3x - 1}{1 - x}$  ?

2. What is  $\lim_{x \rightarrow 2^-} \frac{1}{2 - x}$  ?

3. Suppose  $\lim_{x \rightarrow 1} f(x) = 4$  and  $\lim_{x \rightarrow 1} g(x) = -2$ . What is the value of the following limit?

$$\lim_{x \rightarrow 1} \frac{f(x) \cdot g(x)}{(4 + g(x))^2}$$

.....

1.  $\lim_{x \rightarrow 3} \frac{3x - 1}{1 - x} = \frac{3 \cdot 3 - 1}{1 - 3} = \frac{8}{-2} = -4$

2.  $\lim_{x \rightarrow 2^-} \frac{1}{2 - x} = +\infty$

3. 
$$\lim_{x \rightarrow 1} \frac{f(x) \cdot g(x)}{(4 + g(x))^2} = \frac{\lim_{x \rightarrow 1} f(x) \cdot \lim_{x \rightarrow 1} g(x)}{(4 + \lim_{x \rightarrow 1} g(x))^2} = \frac{4 \cdot (-2)}{(4 - 2)^2} = -2$$

**Math. 125 Quiz #3**      September 17, 2002

For each of the functions below, select one of the properties A-D.

- A. Is continuous for all  $x$ .
- B. Has one or more vertical asymptotes.
- C. Has one or more removable singularities.
- D. None of the above.

1. The function  $[[x]] + 1$ .
2. The function  $\frac{x^2 + 3x + 2}{x + 1}$ .
3. The function  $\frac{(x-3) \sin x}{x^2 - 6x + 9}$ .
4. The function  $x + \sqrt[3]{1 - x}$ .

.....  
1. The function  $[[x]] + 1$ .

Answer: D

There is a jump discontinuity at each integer value.

2. The function  $\frac{x^2 + 3x + 2}{x + 1}$ .

Answer: C

$$\frac{x^2 + 3x + 2}{x + 1} = \frac{(x + 2)(x + 1)}{x + 1} = x + 2$$

Therefore, we have a removable singularity at  $x = -1$ .

3. The function  $\frac{(x-3)\sin x}{x^2-6x+9}$ .

Answer: B

$$\frac{(x - 3)\sin x}{x^2 - 6x + 9} = \frac{(x - 3)\sin x}{(x - 3)(x - 3)} = \frac{\sin x}{x - 3}$$

Therefore, we have a vertical asymptote at  $x = 3$ .

4. The function  $x + \sqrt[3]{1 - x}$ .

Answer: A

Root functions are continuous everywhere in their domain. So  $\sqrt[3]{1 - x}$  is continuous on  $\mathbf{R}$ .  $g(x) = x$  is continuous on  $\mathbf{R}$ . Hence,  $x + \sqrt[3]{1 - x}$  is continuous for all  $x$ .

---

**Math. 125 Quiz #4**

October 1, 2002

Find a formula for the derivative of  $f(x) = \frac{\tan(x) + \sin(x)}{x^2 + 4x - 3}$ .

.....  
We have a quotient so the answer is

(derivative of the numerator)  $\times$  (denominator) - (numerator)  $\times$  (derivative of the denominator) divided by (the denominator) squared

$$f'(x) = \frac{(\sec^2(x) + \cos(x))(x^2 + 4x - 3) - (\tan(x) + \sin(x))(2x + 4)}{(x^2 + 4x - 3)^2}$$

---

**Math. 125 Quiz #5**

October 8, 2002

1.a) Find a formula for  $\frac{dy}{dx}$  if

$$x^4 + 5xy^2 + y^5 = -5 .$$

1.b) Find an equation for the tangent line to the curve at the point  $(-1, -1)$ .

2 ) Suppose that, when  $x = 0$ ,  $f'(0) = -0.5253$ . If  $h(x) = f(x^2 + 10x)$ , find  $h'(-10)$ .

.....  
1a) Using implicit differentiation:

$$4x^3 + (5y^2 + 10xyy') + 5y^4y' = 0$$

or

$$y' = -\frac{4x^3 + 5y^2}{10xy + 5y^4}$$

1b) Line goes thru  $(-1, -1)$  and has slope  $\frac{4(-1)^3 + 5(-1)^2}{10(-1)(-1) + 5(-1)^4} = -\frac{5-4}{10+5} = -\frac{1}{15}$ ,  
so an equation for the tangent line is  $y - (-1) = -\frac{1}{15}((x - (-1)))$ , or  $15y + 15 = -(x + 1)$ .

2) By the Chain Rule,  $h'(x) = f'(x^2 + 10x)(2x + 10)$ , so  
 $h'(-10) = f'(0) \cdot (2(-10) + 10) = -0.5253 \cdot (-10) = 5.253$ .

---

**Math. 125 Quiz #6**      October 15, 2002

Consider a clock whose hour hand has length  $a$  and whose minute hand has length  $b$ .  
At 3 o'clock, how fast is the distance between the tip of the hour hand and the tip of the  
minute hand changing?

Let  $\theta$  be the angle between the minute and hour hand. The distance,  $c$ , between the  
tips is given by the law of cosines

$$c^2 = a^2 + b^2 - 2ab \cos(\theta) .$$

Step 1. It is true that  $\frac{d\theta}{dt}$  is a constant. What constant is it?

**Hint:**  $\theta = \theta_{\text{hr}} - \theta_{\text{min}}$  where  $\theta_{\text{hr}}$  is the angle the hour hand makes with the vertical and  $\theta_{\text{min}}$   
is the angle the minute hand makes with the vertical. Both  $\frac{d\theta_{\text{hr}}}{dt}$  and  $\frac{d\theta_{\text{min}}}{dt}$  are constants.  
What are they?

Step 2. Use this fact and your related rates techniques to answer the original question.  
If the distance is *decreasing* be sure your formula gives a negative value; if increasing, a  
positive one.

.....

$$c^2 = a^2 + b^2 - 2ab \cos(\theta) \Rightarrow 2c \frac{dc}{dt} = -2ab(-\sin(\theta)) \frac{d\theta}{dt} = 2ab \sin(\theta) \frac{d\theta}{dt}$$

$$\Rightarrow \frac{dc}{dt} = \frac{2ab \sin(\theta) \frac{d\theta}{dt}}{2c} = \frac{ab \sin(\theta) (-\frac{11\pi}{6})}{c}$$

At 3:00,  $\theta = \frac{\pi}{2}$  and  $\sin(\theta) = \sin(\frac{\pi}{2}) = 1$ . Therefore, at 3:00,

$$\frac{dc}{dt} = \frac{ab \sin(\frac{\pi}{2}) (-\frac{11\pi}{6})}{c} = \frac{ab(-\frac{11\pi}{6})}{c} = -\frac{11ab\pi}{6c}.$$

**Math. 125 Quiz #7**      November 5, 2002

1. Find all the critical points for the function below. At which critical points is the function value a local maximum? Why? At which critical points is the function value a local minimum? Why?
2. The existence theorem for global max/min does not apply to this function restricted to the interval  $[0, \infty)$  since the interval must be a closed interval for the theorem to apply. Nevertheless, this function does assume a minimum value on this interval. Where is it and why? Why does this function assume no maximum value on this interval?

$$\frac{(x-3)^6}{(x+1)^2}$$

.....

1. Let  $r(x) = \frac{(x-3)^6}{(x+1)^2}$ . Therefore,

$$r'(x) = \frac{(x+1)^2 \cdot 6 \cdot (x-3)^5 - (x-3)^6 \cdot 2 \cdot (x+1)}{(x+1)^4}$$

$$= \frac{(x+1)(x-3)^5 [6(x+1) - 2(x-3)]}{(x+1)^4} = \frac{(x-3)^5 (4x+12)}{(x+1)^3} = \frac{4(x-3)^5 (x+3)}{(x+1)^3}.$$

$r'(x) = 0 \Leftrightarrow (x-3)^5(x+3) = 0 \Leftrightarrow x = -3, 3$ .  $r'(x)$  does not exist  $\Leftrightarrow (x+1)^3 = 0 \Leftrightarrow x = -1$ . So the critical numbers of  $r(x)$  are  $x = -3, -1, 3$ . We have  $x \in (-\infty, -3) \Rightarrow r'(x) < 0$ ,  $x \in (-3, -1) \Rightarrow r'(x) > 0$ ,  $x \in (-1, 3) \Rightarrow r'(x) < 0$  and  $x \in (3, \infty) \Rightarrow r'(x) > 0$ . Hence, local minima occur at  $x = -3, 3$  because the derivative changes from negative to positive at these points. There is no local minimum or maximum at  $x = -1$  since  $r(x)$  is not defined there.

2.  $\lim_{x \rightarrow \infty} r(x) = \infty$  and  $r(0) = 729$ . We suspect that the global minimum occurs at  $x = 3$  since this point is in the interval of interest and a local minimum occurs here. By the first derivative test, the value of  $r(x)$  is increasing on  $(3, \infty)$  so  $r(3) \leq r(x)$  for  $x > 3$ . Likewise, the value of  $r(x)$  is decreasing on  $[0, 3)$  so  $r(x) \geq r(3)$  for  $x$  less than 3 and greater than or

equal to 0. Hence  $r(3) \leq r(x)$  for all  $x \in [0, \infty)$ , which says that the value at  $x = 3$  is a global minimum for the function on the interval  $[0, \infty)$ .

The idea in the last paragraph is a general result. Suppose

- a) a function  $f(x)$  is differentiable on an interval (closed, open, or half open),
- b)  $x = c$  is the *only* critical number in the interval, and
- c) the value of  $f$  is a local maximum (respectively minimum) at  $x = c$ .

Then  $f(c)$  is a global maximum (respectively minimum) on that interval.

No maximum value occurs in  $[0, \infty)$  since  $\lim_{x \rightarrow \infty} r(x) = \infty$ .

**Math. 125 Quiz #8**      November 12, 2002

We wish to use Newton's method to find a solution to the equation

$$\cot x = 2 \tan x .$$

If we make a first approximation of  $x = \frac{\pi}{4}$ , what does Newton's method give for a second approximation?

Leave your answer in terms of sums of fractions and rational multiples of  $\pi$ .

.....

To apply Newton's method, observe that our original equation is equivalent to  $f(x) = 0$  where  $f(x) = \cot x - 2 \tan x$ .

Since  $f'(x) = -\csc^2 x - 2 \sec^2 x$ , we get

$$x_{n+1} = x_n - \frac{\cot x - 2 \tan x}{-\csc^2 x - 2 \sec^2 x} = x_n + \frac{\cot x - 2 \tan x}{\csc^2 x + 2 \sec^2 x} .$$

Taking  $x_0 = \frac{\pi}{4}$ , we get

$$x_1 = \frac{\pi}{4} - \frac{f(\frac{\pi}{4})}{f'(\frac{\pi}{4})}$$

$f(\frac{\pi}{4}) = \cot(\frac{\pi}{4}) - 2 \tan(\frac{\pi}{4}) = 1 - 2 = -1$ ,  $f'(\frac{\pi}{4}) = -\csc^2(\frac{\pi}{4}) - 2 \sec^2(\frac{\pi}{4}) = -2 - 2 \cdot 2 = -6$  so  $x_1 = \frac{\pi}{4} - \frac{-1}{-6} = \frac{\pi}{4} - \frac{1}{6}$ .

**Math. 125 Quiz #9: Solutions**      November 19, 2002

1. Consider the region in the plane below the graph  $y = \csc x$ ; above the  $x$ -axis and between the vertical lines  $x = 0$  and  $x = 2$ .

- a. Divide the interval into 6 equal parts and write down the Riemann sum using the mid-point rule. Do not evaluate the sum or even try to simplify it.

.....

Midpoints are  $\frac{13}{12}, \frac{15}{12}, \dots, \frac{23}{12}$  : The sum is  $\sum_{i=1}^6 \frac{1}{6} \csc\left(\frac{13}{12} + \frac{i-1}{6}\right) = \sum_{i=1}^6 \frac{1}{6} \csc\left(\frac{11}{12} + \frac{i}{6}\right) = \frac{1}{6} \left( \csc\left(\frac{13}{12}\right) + \csc\left(\frac{15}{12}\right) + \dots + \csc\left(\frac{23}{12}\right) \right)$ .

.....

b. Write a definite integral which evaluates the area precisely. Do not evaluate the integral.

.....

$$\int_1^2 \csc x dx.$$

.....

2. What number is represented by the sum

$$\sum_{q=3}^6 (q+2) ?$$

.....

$$\sum_{q=3}^6 (q+2) = (3+2) + (4+2) + (5+2) + (6+2) = 26.$$


---

**Math. 125 Quiz #10**      Decsmbler 10, 2002

1. Write a definite integral that evaluates to the area of the shaded region below. Do not evaluate the integral.
2. Rotate the shaded region about the line  $y = 2$  and write a definite integral that evaluates to the volume of the resulting solid of revolution. Do not evaluate the integral.
3. Rotate the shaded region about the line  $x = -3$  and write a definite integral that evaluates to the volume of the resulting solid of revolution. Do not evaluate the integral.

=800

graph10.eps

.....

Because the curves are given as  $y$  equals a function of  $x$ , the answers will be integrals with respect to  $dx$  and so the first issue is to find the points of intersection of the two curves, or solve  $\sin x = \cos x$  or  $\tan x = 1$ .  $x = \frac{\pi}{4}$  is one solution and  $x = \frac{5\pi}{4}$  is the next and these are all the solutions in the relevant interval.

The area is given by the integral  $\int_{\frac{\pi}{4}}^{\frac{5\pi}{4}} (\sin x - \cos x) dx$ : top curve minus bottom curve.

The volume integral when rotated about  $y = 2$  is given by

$$\pi \int_{\frac{\pi}{4}}^{\frac{5\pi}{4}} \left( (2 - \cos x)^2 - (2 - \sin x)^2 \right) dx .$$

With this set up, we *must* use the washer method. The outer radius is the distance between  $y = 2$  and  $y = \cos x$  because from the graphs, the distance from  $y = 2$  to  $y = \sin x$  is smaller.

The volume integral when rotated about  $x = -3$  is given by

$$2\pi \int_{\frac{\pi}{4}}^{\frac{5\pi}{4}} (x - (-2)) \cdot (\sin x - \cos x) dx .$$

With this set up, we *must* use the shell method.

---