

Exam I
February 15, 2001

11.

- (a) First we need to determine the corner points. They are $(\frac{1}{4}, 4)$, $(1, 4)$, $(4, 1)$ and $(4, \frac{1}{4})$. It is also good to note at this point that the integral will need to be split into two pieces. We also note that we may take the density to be 1 in this problem. Hence if we integrate along the x -axis we get for the area (= mass)

$$\begin{aligned} \text{Area} &= \int_{\frac{1}{4}}^1 \left(4 - \frac{1}{x}\right) dx + \int_1^4 \left(\frac{4}{x} - \frac{1}{x}\right) dx \\ &= [4x - \ln x]_{\frac{1}{4}}^1 + [3 \ln x]_1^4 \\ &= (4 - \ln 1) - \left(1 - \ln \frac{1}{4}\right) + 3 \ln 4 - 3 \ln 1 \\ &= 3 + \ln \frac{1}{4} + 3 \ln 4 = 3 - \ln 4 + 3 \ln 4 = 3 + 2 \ln 4 \end{aligned}$$

If you preferred to integrate up the y -axis you get an identical formula except that everywhere you see an x above, put a y .

- (b) You have four choices. Do you want to compute the moment about the x -axis or the y -axis and do you want to integrate along the x -axis or the y -axis? To compute the moment about the y -axis and integrate along the x -axis, proceed as follows.

$$\begin{aligned} \text{Moment}_y &= \int_{\frac{1}{4}}^1 \underbrace{x}_{\substack{\text{dist.} \\ \text{from} \\ y\text{-axis}}} \underbrace{\left(4 - \frac{1}{x}\right) dx}_{\text{mass of rect.}} + \int_1^4 x \left(\frac{4}{x} - \frac{1}{x}\right) dx \\ &= [2x^2 - x]_{\frac{1}{4}}^1 + [3x]_1^4 = 2 - 1 - \frac{2}{16} + \frac{1}{4} + 12 - 3 = \frac{81}{8} \end{aligned}$$

If you choose instead to calculate the moment about the x -axis while still integrating along the x -axis you would proceed as follows.

$$\begin{aligned}
 \text{Moment}_x &= \int_{\frac{1}{4}}^1 \underbrace{\frac{1}{2} \left(4 + \frac{1}{x}\right)}_{\substack{\text{dist. from} \\ \text{center of mass} \\ \text{to } x\text{-axis}}} \underbrace{\left(4 - \frac{1}{x}\right) dx}_{\text{mass of rect.}} + \int_1^4 \frac{1}{2} \left(\frac{4}{x} + \frac{1}{x}\right) \left(\frac{4}{x} - \frac{1}{x}\right) dx \\
 &= \int_{\frac{1}{4}}^1 8 - \frac{1}{2x^2} dx + \int_1^4 \frac{15}{2x^2} dx \\
 &= \left[8x + \frac{1}{2x}\right]_{\frac{1}{4}}^1 + \left[\frac{-15}{2x}\right]_1^4 = \left(8 + \frac{1}{2}\right) - \left(2 + 2\right) + \left(-\frac{15}{8} - \left(-\frac{15}{2}\right)\right) \\
 &= 4 + \frac{4}{8} + \frac{-15}{8} + \frac{60}{8} = \frac{32}{8} + \frac{4}{8} + \frac{-15}{8} + \frac{60}{8} = \frac{96 - 15}{8} = \frac{81}{8}
 \end{aligned}$$

If you want the integrals for integrating along the y -axis, just switch x and y in the formulae above.

- (c) By symmetry, the moment about the x -axis is the same as the moment about the y -axis and the center of mass has equal x and y coordinates, each one being

$$\frac{\frac{81}{8}}{3 + 2 \ln 4}$$

12. Complete the square: $x^2 + 2x + 2 = (x + 1)^2 + 1$. This suggests the trig. substitution $\tan \theta = x + 1$: $dx = \sec^2 \theta d\theta$ and $x^2 + 2x + 2 = (x + 1)^2 + 1 = \tan^2 \theta + 1 = \sec^2 \theta$. When $x = -1$, $x + 1 = 0 = \tan \theta$, so $\theta = 0$; when $x = 0$, $x + 1 = 1 = \tan \theta$, so $\theta = \frac{\pi}{4}$. Hence

$$\int_{-1}^0 \frac{1}{x^2 + 2x + 2} dx = \int_0^{\frac{\pi}{4}} \frac{\sec^2 \theta}{\sec^2 \theta} d\theta = \int_0^{\frac{\pi}{4}} d\theta = \theta \Big|_0^{\frac{\pi}{4}} = \frac{\pi}{4} - 0 = \frac{\pi}{4}.$$

13. Separating variables gives $\frac{dy}{\sqrt{1-y^2}} = x$: integrating gives

$$\arcsin(y) = \frac{x^2}{2} + C.$$

Now since $y(0) = 0$ we see that the constant $C = 0$.
Hence the unique function is

$$y = \sin\left(\frac{x^2}{2}\right)$$

and $y(1) = \sin\left(\frac{1}{2}\right)$.

14. To find the general solution to the differential equation

$$y' = \frac{2y}{x} + x \quad (x > 0)$$

first put it into standard form:

$$y' + \frac{-2}{x} \cdot y = x$$

so $P(x) = \frac{-2}{x}$ and $Q(x) = x$. Then find the integrating factor $v = e^{\int P(x) dx}$.

$\int \frac{-2}{x} = -2 \ln x + C$ so we may take $v(x) = e^{-2 \ln x}$ which is $v(x) = x^{-2}$. Then we do the second integral $\int v(x) Q(x) dx = \int x^{-2} \cdot x dx = \int x^{-1} dx = \ln x + C$. Since we are told $x > 0$, there is no need for absolute values in the $\ln x$. Finally,

$$y(x) = x^{-2}(\ln x + C)$$

is the general solution.
