

13. Find the area under the region bounded by the curves $y = 2^x$ and $y = 5^x$, between $x = 0$ and $x = 1$.

.....
 Since $5^x > 2^x$ for $x > 0$ the requested area is

$$\int_0^1 5^x - 2^x dx = \left. \frac{5^x}{\ln 5} - \frac{2^x}{\ln 2} \right|_0^1 = \left(\frac{5}{\ln 5} - \frac{2}{\ln 2} \right) - \left(\frac{1}{\ln 5} - \frac{1}{\ln 2} \right) = \left(\frac{4}{\ln 5} - \frac{1}{\ln 2} \right).$$

14. Evaluate the limit

$$\lim_{x \rightarrow 0^+} (\sin x)^{\tan x}$$

.....
 Let $y = \lim_{x \rightarrow 0^+} (\sin x)^{\tan x}$. Then

$$\ln y = \lim_{x \rightarrow 0^+} \tan x \ln(\sin x) = \lim_{x \rightarrow 0^+} \frac{\ln(\sin x)}{\cot x}.$$

When $x \rightarrow 0^+$, $\tan(x) \rightarrow 0$, $\cot(x) \rightarrow \infty$ and $\ln(\sin x) \rightarrow -\infty$.

This is an indeterminate form where we can apply L'Hospital's rule, so

$$\ln y = \lim_{x \rightarrow 0^+} \frac{\frac{\cos x}{\sin x}}{-\csc^2 x} = \lim_{x \rightarrow 0^+} -\sin x \cos x = 0.$$

Since $\ln y = 0$, $y = e^0 = 1$.

15. Use logarithmic differentiation to find the derivative of

$$y = \frac{(x^2 + 1)^{2.2}(x + 1)^{1.2}}{(x^3 - 1)^{0.3}}$$

.....
 This is a logarithmic differentiation problem, so start by taking the log of both sides and simplifying the right hand side.

$$\begin{aligned} \ln y &= \ln \left(\frac{(x^2 + 1)^{2.2}(x + 1)^{1.2}}{(x^3 - 1)^{0.3}} \right) \\ &= \ln((x^2 + 1)^{2.2}) + \ln((x + 1)^{1.2}) - \ln((x^3 - 1)^{0.3}) \\ &= 2.2 \ln(x^2 + 1) + 1.2 \ln(x + 1) - 0.3 \ln(x^3 - 1) \end{aligned}$$

Differentiating both sides with respect to x we get

$$\frac{y'}{y} = 2.2 \frac{2x}{x^2 + 1} + 1.2 \frac{1}{x + 1} - 0.3 \frac{3x^2}{x^3 - 1} = \frac{4x}{x^2 + 1} + \frac{1.2}{x + 1} - \frac{0.9x^2}{x^3 - 1}$$

Now solve for y'

$$\frac{dy}{dx} = \left(\frac{(x^2 + 1)^{2.2}(x + 1)^{1.2}}{(x^3 - 1)^{0.3}} \right) \cdot \left(\frac{4x}{x^2 + 1} + \frac{1.2}{x + 1} - \frac{0.9x^2}{x^3 - 1} \right)$$