

MATH 225: Calculus III**Practice Exam I** *September 2002*

Find the angle between the planes $x - y + 2z = 3$ and $2x + y + z = -5$.

$\pi/3$ $\pi/6$ $\pi/2$ $3\pi/4$ $\pi/4$

Compute the arc length of the curve parameterized by $\mathbf{r}(t) = 2t\mathbf{i} + t^2\mathbf{j} + \frac{1}{3}t^3\mathbf{k}$ for $0 \leq t \leq 1$.

$\frac{7}{3}$ $\frac{\sqrt{46}}{3}$ $\frac{17}{12}$ $\frac{10}{3}$ $\frac{\sqrt{161}}{12}$

Find the parametric equations of the line tangent to the curve $\mathbf{r}(t) = \langle t^2, t, \ln t \rangle$ at the point $(1, 1, 0)$.

$x = 1 + 2t, y = 1 + t, z = t$ $x = 2t, y = 1, z = \frac{1}{t}$ $x = 2 + t, y = 1 + t, z = 1$ $x = 1 + 2t^2, y = 1 + t, z = 1$ $x = 1 + t, y = 1 + 2t, z = -t$

The position of a moving particle at time t is given by $\mathbf{r}(t) = \langle \cos(2t), t, \sin(2t) \rangle$. Determine the tangential and normal components of the acceleration at $t = 0$.

$a_T = 0, a_N = 4$ $a_T = -4, a_N = 0$ $a_T = \sqrt{5}, a_N = \sqrt{11}$ $a_T = \sqrt{5}, a_N = 0$ $a_T = 1, a_N = \sqrt{5}$

Determine which of the following symmetric equations gives a line that is parallel to the vector $\mathbf{i} + 2\mathbf{j} + (1/2)\mathbf{k}$.

$2(x + 1) = y - 3 = 4(z - 1)$ $x + 1 = 2y + 1 = (1/2)z + 1$ $2x = y = (1/2)z$ $x = 2y$ and $z = 1$ $y = (1/4)z$ and $x = 0$

Find a vector that is orthogonal to the plane containing the points $(1, 0, 1)$, $(2, -1, 1)$, and $(0, 0, 3)$.

$2\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ $-\mathbf{i} + \mathbf{k}$ $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ $2\mathbf{i} + 3\mathbf{j} + 4\mathbf{k}$ $\mathbf{i} + \mathbf{j} - \mathbf{k}$

Find the directional derivative of $f(x, y) = e^{x \sin(y)}$ at $P(1, 0)$ in the direction of $Q(-2, 4)$.

$4/5$ $-3/5$ -2 0 4

8. Find a vector function that traces out the curve of intersection of the ellipsoid $(x - 1)^2 + 3(y - 1)^2 + (z - 1)^2 = 1$ and the plane $y = x$.
9. The force acting on a moving particle of mass $m = 2$ at time t is $\mathbf{F}(t) = e^t\mathbf{i} + e^{-t}\mathbf{j} + t\mathbf{k}$. If the particle's initial position and velocity are $\mathbf{r}(0) = \langle 0, 0, 1 \rangle$ and $\mathbf{v}(0) = \langle 0, 1, 0 \rangle$, respectively, determine the particle's position vector, $\mathbf{r}(t)$, for any $t \geq 0$.
10. Let $\mathbf{r}(t) = t\mathbf{i} + t\mathbf{j} + \frac{t^2}{2}\mathbf{k}$. Find the unit tangent vector, $\mathbf{T}(0)$, the unit normal vector, $\mathbf{N}(0)$, and the unit binormal vector $\mathbf{B}(0)$ at $t = 0$.

11. Evaluate the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{y^4 + 2x^3y}{(x^2 + y^2)^2}$$

or show it does not exist.

12. A triangular sheet of glass is expanding. When the base is 2 in and the height is 4 in, the base is increasing at the rate of 0.25 in/hr and the height at 0.5 in/hr. At what rate is the area of the triangle increasing?