

Multiple Choice 8 points each

1. The Laplace transform of $f(t) = t^3 e^{2t}$ is

- (a) $\frac{3}{(s-2)^4}$ (b) $\frac{6}{(s-2)^4}$ (c) $\frac{6}{(s-2)^3}$ (d) $\frac{6}{(s+2)^4}$ (e) $\frac{1}{(s-2)^4}$

2. If $f(t)$ has Laplace transform $F(s) = \frac{s}{s^3 + 1}$ then the Laplace transform of $e^{2t}f(t)$ is

- (a) $\frac{2s}{4s^3 + 1}$ (b) $\frac{s}{s^3 + 2}$ (c) $\frac{s}{(s-2)^3 + 1}$
(d) $\frac{s+2}{(s+2)^3 + 1}$ (e) $\frac{s-2}{(s-2)^3 + 1}$

3. The inverse Laplace transform of $F(s) = \frac{2e^{-2s}}{s^2 - 4}$ is

- (a) $u_2(t) \sinh 2(t-2)$ (b) $u_2(t) \sin(2t-2)$ (c) $u_2(t) \sinh(2t+2)$
(d) $u_2(t) \cosh 2(t-2)$ (e) $u_2(t) \cosh(2t-2)$

4. The value of the improper integral

$$\int_0^{\infty} e^{(t-2)^2} \sin \frac{\pi t}{6} \delta(t-3) dt$$

- (a) e (b) 0 (c) $\frac{e^2}{6}$ (d) ∞ (e) $e^{(t-2)^2} \sin \frac{\pi t}{6}$

5. The solution of the initial value problem

$$y'' + y = \delta(t - \pi) \cos t ; \quad y(0) = 0, y'(0) = 1 \quad \text{is}$$

- (a) $2 \cos t$ (b) $2 \sin t$ (c) $\sin t + u_{\pi}(t) \sin t$
 (d) $\sin t + u_{\pi}(t) \cos t$ (e) $\cos t + u_{\pi}(t) \cos t$

6. The Laplace transform of $f(t) = \int_0^t e^{-(t-\tau)} \sin \tau \, d\tau$

- (a) $\frac{1}{(s-1)(s^2+1)}$ (b) $\frac{1}{s(s^2+1)}$ (c) $\int_0^t e^{-s} \sin s \, ds$
 (d) $\frac{1}{(s+1)(s^2+1)}$ (e) $\frac{s}{(s+1)(s^2+1)}$

7. The largest linearly independent subset of the set of vectors $X^{(1)} = (1,1,1)$, $X^{(2)} = (5,-2,5)$, $X^{(3)} = (1,-1,1)$ $X^{(4)} = (1,0,1)$ has exactly

- (a) two members (b) one member (c) three members
 (d) four members (e) no members

8. One solution of $X' = \begin{pmatrix} 1 & -4 \\ 4 & -7 \end{pmatrix} X$ is

$X^{(1)} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^{-3t}$. A linearly independent second solution $X^{(2)}$ is

- (a) $\begin{pmatrix} 1 \\ 1 \end{pmatrix} te^{-3t} + \begin{pmatrix} 0 \\ 3 \end{pmatrix} e^{-3t}$ (b) $\begin{pmatrix} 4 \\ 1 \end{pmatrix} te^{-3t} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^{-3t}$

$$(c) \begin{pmatrix} 1 \\ 3 \\ 4 \end{pmatrix} te^{-3t} - \begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} e^{-3t}$$

$$(d) \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} te^{-3t} - \begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} e^{-3t}$$

$$(e) \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} te^{-3t} + \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} e^{-3t}$$

9. A real solution of $X' = \begin{pmatrix} -1 & -4 \\ 1 & -1 \end{pmatrix} X$ is

$$(a) \begin{pmatrix} -2 \sin 2t \\ \cos 2t \end{pmatrix} e^{-t}$$

$$(b) \begin{pmatrix} 2 \cos 2t \\ -\sin 2t \end{pmatrix} e^{-t}$$

$$(c) \begin{pmatrix} \cos 2t \\ \sin 2t \end{pmatrix} e^{-t}$$

$$(d) \begin{pmatrix} 2 \cos 2t \\ \sin 2t \end{pmatrix} e^{-t}$$

$$(e) \begin{pmatrix} \sin 2t \\ 2 \cos 2t \end{pmatrix} e^t$$

10. A fundamental system of solutions of $X' = \begin{pmatrix} 2 & 1 \\ 0 & -3 \end{pmatrix} X$ is

$X^{(1)} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{2t}$, $X^{(2)} = \begin{pmatrix} 1 \\ -5 \end{pmatrix} e^{-3t}$. A fundamental matrix $\Phi(t)$ such that

$\Phi(0) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is

$$(a) \begin{pmatrix} e^{-3t} & 0 \\ 0 & e^{2t} \end{pmatrix}$$

$$(b) \begin{pmatrix} e^{2t} & 5(e^{2t} - e^{-3t}) \\ 0 & e^{-3t} \end{pmatrix}$$

$$(c) \begin{pmatrix} (2e^{2t} - e^{-3t}) & 0 \\ (e^{-3t} - e^{2t}) & e^{-3t} \end{pmatrix}$$

$$(d) \begin{pmatrix} e^{2t} & \frac{1}{5}(e^{2t} - e^{-3t}) \\ 0 & e^{-3t} \end{pmatrix}$$

$$(e) \begin{pmatrix} e^{2t} & (e^{2t} - 1) \\ (e^{-3t} - e^{2t}) & e^{-3t} \end{pmatrix}$$

11. The point $(-3,7)$ is a critical point for the nonlinear system

$$x' = x^2 + 5x + 6$$

$$y' = x + y - 4$$

This point is

(a) An unstable improper node

(b) A stable spiral point

(c) A saddle point

(d) a center

(e) A stable proper node

12. Suppose that $\phi(x)$ satisfies the initial value problem

$$y' = y + 4t \quad y(0) = 3$$

Use Euler's method with stepsize $h = 1$ and two iterations to estimate $\phi(2)$.

The result is $\phi(2) \cong$

(a) 16

(b) 4

(c) 0

(d) -7

(e) 2

13. The value of the coefficient b_7 in the sine series with period 2π

$$\sum_{n=1}^{\infty} b_n \sin nx \quad \text{for } f(x) = 3\pi \quad \text{on the interval } 0 \leq x < \pi \text{ is}$$

- (a) $\frac{4}{7}$ (b) $\frac{12}{7}$ (c) $\frac{-4}{7}$ (d) 0 (e) $\frac{3\pi}{7}$

14. Let $f(x) = \begin{cases} 5 \cos \frac{x}{3} & 0 \leq x < \frac{3\pi}{2} \\ 0 & \frac{3\pi}{2} \leq x < 3\pi \end{cases}$

be periodic with period 3π . The value of $f(x)$ at $x = 3\pi$ so that the Fourier series for f converges to f at every point is $f(3\pi) =$

- (a) -5 (b) 5 (c) 0 (d) $\frac{-5}{2}$ (e) $\frac{5}{2}$

15. For the heat conduction problem with insulated ends

$$u_{xx} = u_t, \quad u_x(0,t) = 0, \quad u_x(3,t) = 0$$

the solutions are of the form (set $\lambda = \frac{n\pi}{3}$) : $u_n(x,t) =$

- (a) $e^{-\lambda^2 t} \cos \lambda x$ (b) $e^{-\lambda^2 t} \sin 3\lambda x$ (c) $e^{-3\lambda^2 t} \cos \lambda t$
 (d) $e^{-\lambda^2 t^2} \sin \lambda x$ (e) $e^{-\lambda^2 x^2} \cos \lambda t$

Partial Credit Problems

16. A 50 cm bar of metal for which $\alpha^2 = 2$ has one end held at 10°C and the other held at 35°C . The bar is initially heated in such a way that

$$u(x,0) = f(x) = 3 \sin\left(\frac{5\pi x}{50}\right).$$

(a) (18 points) Solve the heat conduction problem for $u(x,t)$

(b) (6 points) find the steady state temperature distribution in the bar

17. Let $f(x) = x^2 + 1$ for $0 \leq x \leq 2$.

(6 points) Define a periodic extension of f with period 4 which is neither even nor odd. Sketch the graph for three periods; one of which is to the left of $x = 0$ and one is to the right.