

PRACTICE TEST 2

1. (9 points) Compute the Wronskian of the vectors

$$\mathbf{x}^{(1)} = \begin{bmatrix} t^2 \\ t \\ 0 \end{bmatrix} \quad \mathbf{x}^{(2)} = \begin{bmatrix} -t \\ 1 \\ 0 \end{bmatrix} \quad \mathbf{x}^{(3)} = \begin{bmatrix} 0 \\ 0 \\ e^t \end{bmatrix} .$$

- a. $(t^3 - t^2)e^t$ b. 0 c. $(t^2 + 1)e^t$ d. $2t^2e^t$ e. $-2te^t$

2. (9 points) Find all of the eigenvalues of the matrix $\begin{bmatrix} -1 & -3 & 2 \\ 0 & 2 & 0 \\ 3 & 0 & -2 \end{bmatrix}$.

- a. 1, 2 and -4 b. 1, 2 and 3 c. 1, 2 and -3
d. -1, -2 and 4 e. 1, 2 and 4

3. (9 points) The matrix $A = \begin{bmatrix} -2 & 1 & 4 \\ 0 & -2 & -3 \\ 0 & 0 & 2 \end{bmatrix}$ has $r = -2$ as a repeated

eigenvalue of multiplicity two. Find all of the eigenvectors \mathbf{v} of A corresponding to this eigenvalue (the arbitrary constants appearing in each of the answers may be assumed to be not all zero).

a. $\mathbf{v} = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

b. $\mathbf{v} = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

c. $\mathbf{v} = c_1 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

d. $\mathbf{v} = c_1 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

e. $\mathbf{v} = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$

4. (9 points) Find the inverse of the matrix $\begin{bmatrix} 1 & 2 \\ -3 & 4 \end{bmatrix}$.

November 30, 2004

a. $\begin{bmatrix} \frac{-1}{10} & \frac{-2}{10} \\ \frac{3}{10} & \frac{-4}{10} \end{bmatrix}$

b. $\begin{bmatrix} \frac{1}{10} & \frac{-2}{10} \\ \frac{3}{10} & \frac{4}{10} \end{bmatrix}$

c. $\begin{bmatrix} \frac{4}{10} & \frac{-2}{10} \\ \frac{3}{10} & \frac{1}{10} \end{bmatrix}$

d. $\begin{bmatrix} \frac{4}{10} & \frac{-2}{10} \\ \frac{-3}{10} & \frac{1}{10} \end{bmatrix}$

e. $\begin{bmatrix} \frac{3}{10} & \frac{-4}{10} \\ \frac{1}{10} & \frac{-2}{10} \end{bmatrix}$

5.(9 points) Find the general solution of $x' = \begin{bmatrix} 1 & -2 \\ 2 & 1 \end{bmatrix} x$.

NOTE: Be aware that the general solution that you find should be one of the answers below. . . perhaps after a suitable re-naming of your arbitrary constants.

a. $e^t \begin{bmatrix} c_1 \sin t + c_2 \cos t \\ -c_1 \cos t + c_2 \sin t \end{bmatrix}$

b. $e^{-t} \begin{bmatrix} -c_1 \sin 2t + c_2 \cos 2t \\ c_1 \cos 2t + c_2 \sin 2t \end{bmatrix}$

c. $e^t \begin{bmatrix} c_1 \sin 2t + c_2 \cos 2t \\ c_1 \cos 2t + c_2 \sin 2t \end{bmatrix}$

d. $e^{2t} \begin{bmatrix} -c_1 \sin t + c_2 \cos t \\ c_1 \cos t + c_2 \sin t \end{bmatrix}$

e. $e^t \begin{bmatrix} -c_1 \sin 2t + c_2 \cos 2t \\ c_1 \cos 2t + c_2 \sin 2t \end{bmatrix}$

6. (9 points) A fundamental matrix for the system $x' = \begin{pmatrix} 1 & 0 \\ 0 & -2 \end{pmatrix} x$ is

a. $\begin{pmatrix} e^t & e^{-2t} \\ e^t & e^{-2t} \end{pmatrix}$

b. $\begin{pmatrix} e^t & e^{-2t} \\ e^{-2t} & e^t \end{pmatrix}$

c. $\begin{pmatrix} e^t & 0 \\ 0 & e^{-2t} \end{pmatrix}$

d. $\begin{pmatrix} e^{-2t} & 0 \\ 0 & e^t \end{pmatrix}$

e. $\begin{pmatrix} e^{-t} & 0 \\ 0 & e^{2t} \end{pmatrix}$

7. (9 points) Using row reduction on the augmented matrix (or otherwise) find the number k such that the system

$$x_1 - 2x_2 + 3x_3 = k$$

$$-x_1 + x_2 - 2x_3 = -1$$

$$2x_1 - x_2 + 3x_3 = 0$$

has a solution.

- a. $k=2$ b. $k=3$ c. $k=1$ d. $k=0$ e. $k=-1$

8.(9 points) The eigenvalues of the matrix $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & 1 \end{pmatrix}$ are

- a. All real and unequal b. All real and equal
 c. $1 + 2i, 1 - 2i$ and 1 d. $1 + i, 1 - i$ and 1 e. $1, i$ and $-i$

PARTIAL CREDIT

9. (14 points) Find the general solution of the system $\mathbf{x}' = \begin{bmatrix} 2 & 0 \\ -1 & 2 \end{bmatrix} \mathbf{x}$.

10.(14 points) Find the general solution of the non-homogeneous system

$$\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 2e^{-2t} \\ 0 \end{bmatrix}$$

evaluating all integrals that occur.

Hint: A fundamental matrix of the system $\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x}$

is $\psi(t) = \begin{bmatrix} e^{-3t} & e^{-t} \\ -e^{-3t} & e^{-t} \end{bmatrix}$.

November 30, 2004