

Math 325  
Spring, 2001

## SOLVING FIRST ORDER LINEAR CONSTANT COEFFICIENT EQUATIONS

In section 2.1 of Boyce and DiPrima, you learned how to solve a first order linear ordinary differential equation using an integrating factor (typically called  $\mu$ ). Such an equation has the form  $y' + p(t)y = g(t)$ . This method works for any first order linear ODE.

However, if the equation happens to be constant coefficient and the function  $g$  is of a particularly simple form, there is another way to think about the problem. The equation has the form

$$y' + ay = g(t) \tag{1}$$

where  $a$  is a constant. You can think of this as a special case of an  $n$ th order linear inhomogeneous ODE (with  $n = 1$ ). If you think of it that way, you can solve it the same way you solve higher order constant coefficient linear ODEs. Here's a sketch.

*Step 1* Solve the corresponding homogeneous equation

$$y' + ay = 0 \tag{2}$$

by looking for a solution of the form  $y = Ce^{rt}$ . You find that  $r = -a$ . So the general solution to (??) is

$$y_c = Ce^{-at}.$$

Now, back to the original equation, (??). The general solution will be of the form

$$y = y_c + y_p$$

where  $y_p$  is a particular solution, that is, one solution you will find somehow. Step 2 will apply if  $g(t)$  is of a particularly nice form. Suppose

$$g(t) = p(t)e^{-at}$$

where  $p(t)$  is a polynomial of degree  $k$ .

*Step 2* Use the **method of undetermined coefficients**. Look for a particular solution of the form

$$y_p = t(A_0t^k + A_1t^{k-1} + \dots A_{k-1}t + A_k)e^{-at},$$

that is,  $t$  times a general polynomial of degree  $k$ , with undetermined coefficients which you need to determine, times an exponential. (You need the factor of  $t$  in front because the exponential term solves the homogeneous equation (??).) Plug  $y_p$  into the original equation (??). Then equate corresponding terms. This will give you  $k+1$  equations for the  $k+1$  undetermined coefficients  $A_0, \dots, A_k$ . Solve these equations to determine the coefficients. Now you have found  $y_p$ .

(You can actually handle somewhat more general forms of  $g(t)$ , any form that can be handled for  $n$ th order equations by the method of undetermined coefficients, but this is the form of  $g(t)$  which comes up when you are solving a system of the form  $\mathbf{x}' = J\mathbf{x}$  where  $J$  is a matrix in Jordan canonical form.)

*Step 3* The general solution to (??) is

$$y = Ce^{-at} + t(A_0t^k + A_1t^{k-1} + \dots A_{k-1}t + A_k)e^{-at}$$

where  $A_0, \dots, A_k$  are the coefficients you found in Step 2.