

**Rudin, page 239/17:**

a) The range is all  $\mathbb{R}^2$  except  $(0, 0)$ . Given  $(s, t) \in \mathbb{R}^2$ , just let  $x = \frac{1}{2} \log(s^2 + t^2)$  and choose  $y$  so that  $\cos y = e^{-x}s$  and  $\sin y = e^{-x}t$ . This can be done since  $(e^{-x}s)^2 + (e^{-x}t)^2 = 1$ .

b) The Jacobian of  $f$  is

$$\det \begin{pmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{pmatrix} = e^{2x}(\cos^2 y + \sin^2 y) = e^{2x} \neq 0$$

for any  $(x, y) \in \mathbb{R}^2$ . So  $f$  is locally invertible by the inverse function theorem. However,  $f$  is not globally invertible since it's not injective:  $f(x, y + 2n\pi) = f(x, y)$  for every  $n \in \mathbb{Z}$ .

c)

$$g(s, t) = \left( \frac{1}{2} \log(s^2 + t^2), \tan^{-1}(t/s) \right),$$

(choosing  $\tan^{-1}$  to have range  $(-\pi/2, \pi/2)$ ). Then

$$g'(s, t) = \begin{pmatrix} \frac{s}{s^2+t^2} & \frac{t}{s^2+t^2} \\ \frac{-t}{s^2+t^2} & \frac{s}{s^2+t^2} \end{pmatrix}.$$

So

$$g'(f(x, y)) = \begin{pmatrix} e^{-x} \cos y & e^{-x} \sin y \\ -e^{-x} \sin y & e^{-x} \cos y \end{pmatrix} = f'(x)^{-1}.$$

d) For given  $y$ , the set

$$\{f(x, y) : x \in \mathbb{R}\} = \{e^x(\cos y, \sin y) : x \in \mathbb{R}\}$$

consists of all positive multiples of the fixed vector  $(\cos y, \sin y)$ . That is, the image of a horizontal line is a ray beginning at  $(0, 0)$  (but not including this point).

For given  $x$ , the set

$$\{f(x, y) : y \in \mathbb{R}\} = \{e^x(\cos y, \sin y) : y \in \mathbb{R}\}$$

consists of all points at distance  $e^x$  from  $(0, 0)$ . That is, the image of a vertical line is a circle centered at the origin.