

Multiple Choice

1.(6 pts) If the scalar projection of \mathbf{b} onto \mathbf{a} is $\text{Comp}_{\mathbf{a}} \mathbf{b} = 1$, what is $\text{Comp}_{2\mathbf{a}} 3\mathbf{b}$?

- (a) 2 (b) 5 (c) $\frac{3}{2}$ (d) 6 (e) 3

2.(6 pts) Which of the following expressions gives the length of the curve defined by $\mathbf{r}(t) = \ln(t)\mathbf{i} - t\mathbf{j} + t^2\mathbf{k}$ between the points $(0, -1, 1)$ and $(1, -e, e^2)$?

- (a) $\int_1^e \sqrt{\ln^2(t) + t^2 + t^4} dt$ (b) $\int_1^{e^2} \sqrt{1/t + 1 + 4t^2} dt$
(c) $\int_1^e \sqrt{1/t^2 + 1 + 4t^2} dt$ (d) $\int_1^e \sqrt{1/t - 1 + 2t} dt$
(e) $\int_1^e \sqrt{\ln(t) - t + t^2} dt$

3.(6 pts) Find the distance from the point $(1, 2, 3)$ to the plane $x + 2y - 2z = -7$.

- (a) $\sqrt{3}$ (b) 1 (c) 6 (d) 2 (e) $\sqrt{6}$

4.(6 pts) Suppose the position function $\mathbf{r}(t) = \langle t^3/3, t^2/2, t \rangle$. Find the normal component of the acceleration vector at $t = 1$.

- (a) $a_N = \sqrt{2}$ (b) $a_N = \sqrt{3}$ (c) $a_N = \sqrt{5}$
(d) $a_N = 1$ (e) $a_N = 0$

5.(6 pts) Find the area of the triangle with vertices $(4, 2, 2)$, $(3, 3, 1)$ and $(5, 5, 1)$.

- (a) 0 (b) 4 (c) $\sqrt{3}$ (d) $\sqrt{6}$ (e) 2

6.(6 pts) For $f(x, y) = e^{xy^2}$, calculate the second partial derivative f_{xy} .

- (a) $2y^3e^{xy^2} + 2ye^{xy^2}$ (b) $2xy^3e^{xy^2} + 2ye^{xy^2}$
 (c) $4xy^2e^{xy^2}$ (d) $4xye^{xy^2}$
 (e) $2xye^{xy^2} + 2ye^{xy^2}$

7.(6 pts) Find the vector equation of the line passing through the point $(1, 1, 1)$ and $(1, 2, 3)$

- (a) $\langle x, y, z \rangle = \langle 1, 1, 1 \rangle + t\langle 0, 1, 2 \rangle$
 (b) $\langle x, y, z \rangle = \langle 0, 1, 2 \rangle + t\langle 1, 2, 3 \rangle$
 (c) $\langle x, y, z \rangle = \langle 1, 1, 1 \rangle + t\langle 0, 0, 1 \rangle$
 (d) None of the above
 (e) $\langle x, y, z \rangle = \langle 1, 1, 1 \rangle + t\langle 1, 2, 3 \rangle$

8.(6 pts) Consider a helix curve

$$\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle.$$

Find the equation of the osculating plane of the curve at the point $(0, 0, 0)$

- (a) $x = 0$
 (b) $-y + z = 0$
 (c) None of the above
 (d) $y + z = 0$
 (e) $x + y + z = 0$


9.(6 pts) Given a space curve

$$\mathbf{r}(t) = \langle 2 \cos t, e^t, t \rangle.$$

Which of the following points is in the tangent line of the curve at the point $(2, 1, 0)$?

- (a) (1, 1, 1) (b) (2, 1, 1) (c) (1, 2, 0)
 (d) (2, 2, 1) (e) (0, 1, 2)

10.(6 pts) Which of the following functions has this contour map



- (a) $f(x, y) = xy$ (b) $f(x, y) = y - \frac{xy - 1}{x}$
 (c) $f(x, y) = y - \frac{1}{x^2}$ (d) $f(x, y) = \frac{1}{x}$
 (e) $f(x, y) = \frac{1}{x^2}$

11.(6 pts) Given three points $P(2, 0, 2)$, $Q(1, 1, 0)$ and $R(1, 2, 3)$.

- (a) Find an equation of the plane through P , Q and R .
 (b) Find an equation of the line through the point $(1, 1, 1)$ perpendicular to the plane in part (a).

12.(6 pts) (a) Find an equation for the line of intersection of the planes $x - 3y + 2z = 0$ and $2x - 3y + z = 0$.

(b) Does the line from part (a) intersect the line with equations $x = 1 + t$, $y = 3 - t$, $z = 1 + t$? If so, where do they intersect?

13.(6 pts) A particle has the acceleration

$$\mathbf{a}(t) = 2\mathbf{j} + 6t\mathbf{k}.$$

At the time $t = 0$, the particle's position is at the origin and its velocity is $\mathbf{v}(0) = \mathbf{i} + \mathbf{k}$. Find the position function $\mathbf{r}(t)$ of the particle.

Name: _____

Instructor: ANSWERS

Math 20550, Practice Exam 1
February 19, 2019

- The Honor Code is in effect for this examination. All work is to be your own.
- No calculators.
- The exam lasts for 1 hour and 15 minutes.
- Be sure that your name is on every page in case pages become detached.
- Be sure that you have all 6 pages of the test.
- Each multiple choice question is 6 points. Each partial credit problem is 12 points.
You will receive 4 extra points.

PLEASE MARK YOUR ANSWERS WITH AN X, not a circle!					
1.	(a)	(b)	(c)	(d)	(●)
2.	(a)	(b)	(●)	(d)	(e)
3.	(a)	(b)	(c)	(●)	(e)
4.	(●)	(b)	(c)	(d)	(e)
5.	(a)	(b)	(c)	(●)	(e)
6.	(a)	(●)	(c)	(d)	(e)
7.	(●)	(b)	(c)	(d)	(e)
8.	(a)	(●)	(c)	(d)	(e)
9.	(a)	(b)	(c)	(●)	(e)
10.	(a)	(b)	(●)	(d)	(e)

Please do NOT write in this box.	
Multiple Choice	_____
11.	_____
12.	_____
13.	_____
Extra Points.	<u>4</u> _____
Total:	_____

11.

$$\mathbf{PQ} = (2, 0, 2) - (1, 1, 0) = \langle 1, -1, 2 \rangle$$

$$\mathbf{RQ} = (1, 2, 3) - (1, 1, 0) = \langle 0, 1, 3 \rangle$$

So for part (a) we compute $\mathbf{PQ} \times \mathbf{RQ}$ which is $\langle -3 - 2, -(3 - 0), 1 \rangle = \langle -5, -3, 1 \rangle$. So the equation of the plane through P , Q , and R is

$$\langle -5, -3, 1 \rangle \cdot \langle x, y, z \rangle = \langle -5, -3, 1 \rangle \cdot \langle 1, 1, 0 \rangle$$

$$-5x - 3y + z = -8.$$

For part (b) we use the computations from part (a) to see that the equation of the line should be

$$\langle x, y, z \rangle = \langle 1, 1, 1 \rangle + t\langle -5, -3, 1 \rangle.$$

12. For part (a) we take the cross product of the normal vectors of each plane.

$$\langle 1, -3, 2 \rangle \times \langle 2, -3, 1 \rangle = \langle -3 + 6, -(1 - 4), -3 + 6 \rangle = \langle 3, 3, 3 \rangle.$$

This is the direction of the line. So now we just need a point in the intersection of both planes, since both are of the form $ax + by + cz = 0$ we can easily see that $(0, 0, 0)$ is in the intersection. So the equation for the line of intersection is

$$\langle x, y, z \rangle = t\langle 3, 3, 3 \rangle.$$

For part (b) to see if this line intersects the given line, we must solve a system of equations. Changing the parameter in the given line to s we get

$$1 + s = 3t$$

$$3 - s = 3t$$

$$1 + s = 3t$$

Solving the first equation for s to get $s = 3t - 1$ and plugging into the 2nd equation to get $3 - (3t - 1) = 3t$ we can solve to see $t = 2/3$. Which would mean that $s = 1$. Checking these values in the equation we see that this is a solution $1 + 2 = 3(2/3)$. So using either the $t = 2/3$ or $s = 1$ value we can compute that the intersection point is $(2, 2, 2)$.

13. To find $\mathbf{v}(t)$ we integrate $\mathbf{a}(t)$, to get

$$\mathbf{v}(t) = \langle c, 2t + d, 3t^2 + e \rangle$$

where c , d , and e are constants.

To find the constants we evaluate at 0. To see that $\mathbf{v}(0) = \langle c, d, e \rangle$ which should equal $\langle 1, 0, 1 \rangle$. So $c = 1$, $d = 0$, and $e = 1$.

So now $\mathbf{v}(t) = \langle 1, 2t, 3t^2 + 1 \rangle$.

To find $\mathbf{r}(t)$ we repeat this process,

$$\mathbf{r}(t) = \langle t + c, t^2 + d, t^3 + t + e \rangle.$$

Now using the initial data we have that

$$\langle 0, 0, 0 \rangle = \mathbf{r}(0) = \langle c, d, e \rangle.$$

So $c = 0$, $d = 0$, and $e = 0$.

So the final answer is $\mathbf{r}(t) = \langle t, t^2, t^3 + t \rangle$.