

AME 60611  
Examination 2  
Prof. J. M. Powers  
21 November 2008

1. (20) Given  $x \in \mathbb{R}^1$ ,  $f : \mathbb{R}^1 \rightarrow \mathbb{R}^1$ ,

$$f(x) = e^x, \quad x \in [-1, 1],$$

find the first two terms in a Fourier-Legendre expansion of  $f(x)$ .  
The first two Legendre polynomials are  $P_0(s) = 1$ ,  $P_1(s) = s$ ,  
 $s \in [-1, 1]$ .

2. (20) If  $\mathbf{u} = x\mathbf{i} + y\mathbf{j}$ , evaluate

$$I = \oint_C \mathbf{u}^T \cdot d\mathbf{r},$$

where  $C$  is a single loop around the closed contour described by  
the ellipse,  $x^2 + 2y^2 = 1$ .

3. (20) For  $x \in \overline{\mathbb{L}}_2[0, 1]$ ,  $y \in \overline{\mathbb{L}}_2[0, 1]$ ,  $t \in \mathbb{R}^1$ , find the angle between  
the two functions

$$x(t) = it,$$

$$y(t) = i.$$

4. (20) For  $\mathbf{A} : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ , find the vector  $\mathbf{x} \in \mathbb{R}^3$  of minimum  $\|\mathbf{x}\|_2$   
which minimizes  $\|\mathbf{A} \cdot \mathbf{x} - \mathbf{b}\|_2$  when

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 1 \\ 3 & 6 & 3 \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}.$$

5. (20) Find the singular value decomposition of  $\mathbf{A}$  where

$$\mathbf{A} = (3 \quad 4i \quad 0).$$