

Verified Calculation of Multiscale Combustion in Gaseous Mixtures

Joseph M. Powers

DEPARTMENT OF AEROSPACE & MECHANICAL ENGINEERING

DEPARTMENT OF APPLIED & COMPUTATIONAL MATHEMATICS & STATISTICS

UNIVERSITY OF NOTRE DAME

NOTRE DAME, INDIANA, USA

delivered to

Purdue University

19 January 2012



Acknowledgments

- Samuel Paolucci, U. Notre Dame
- Ashraf al-Khateeb, Khalifa U.
- Zach Zikoski, Post-doctoral student, U. Notre Dame
- Christopher Romick, Ph.D. student, U. Notre Dame
- Tariq Aslam, Los Alamos National Laboratory
- Stephen Voelkel, NSF-REU undergraduate student, U. Notre Dame
- Karel Matouš, U. Notre Dame
- NSF
- NASA
- DOE


Outline

- Part I: Preliminaries
- Part II: Fundamental linear analysis of length scales of reacting flows with detailed chemistry and multicomponent transport. (*with al-Khateeb and Paolucci*)
- Part III: Direct Numerical Simulation (DNS) of complex reacting and inert flows with a) traditional methods, and b) a wavelet-based adaptive algorithm implemented in a massively parallel computing architecture. (*1D detonations with Romick and Aslam; 2D detonations of Zikoski and Paolucci, inert implosions with Voelkel and Romick using Zikoski's algorithm*)

Part I: Preliminaries

Some Semantics

- *Verification*: Solving the equations right—a math exercise.
- *Validation*: Solving the right equations—a physics exercise.
- *DNS*: a verified and validated computation that resolves *all* ranges of relevant continuum physical scales present.



“Research needs for future internal combustion engines,” *Phys. Today*, 2008.

Hypothesis

DNS of fundamental compressible reactive flow fields (thus, detailed kinetics, viscous shocks, multi-component diffusion, etc. are represented, verified, and validated) is on a trajectory toward realization via advances in

- adaptive refinement algorithms, and
- massively parallel architectures.

Corollary I

A variety of modeling compromises, e.g.

- shock-capturing (FCT, PPM, ENO, WENO, etc.),
- implicit chemistry with operator splitting,
- low Mach number approximations,
- turbulence modeling (RANS, $k - \epsilon$, LES, etc.), or
- reduced/simplified kinetics, flamelet models,

need not be invoked *when and if* this difficult goal of DNS is realized; **simple low order explicit discretizations suffice if spatio-temporal grid resolution is achieved.**

Corollary II

Micro-device level DNS is feasible today; macro-device level DNS remains in the distant future.

Corollary III

A variety of challenging fundamental unsteady multi-dimensional compressible reacting flows are now becoming amenable to DNS, especially in the weakly unstable regime; **we would do well as a community to direct more of our efforts towards *unfiltered* simulations so as to more starkly expose the richness of unadulterated continuum scale physics.**

[Example (only briefly shown today): ordinary WENO shock-capturing applied to unstable detonations can dramatically corrupt the long time limit cycle behavior; retention of physical viscosity allows relaxation to a unique dissipative structure in the unstable regime.]

Part II: Fundamental Linear Analysis of Length Scales

Motivation

- To achieve DNS, the interplay between chemistry and transport needs to be captured.
- The interplay between reaction and diffusion length and time scales is well summarized by the classical formula

$$l \sim \sqrt{D\tau}.$$

- Segregation of chemical dynamics from transport dynamics is a prevalent notion in reduced kinetics combustion modeling.
- But, can one rigorously mathematically verify an NS model without resolving the small length scale induced by fast reaction? *Answer: no.*
- Do micro-scales play a role in macro-scale non-linear dynamics? *Answer: in some cases, yes, see Romick, Aslam, & Powers, 2012, JFM, in review.*

Illustrative Linear Model Problem

A linear one-species, one-dimensional unsteady model for reaction, advection, and diffusion:

$$\frac{\partial \psi}{\partial t} + u \frac{\partial \psi}{\partial x} = D \frac{\partial^2 \psi}{\partial x^2} - a\psi,$$
$$\psi(0, t) = \psi_u, \quad \left. \frac{\partial \psi}{\partial x} \right|_{x=L} = 0, \quad \psi(x, 0) = \psi_u.$$

Time scale spectrum

For the spatially homogenous version: $\psi_h(t) = \psi_u \exp(-at)$,

reaction time constant: $\tau = \frac{1}{a} \implies \Delta t \ll \tau.$

Length Scale Spectrum

- The steady structure:

$$\psi_s(x) = \psi_u \left(\frac{\exp(\mu_1 x) - \exp(\mu_2 x)}{1 - \frac{\mu_1}{\mu_2} \exp(L(\mu_1 - \mu_2))} + \exp(\mu_2 x) \right),$$

$$\mu_1 = \frac{u}{2D} \left(1 + \sqrt{1 + \frac{4aD}{u^2}} \right), \quad \mu_2 = \frac{u}{2D} \left(1 - \sqrt{1 + \frac{4aD}{u^2}} \right),$$

$$l_i = \left| \frac{1}{\mu_i} \right|.$$

- For fast reaction ($a \gg u^2/D$):


$$l_1 = l_2 = \sqrt{\frac{D}{a}} = \sqrt{D\tau} \implies \Delta x \ll \sqrt{D\tau}.$$

Spatio-Temporal Spectrum

$$\psi(x, t) = \Psi(t)e^{ikx} \quad \Rightarrow \quad \frac{d\Psi}{dt} = \left(-a \left(1 + \frac{iku}{a} + \frac{Dk^2}{a} \right) t \right) \Psi.$$

$$\Psi(t) = C \exp \left(-a \left(1 + \frac{iku}{a} + \frac{Dk^2}{a} \right) t \right).$$

- For long length scales: $\lim_{k \rightarrow 0} \tau = \lim_{\lambda \rightarrow \infty} \tau = \frac{1}{a},$
 - For fine length scales: $\lim_{k \rightarrow \infty} \tau = \lim_{\lambda \rightarrow 0} \tau = \frac{\lambda^2}{4\pi^2} \frac{1}{D},$
- $\left. \vphantom{\begin{matrix} \lim_{k \rightarrow 0} \tau \\ \lim_{k \rightarrow \infty} \tau \end{matrix}} \right\} \mathcal{S}_t = \left(\frac{2\pi}{\lambda} \sqrt{\frac{D}{a}} \right)^2.$
- Balance between reaction and diffusion at $k \equiv \frac{2\pi}{\lambda} = \sqrt{\frac{a}{D}} = 1/\ell,$



- Similar to $H_2 - air$: $\tau = 1/a = 10^{-8}$ s, $D = 10$ cm²/s,
- $\ell = \sqrt{\frac{D}{a}} = \sqrt{D\tau} = 3.2 \times 10^{-4}$ cm.

Laminar Premixed Flames

Adopted Assumptions:

- One-dimensional,
- Low Mach number,
- Neglect thermal diffusion effects and body forces.

Governing Equations:

$$\begin{aligned}\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial x} (\rho u) &= 0, \\ \rho \frac{\partial h}{\partial t} + \rho u \frac{\partial h}{\partial x} + \frac{\partial j^q}{\partial x} &= 0, \\ \rho \frac{\partial y_l}{\partial t} + \rho u \frac{\partial y_l}{\partial x} + \frac{\partial j_l^m}{\partial x} &= 0, \quad l = 1, \dots, L - 1, \\ \rho \frac{\partial Y_i}{\partial t} + \rho u \frac{\partial Y_i}{\partial x} + \frac{\partial j_i^m}{\partial x} &= \dot{\omega}_i \bar{m}_i, \quad i = 1, \dots, N - L.\end{aligned}$$

- **Unsteady spatially homogeneous** reactive system:

$$\frac{d\mathbf{z}(t)}{dt} = \mathbf{f}(\mathbf{z}(t)), \quad \mathbf{z}(t) \in \mathbb{R}^N, \quad \mathbf{f} : \mathbb{R}^N \rightarrow \mathbb{R}^N.$$

$$\mathbf{0} = (\mathbf{J} - \lambda\mathbf{I}) \cdot \mathbf{v}.$$

$$\mathcal{S}_t = \frac{\tau_{slowest}}{\tau_{fastest}}, \quad \tau_i = \frac{1}{|\operatorname{Re}(\lambda_i)|}, \quad i = 1, \dots, R \leq N - L.$$

- **Steady spatially inhomogeneous** reactive system:

$$\tilde{\mathbf{B}}(\tilde{\mathbf{z}}(x)) \cdot \frac{d\tilde{\mathbf{z}}(x)}{dx} = \tilde{\mathbf{f}}(\tilde{\mathbf{z}}(x)), \quad \tilde{\mathbf{z}}(x) \in \mathbb{R}^{2N+2}, \quad \tilde{\mathbf{f}} : \mathbb{R}^{2N+2} \rightarrow \mathbb{R}^{2N+2}.$$

$$\tilde{\lambda}\tilde{\mathbf{B}} \cdot \tilde{\mathbf{v}} = \tilde{\mathbf{J}} \cdot \tilde{\mathbf{v}}.$$




$$\mathcal{S}_x = \frac{\ell_{coarsest}}{\ell_{finest}}, \quad \ell_i = \frac{1}{|\operatorname{Re}(\tilde{\lambda}_i)|}, \quad i = 1, \dots, 2N - L.$$

Laminar Premixed Hydrogen–Air Flame

- Standard detailed mechanism^a; $N = 9$ species, $L = 3$ atomic elements, and $J = 19$ reversible reactions,
- stoichiometric hydrogen-air: $2H_2 + (O_2 + 3.76N_2)$,
- adiabatic and isobaric: $T_u = 800\text{ K}$, $p = 1\text{ atm}$,
- calorically imperfect ideal gases mixture,
- neglect Soret effect, Dufour effect, and body forces,
- CHEMKIN and IMSL are employed.

^aJ. A. Miller, R. E. Mitchell, M. D. Smooke, and R. J. Kee, *Proc. Combust. Ins.* **19**, p. 181, 1982.




● Unsteady spatially homogeneous reactive system:



$\tau_{slowest} = 1.8 \times 10^{-2} \text{ s}$
 $\tau_{fastest} = 1.0 \times 10^{-8} \text{ s}$

$\left. \begin{array}{l} \tau_{slowest} = 1.8 \times 10^{-2} \text{ s} \\ \tau_{fastest} = 1.0 \times 10^{-8} \text{ s} \end{array} \right\} \mathcal{S}_t \sim \mathcal{O}(10^4).$

• **Steady spatially inhomogeneous** reactive system:^a



$$\left. \begin{array}{l} \ell_{\text{coarsest}} = 2.6 \times 10^0 \text{ cm} \\ \ell_{\text{finest}} = 2.4 \times 10^{-4} \text{ cm} \end{array} \right\} \mathcal{S}_x \sim \mathcal{O}(10^4).$$

^aA. N. Al-Khateeb, J. M. Powers, and S. Paolucci, *Comm. Comp. Phys.* 8(2): 304, 2010.

Spatio-Temporal Spectrum

- PDEs \longrightarrow $2N + 2$ PDAEs,

$$\mathbf{A}(\mathbf{z}) \cdot \frac{\partial \mathbf{z}}{\partial t} + \mathbf{B}(\mathbf{z}) \cdot \frac{\partial \mathbf{z}}{\partial x} = \mathbf{f}(\mathbf{z}).$$

- Spatially homogeneous system at chemical equilibrium subjected to a spatially inhomogeneous perturbation, $\mathbf{z}' = \mathbf{z} - \mathbf{z}^e$,

$$\mathbf{A}^e \cdot \frac{\partial \mathbf{z}'}{\partial t} + \mathbf{B}^e \cdot \frac{\partial \mathbf{z}'}{\partial x} = \mathbf{J}^e \cdot \mathbf{z}'.$$


- Spatially discretized spectrum,

$$\mathcal{A}^e \cdot \frac{d\mathbf{Z}}{dt} = (\mathcal{J}^e - \mathcal{B}^e) \cdot \mathbf{Z}, \quad \mathbf{Z} \in \mathbb{R}^{2N(N+1)}.$$

- The time scales of the generalized eigenvalue problem,

$$\tau_i = \frac{1}{|\operatorname{Re}(\lambda_i)|}, \quad i = 1, \dots, (\mathcal{N} - 1)(N - 1).$$

- $D_{mix} = \frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \mathcal{D}_{ij},$
- $\ell_1 = \sqrt{D_{mix}\tau_s} = 1.1 \times 10^{-1} \text{ cm},$
- $\ell_2 = \sqrt{D_{mix}\tau_f} = 8.0 \times 10^{-4} \text{ cm} \approx \ell_{finest} = 2.4 \times 10^{-4} \text{ cm}.$





Conclusions: Part II

- Time and length scales are coupled.
- Coarse wavelength modes have time scales dominated by reaction.
- Short wavelength modes have time scales dominated by diffusion.
- Fourier modal analysis reveals a cutoff length scale for which time scales are dictated by a balance between **transport** and **chemistry**.
- Fine scales, temporal and spatial, are essential to resolve reacting systems; the finest length scale is related to the finest time scale by $\ell \sim \sqrt{D\tau}$.
- For a $p = 1 \text{ atm}$, $H_2 + \text{air}$ laminar flame, the length scale where fast reaction balances diffusion is $\sim 2 \mu\text{m}$, the necessary scale for a DNS.

Part III: DNS of Complex Reacting and Inert Flows

Effect of Diffusion on Detonation Dynamics: 1D, 1-Step



- Small physical diffusion significantly delays transition to instability.
- In the unstable regime, small diffusion has a large role in determining role for the long time dynamics.
- Romick, Aslam, Powers, *JFM*, in review, 2012.

Effect of Diffusion on Detonation Dynamics: 1D, N -Step

Case Examined

- Romick, Aslam, Powers, AIAA ASM, 2012
- Overdriven detonations with ambient conditions of 0.421 atm and 293.15 K
- Initial stoichiometric mixture of $2H_2 + O_2 + 3.76N_2$
- Detailed kinetics: 9 species, 19 reversible reactions
- a) Inviscid via shock-fitting, and b) viscous (multicomponent diffusion of a viscous, heat conducting fluids) via WAMR studied
- $D_{CJ} \sim 1961 \text{ m/s}$
- Overdrive is defined as $f = D_o^2 / D_{CJ}^2$
- Overdrives of $1.025 < f < 1.150$ were examined

Continuum Scales

- The mean-free path scale is the cut-off minimum length scale associated with continuum theories.
- A simple estimate for this scale is given by Vincenti and Kruger (1967):

$$\lambda = \frac{\bar{M}}{\sqrt{2}\pi\mathcal{N}_A\rho d^2} \sim \mathcal{O}(10^{-6} \text{ cm}) . \quad (1)$$


- The finest reaction length scale is $L_r \sim \mathcal{O}(10^{-4} \text{ cm})$.
- A simple estimate of a viscous length scale is:

$$L_\mu = \frac{\nu}{c} = \frac{6 \times 10^{-1} \text{ cm}^2/\text{s}}{9 \times 10^4 \text{ cm}/\text{s}} \sim \mathcal{O}(10^{-5} \text{ cm}) . \quad (2)$$

- $\lambda < L_\mu < L_r$


Inviscid Steady-State: Pressure

$$f = 1.15$$




Inviscid Transient Behavior: Stable Detonation

$$f = 1.15$$




Near Neutral Stability




Inviscid Transient Behavior: Unstable Detonation

$$f = 1.10$$



- Frequency of 0.97 MHz agrees well with both the frequency, 1.04 MHz , observed by Lehr (*Astro. Acta*, 1972) in experiments and the frequency, 1.06 MHz , predicted by Yungster and Radhakrishnan.
- The maximum detonation front pressure predicted, 13.5 atm , is similar to the value of 14.0 atm found by Daimon and Matsuo.

Validation: Recovering Lehr's High Frequency Instability




Lehr, *Astro. Acta*, 1972

- Experiment of shock-induced combustion in flow around a projectile in an ambient stoichiometric mixture of $2H_2 + O_2 + 3.76N_2$ at 0.421 atm .
- Projectile velocity yields an equivalent overdrive of $f \approx 1.1$
- The observed frequency was approximately 1.04 MHz
- Compare to 1D computation' prediction: 0.97 MHz





Unstable, Inviscid Detonation: $x-t$ Diagram

$$f = 1.10$$







A $x-t$ diagram of density in a Galilean reference frame traveling at 2057 m/s .

Inviscid Transient Behavior: Various Overdrives





Inviscid Phase Portraits: Various Overdrives




Stable, Viscous Detonation: Long Time Structure

$$f = 1.15$$





Stable, Viscous Detonation: Transient Behavior

$$f = 1.15$$




Unstable, Viscous Detonation: Long Time Structure

$$f = 1.10$$



Unstable, Viscous Detonation: Transient Behavior


$$f = 1.10$$



The addition of viscous effects have a stabilizing effect, decreasing the amplitude of the oscillations by $\sim 25\%$.

Unstable, Viscous Detonation: $x-t$ Diagram

$$f = 1.10$$




A $x-t$ diagram of density in a Galilean reference frame traveling at 2057 m/s.

2D Viscous Detonation in Hydrogen-Air

- Wavelet Adaptive Multilevel Representation (WAMR, Zikoski and Paolucci), resolves multi-scale solutions in an adaptive fashion.
- User-defined error control guarantees a verified solution.
- The algorithm has been implemented with an MPI-based domain decomposition in a massively parallel computational architecture with linear scaling to at least 10^3 processors.

2-D VISCOUS DETONATION

Initial Conditions:



Domain: $[0, 60] \times [0, 6]$ cm

Front: $x = 15.0$ cm

Unreacted pocket:

$[1.05 \times 1.43]$ cm

at $x = 14.7$ cm

$P = 4.7 \times 10^5$ dyne/cm²

$T = 2100$ K

128 cores

391 hrs runtime

$2H_2 : O_2 : 7Ar$ mixture
9 species, 37 reactions

Wavelet parameters:


$$\epsilon = 1 \times 10^{-3}$$

$$p = 6, \quad n = 5$$


$$[N_x \times N_y]_{j_0} = [600 \times 60]$$

$$J - j_0 = 10$$

2-D VISCOUS DETONATION (CONT.)



2-D VISCOUS DETONATION (CONT.)



Inert Viscous Cylindrical Implosion

- WAMR algorithm employed
- $100 \mu m \times 100 \mu m$ square domain,
- Pure argon,
- Initial uniform temperature, $T = 300 K$,
- Initial pressure ratio is $4 atm : 0.2 atm$ between argon on either side of an octagonal diaphragm,
- $T_{max}(r = 0, t \sim 40 ns) \sim 2400 K$.

Conclusions

- Verified 1D and 2D combustion physics spanning over five orders of magnitude—from near mean-free path scales (10^{-4} *cm*) to small scale device scales (10 *cm*)—can be calculated today with modern adaptive algorithms working within a massively parallel computing architecture.
- Micro-scale viscous shock dynamics can influence oscillatory detonation dynamics on the macro-scale.
- Some 1D detonations can be validated; others await 3D extension.
- Realization of verified and validated DNS would remove the need for common, but problematic, modeling assumptions (shock-capturing, turbulence modeling, implicit chemistry with operator splitting, reduced kinetics/flamelets).
- Such 3D V&V could be viable in an exascale environment; however, routine desktop DNS calculations remain difficult to envision at macro-device scales.