

The Elusive *closo*-Ge₁₀²⁻ Zintl Ion: Finally “Captured” as a Ligand in the Complex [Ge₁₀Mn(CO)₄]³⁻

Daniel Rios and Slavi C. Sevov*

Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, Indiana 46556

Received May 4, 2010

Although often seen in mass spectra, the otherwise elusive *closo*-Ge₁₀²⁻ Zintl ion has been finally structurally characterized as coordinated to manganese in [Ge₁₀Mn(CO)₄]³⁻ made by the reaction of *nido*-Ge₉⁴⁻ with Mn₂(CO)₁₀ in ethylenediamine. The new cluster is a bicapped square antiprism, as expected for a *closo*-deltahedron of 10 atoms.

Fragmentation of the well-known nine-atom *nido*-deltahedral Zintl ions of group 14, Ge₉⁴⁻, Sn₉⁴⁻, and Pb₉⁴⁻, is not an uncommon phenomenon, especially for the heavier congeners, and is manifested by the observed reassembly of the fragments into larger clusters. Thus, the reactions of Pb₉⁴⁻ clusters with [Pt(PPh₃)₄] and Ni(COD)₂ result in 12-atom icosahedra, [Pt@Pb₁₂]²⁻, and 10-atom bicapped prisms, [Ni@Pb₁₀]²⁻, respectively.^{1,2} Similarly, the reactions of nine-atom Sn₉⁴⁻ clusters with Ni(COD)₂ and [(COD)IrCl]₂ lead to the 17-atom species [Ni₂Sn₁₇]⁴⁻ with a complex structure and the icosahedral [Ir@Sn₁₂]³⁻ (after oxidation of the initially formed and structurally characterized [Sn₉Ir(COD)]³⁻), respectively.^{3–5} The recent synthesis of ferrocene-like [Fe@Ge₁₀]³⁻ and [Co@Ge₁₀]³⁻ from Ge₉⁴⁻ ions showed that germanium is not immune to such fragmentations either.^{6,7} What is common in all these reactions is that they all contain a central transition metal, and the product is not a nine-atom cluster or a multiple of it as in, for example, [Ni₃Ge₁₈]⁴⁻ and [Pd₂Ge₁₈]⁴⁻.^{8,9} The mechanisms for the formation of these species most likely involve some fragmentation of the nine-atom clusters, perhaps

driven by nucleation of the larger clusters around the central transition metals.¹⁰ However, the successful synthesis and characterization of *closo*-Pb₁₀²⁻ by soft oxidation of Pb₉⁴⁻ with Au(I) salt indicates that such a central atom is not always necessary.¹¹ This is further supported by the observed formation of heteroatomic nine-atom clusters Ge_{9-x}Sn_x⁴⁻ (*x* = 0–9) from the corresponding homoatomic Ge₉⁴⁻ and Sn₉⁴⁻, which presumably involves fragmentation and reassembly as well.¹²

The same mechanism is most likely responsible for the formation of the 10-atom germanium species often observed in our electrospray mass spectra. Thus, the reactions of Ge₉⁴⁻ clusters (derived from the intermetallic precursor K₄Ge₉) with aldehydes and ketones in dimethylformamide (DMF) exhibit almost exclusively a peak at *m/z* = 727 which corresponds to Ge₁₀⁻.¹³ Clearly, some oxidation and fragmentation of Ge₉⁴⁻ must take place in such reactions. However, our numerous attempts to crystallize and structurally characterize this elusive 10-atom cluster by employing different solvents and cation sequestering agents had been unsuccessful until now. Here, we report the isolation and structure of the deltahedral *closo*-Ge₁₀²⁻ Zintl ion as a ligand in the manganese tetracarbonyl complex [Ge₁₀Mn(CO)₄]³⁻ (Figure 1).

The new compound, [K(2,2,2-crypt)]₃[Ge₁₀Mn(CO)₄]·tol, was crystallized as dark purple blocks from the soft oxidation of Ge₉⁴⁻ in an ethylenediamine solution of K₄Ge₉ with Mn₂(CO)₁₀ in the presence of 2,2,2-crypt. Some of the purple crystals were dissolved in DMF, and the resulting light purple solution was used for carrying out electrospray mass spectrometry in the negative-ion mode. The spectra (in the Supporting Information) showed not only the [Ge₁₀Mn(CO)₄]⁻ anion at *m/z* = 894 but also its derivatives after the loss of one or more carbon monoxide ligands in the process of solvent removal, i.e., [Ge₁₀Mn(CO)₃], [Ge₁₀Mn(CO)₂], [Ge₁₀Mn(CO)], and [Ge₁₀Mn] at *m/z* = 866, 838, 810, and 782, respectively.

*To whom correspondence should be addressed. E-mail: ssevov@nd.edu.

(1) Esenturk, N. E.; Fettinger, J.; Lam, Y.; Eichhorn, B. *Angew. Chem., Int. Ed.* **2004**, *43*, 2132.

(2) Esenturk, N. E.; Fettinger, J.; Eichhorn, B. *Chem. Commun.* **2005**, 247.

(3) Esenturk, N. E.; Fettinger, J.; Eichhorn, B. *J. Am. Chem. Soc.* **2006**, *128*, 12.

(4) Downing, D. O.; Zavalij, P.; Eichhorn, B. W. *Eur. J. Inorg. Chem.* **2010**, 890.

(5) Wang, J. Q.; Stegmaier, S.; Wahl, B.; Fässler, T. F. *Chem.—Eur. J.* **2010**, *16*, 1793.

(6) Zhou, B.; Denning, S. M.; Kays, L. D.; Goicoechea, J. M. *J. Am. Chem. Soc.* **2009**, *131*, 2802.

(7) Wang, J. Q.; Stegmaier, S. D.; Fässler, T. F. *Angew. Chem., Int. Ed.* **2009**, *121*, 2032.

(8) Goicoechea, J. M.; Sevov, S. C. *Angew. Chem., Int. Ed.* **2005**, *44*, 4026.

(9) Goicoechea, J. M.; Sevov, S. C. *J. Am. Chem. Soc.* **2005**, *127*, 7676.

(10) Fässler, T. F.; Hoffmann, S. D. *Angew. Chem., Int. Ed.* **2004**, *43*, 6242.

(11) Spiekermann, A.; Hoffmann, S. D.; Fässler, T. F. *Angew. Chem., Int. Ed.* **2006**, *45*, 3459.

(12) Gillett-Kunnath, M. M.; Petrov, I.; Sevov, S. C. *Inorg. Chem.* **2010**, *49*, 721.

(13) Schneeberger, P. M. In *Functionalization and Reactivity of Germanium Clusters*, M.S. Thesis, University of Notre Dame, Notre Dame, IN, 2009.




Figure 1. The complex anion $[\text{Ge}_{10}\text{Mn}(\text{CO})_4]^{3-}$ is composed of a deltahedral *closo*- Ge_{10}^{2-} Zintl ion coordinated as a ligand to a $\text{Mn}(\text{CO})_4^-$ fragment to form an 18-electron Mn complex ($d_{\text{Ge-Mn}} = 2.335 \text{ \AA}$). The Ge_{10}^{2-} ion is a bicapped square antiprism with average Ge–Ge distances (\AA) of 2.575 at the capping atoms, 2.547 between the staggered squares, and 2.812 within the squares.

Although a much earlier report claims the synthesis of Ge_{10}^{2-} ,¹⁴ the structure reported there is so greatly disordered that even the nuclearity of the cluster is uncertain, and the cluster has been widely viewed to be the nine-atom *closo*- Ge_9^{2-} .¹⁵

Ge_{10}^{2-} is a 10-atom deltahedron with 42 electrons (four from each germanium and two from the charge), and as such it satisfies Wade-Mingos' rule for a *closo*-deltahedron, which prescribes $4n + 2$ electrons for *closo*-deltahedra with n vertices. Its geometry, as expected for a 10-atom *closo*-deltahedron, is that of a bicapped square antiprism with pseudo- D_{4d} symmetry. The distances with the cluster are very similar to other analogous species with 10 vertices and the same electron count, e.g., $[\text{Ge}_9\text{Zn-Ph}]^{3-}$ with the same geometry (Zn is one of the capping atoms) and 42 cluster electrons (9×4 from Ge + 2 from Zn + 1 from Ph + 3 from charge).¹⁶ Three types of distances can be defined depending on the types of atoms they connect, and their averages in Ge_{10}^{2-} are as follows: 2.575 \AA at the capping atoms, 2.547 \AA between the staggered squares, and 2.812 \AA within the squares. For comparison, the corresponding distances in $[\text{Ge}_9\text{Zn-Ph}]^{3-}$ are 2.564 (at the only one Ge capping atom), 2.551, and 2.804 \AA .

In $[\text{Ge}_{10}\text{Mn}(\text{CO})_4]^{3-}$, the *closo*- Ge_{10}^{2-} cluster coordinates to the Mn center via the lone pair of one of the capping atoms of the bicapped square antiprism and thus acts as a normal two-electron ligand. From an organometallic point of view, the Mn center itself achieves an 18-electron configuration by combining the electrons from five ligands (four CO and one cluster), the metal center with seven electrons, and one negative charge, i.e., $[(\text{Ge}_{10}^{2-})\text{Mn}(\text{CO})_4]^-$ (HOMOs are d_{xy} and $d_{x^2-y^2}$ on Mn, nearly degenerate). Thus, the -3 charge of the

complex can be viewed as distributed between the cluster and the Mn center. The Mn–Ge distance of 2.3347(8) \AA cannot be compared with that in a similar anionic complex because there are none available. The only available related compound is the 19-electron complex $(\text{CF}_3)_3\text{GeMn}(\text{CO})_5$, and its Ge–Mn distance of 2.4132(9) \AA is understandably quite longer.¹⁷

The infrared spectrum of the new compound (crystals in Nujol) showed the C–O stretching modes at very low frequencies of 1892, 1789, and 1775 cm^{-1} . As expected, they are lower than those in neutral species such as $\text{HMn}(\text{CO})_5$ (2094, 2001, and 1970 cm^{-1}) and $\text{Mn}_2(\text{CO})_{10}$ (2044, 2013, and 1983 cm^{-1}),^{18,19} but they are also lower than the frequencies of even the anionic $[\text{Mn}(\text{CO})_5]^-$ (1893 and 1860 cm^{-1}) and $[(\text{Ph}_3\text{P})\text{Mn}(\text{CO})_4]^-$ (1940, 1846, and 1815 cm^{-1}).²⁰ This suggests very strong electron donation from the negatively charged cluster to the Mn center and subsequently to the carbonyls' π^* orbitals. This is supported further by the much shorter Mn–C and much longer C–O distances in $[\text{Ge}_9\text{-Mn}(\text{CO})_4]^{3-}$, 1.784 and 1.161 \AA (averaged), respectively, when compared with both the neutral and the anionic species: $d_{\text{Mn-C}}/d_{\text{C-O}}(\text{av.}) = 1.847/1.123$, $1.846/1.130$, $1.798/1.156$, and $1.796/1.159 \text{ \AA}$ in $\text{HMn}(\text{CO})_5$,²¹ $\text{Mn}_2(\text{CO})_{10}$,²² $[\text{Mn}(\text{CO})_5]^-$,²³ and $[(\text{Ph}_3\text{P})\text{Mn}(\text{CO})_4]^-$,²⁴ respectively.

All operations were carried out under an inert atmosphere or vacuum using standard Schlenk-line or glovebox techniques. Ethylenediamine (Alfa-Aesar, 99%) was distilled over sodium metal and stored in a gastight ampule under nitrogen in the glovebox. Dried toluene was stored over molecular sieves in the glovebox. 2,2,2-Crypt (4,7,13,16,21,24-hexaoxa-1,10-diazabicyclo-[8.8.8]-hexacosane, Acros, 98%) and $\text{Mn}_2(\text{CO})_{10}$ (Aldrich, 99%) were used as received after carefully drying them under a vacuum. K_4Ge_9 was synthesized by heating a stoichiometric mixture of the elements (K: 99+%, Strem; Ge: 99.999%, Alfa Aesar) at 900 $^\circ\text{C}$ for 2 days in sealed niobium containers jacketed in evacuated fused silica ampules. K_4Ge_9 (70 mg, 0.086 mmol) and 2,2,2-crypt (129 mg, 0.344 mmol) were dissolved in 2 mL of ethylenediamine (red solution) in a test tube inside a glovebox. $\text{Mn}_2(\text{CO})_{10}$ (130 mg, 0.112 mmol) was added, and the reaction mixture was stirred for an additional 3 h. The resulting dark purple solution was filtered from the substantial amount of precipitate (most likely clusters oxidized to neutral Ge) and layered with toluene. It yielded nicely shaped black-purple block crystals of $[\text{K}(2,2,2\text{-crypt})]_3[\text{Ge}_{10}\text{Mn}(\text{CO})_4] \cdot \text{tol}$ after 2 weeks (ca. 46% crystalline yield).

Single-crystal X-ray diffraction data of $[\text{K}(2,2,2\text{-crypt})]_3[\text{Ge}_{10}\text{Mn}(\text{CO})_4] \cdot \text{tol}$ were collected on a Bruker APEX diffractometer with a CCD area detector and Mo $\text{K}\alpha$ radiation at 100 K. The structure was solved by direct methods and was refined on F^2 using the SHELXTL V5.1 package. Crystal data: monoclinic, $P2_1/n$, $a = 15.3123(6)$, $b = 20.9401(8)$, and

(17) Brauer, J. D.; Eujen, R. *Organometallics* **1983**, *2*, 263.

(18) Rest, A. J.; Turner, J. J. *J. Chem. Soc., Chem. Commun.* **1969**, 375.

(19) Flitcroft, N.; Huggins, D. K.; Kaesz, H. D. *Inorg. Chem.* **1964**, *3*, 1123.

(20) Faltyněk, R. A.; Wrighton, M. S. *J. Am. Chem. Soc.* **1978**, 2701.

(21) La Placa, S. J.; Hamilton, W. C.; Ibers, J. A.; Davison, A. *Inorg. Chem.* **1969**, *8*, 1928.

(22) Churchill, M. R.; Amoh, K. N.; Wasserman, H. J. *Inorg. Chem.* **1981**, *20*, 1609.

(23) Corraine, M. S.; Lai, C. K.; Zhen, Y.; Churchill, M. R.; Buttrey, L. A.; Ziller, J. W.; Atwood, J. D. *Organometallics* **1992**, *11*, 35.

(24) Riley, O. E.; Davis, R. E. *Inorg. Chem.* **1980**, *19*, 159.

(14) Belin, C.; Mercier, H.; Angilella, V. *New J. Chem.* **1991**, *15*, 931.

(15) Fässler, T. F.; Hunziker, M.; Spahr, M.; Lueken, H. *Z. Anorg. Allg. Chem.* **2000**, *626*, 692.

(16) Goicoechea, J. M.; Sevov, S. C. *Organometallics* **2006**, *25*, 4530.

Communication

$c = 28.0514(11) \text{ \AA}$, $\beta = 92.818(2)^\circ$, $V = 8983.6(6) \text{ \AA}^3$, $Z = 4$, $R1/wR2 = 4.12/9.51\%$ for the observed data ($I > 2\sigma(I)$), $R1/wR2 = 6.56/10.53\%$ for all data.

Acknowledgment. This material is based upon work supported by the Department of Energy, Office of Basic Energy Sciences, under Award Number DE-FG02-

07ER46476. We thank M. M. Gillett-Kunnath for helping with the IR spectra.

Supporting Information Available: X-ray crystallographic file in CIF format and electrospray mass spectrum in negative ion mode of $[\text{K}(2,2,2\text{-crypt})]_3[\text{Ge}_{10}\text{Mn}(\text{CO})_4] \cdot \text{tol}$ are available free of charge via the Internet at <http://pubs.acs.org>.