

Ample Vector Bundles and Branched Coverings

Thomas Peternell Andrew J. Sommese

Dedicated to Robin Hartshorne on his 60th birthday

Introduction

Let $f : X \rightarrow Y$ be a finite surjective morphism of degree d between n -dimensional connected projective manifolds X and Y . Let $\mathcal{E} := [(f_*\mathcal{O}_X)/\mathcal{O}_Y]^*$. It is a beautiful theorem of Lazarsfeld [La80] that when Y is \mathbb{P}^n , then for any such f and X , the vector bundle $\mathcal{E}(-1)$ is spanned. There have been many partial generalizations of this result for homogeneous Y , e.g., [De95, De96, Ki96a, Ki96b, Ma97, KM98]. In this article we investigate spannedness and ampleness properties for not necessarily homogeneous Y .

\mathcal{E} has a number of tendencies towards positivity. For example, using relative duality $\mathcal{E} \cong f_*\omega_{X/Y}/\mathcal{O}_Y$, and thus \mathcal{E} is weakly positive in the sense of Viehweg, e.g., [Vi95]. In particular this implies the nefness of \mathcal{E} for Y a curve. We also prove this for Y an abelian surface. Moreover, $\mathcal{E} \otimes \mathcal{E}$ always has a nontrivial section, coming from a natural linear map from \mathcal{E}^* to \mathcal{E} induced by f , which is an isomorphism over a dense Zariski open set of Y , see Lemma (1.13).

Though it follows from the weak positivity that \mathcal{E} is nef when $\dim Y = 1$, it is easy to construct examples with \mathcal{E} not even nef. Indeed, if f expresses X as a double cover of Y , then $\mathcal{E} \cong L$ where the discriminant locus is an element of $|2L|$. Thus if Y possesses a nonnef line bundle with some even power having a smooth section, then we can construct a covering with \mathcal{E} not nef.

Even when \mathcal{E} is nef, ampleness can fail in a number of different ways. For example, if Y possesses an unramified finite cover $f : X \rightarrow Y$ of degree $d > 1$, then \mathcal{E} is nef with no sections, but with all Chern classes zero (see Theorem (1.20)). Thus the best possible result for ampleness would be, “if \mathcal{E} is nef, and f does not factor through an unramified covering of Y , then \mathcal{E} is ample.” The first place to check this result is for curves, since in this case \mathcal{E} is always nef. It is, in fact, true for $Y = \mathbb{P}^1$ by Lazarsfeld’s result, and for Y an elliptic curve by a result of Debarre [De96]. Our Theorem (2.1) shows that this result holds for arbitrary smooth curves.

Unfortunately, in dimensions ≥ 2 , the double cover construction shows that \mathcal{E} can fail to be ample, even “if \mathcal{E} is nef, and f does not factor through an unbranched covering of Y ”. Some of the restrictions imposed by the hypothesis that \mathcal{E} is ample for all covers are given in Theorem (1.27) and Corollary (1.28). For example, the only smooth connected projective surfaces S , which stand any chance of having the property that for all finite surjective morphisms from connected projective surfaces

to S , the associated bundle \mathcal{E} is ample, are \mathbb{P}^2 ; a minimal $K3$ -surface with no -2 curves; or a surface with an ample canonical bundle, for which there are neither nontrivial unramified covers over it, nor nonfinite morphisms from it onto positive dimensional varieties.

In §3, we show in Theorem (3.2) that \mathcal{E} is always spanned for Y a smooth Del Pezzo surface with $K_Y^2 \geq 5$. We show in Theorem (3.3) that \mathcal{E} is spanned over the Hirzebruch surfaces \mathbb{F}_r , under the added condition that if $r \geq 2$, then the section E , of the projection $p : Y \rightarrow \mathbb{P}^1$, with negative self-intersection does not lie in the discriminant locus of the covering. We also show in Theorem (3.7) that \mathcal{E} is ample over the Hirzebruch surfaces \mathbb{F}_r , under the further condition that the general fiber F , of the projection $p : Y \rightarrow \mathbb{P}^1$, has connected inverse image under the given covering. The conditions are also shown to be necessary.

In §4, we show in Theorem (4.2) that \mathcal{E} is always spanned for Y a smooth Del Pezzo manifold of dimension $n \geq 3$ when $b := H^n \geq 5$. Here by definition $-K_Y = (n - 1)H$ for some ample line bundle H on Y . We also show Theorem (4.3), which states that $h^0(\mathcal{E}) \neq 0$ under very mild conditions, i.e., if b times the degree of the covering is at least five. Finally we ask when $h^0(\mathcal{E} \otimes (-H)) \neq 0$. Using adjunction theory, we show in Theorem (4.7) and (4.10) that this is so for branched covers of Del Pezzo manifolds, except in very rare circumstances.

For our induction arguments, we were forced to consider branched covers over manifolds Y with domains X merely normal Cohen-Macaulay with finite irrational locus, i.e., for each point x of X , the local rings of germs of holomorphic (or algebraic) functions at x are Cohen-Macaulay and normal, with the set of nonrational singularities of X finite. One interesting aspect of our work is that many results, e.g., Lazarsfeld's original result, hold at this level of generality.

We dedicate this paper to Robin Hartshorne. His work has significantly enriched algebraic geometry. It has also been a stimulus for our work: indeed, a main result of this paper is based primarily on Robin Hartshorne's beautiful characterization [Ha71] of ample vector bundles on curves.

We thank the Department of Mathematics and the Duncan Chair of the University of Notre Dame for making our collaboration possible. We also thank Meeyoung Kim and Robert Lazarsfeld for helpful comments on this paper. In particular, Robert Lazarsfeld suggested using Castelnuovo-Mumford regularity in Theorem 1.17 to improve our original result about nefness of \mathcal{E} off the branch locus. He also suggested Problem 1.19 and then found the elegant solution, which appears as an appendix to this paper. The second author thanks the Department of Mathematics of the K.T.H. (Royal Institute of Technology) of Stockholm, Sweden, where this paper was finished.

Contents

1	General results on branched coverings	3
2	Coverings of curves	13

3	Coverings over surfaces	14
4	Coverings over Del Pezzo Manifolds	18

1 General results on branched coverings

In this paper we work over \mathbb{C} . By a variety we mean a complex analytic space, which might be neither reduced or irreducible. Let $f : X \rightarrow Y$ be a surjective finite morphism between n -dimensional projective manifolds. Let $\mathcal{E} := [(f_*\mathcal{O}_X)/\mathcal{O}_Y]^*$. Here are some of the known results on the structure of \mathcal{E} .

1. If $Y = \mathbb{P}^n$, then \mathcal{E} is ample and spanned by Lazarsfeld [La80].
2. If Y is a quadric, then \mathcal{E} is spanned by [Ki96b] and \mathcal{E} is ample if $3 \leq n \leq 6$ by [KM98].
3. If Y is a Grassmannian, then \mathcal{E} is spanned, and k -ample [Ki96a], and in fact ample [Ma97].
4. By [KM98], it follows that if Y is a Lagrangian Grassmannian, then \mathcal{E} is ample; and that for some other homogeneous Y , \mathcal{E} is spanned.
5. \mathcal{E} is ample if Y is an elliptic curve and if f does not factor through an étale map [De96].

We need generalizations of Lazarsfeld's theorem for mildly singular X .

The general setup of these results carries over to Cohen-Macaulay X . Note that reduced curves have at worst Cohen-Macaulay singularities.

Setup 1.1 *Let $f : X \rightarrow Y$ be a finite, degree d , morphism from a reduced pure n -dimensional projective variety X onto an n -dimensional projective manifold Y . Assume that X has at worst Cohen-Macaulay singularities. Since f is flat, $f_*(\mathcal{O}_X)$ is locally free of rank d . By the trace map, $f_*(\mathcal{O}_X)$ has a canonical direct summand \mathcal{O}_Y ; therefore we can define a locally free sheaf \mathcal{E}_f of rank $d - 1$ by setting*

$$f_*(\mathcal{O}_X) = \mathcal{E}_f^* \oplus \mathcal{O}_Y.$$

When the map f is obvious from the context, we usually denote the sheaf \mathcal{E}_f as \mathcal{E} .

We make the above singularity assumption on X , not because we want to state the most general theorems, but rather because we will need the more general choice of X in induction procedures. We will often make the added assumption that X is normal with the irrational locus $\mathcal{I}(X)$ finite, i.e., with the locus of nonrational singularities finite. This assumption guarantees that the crucial vanishing theorems are still valid [So85].

We need the following slight variant of the results in [So85].

Lemma 1.2 *Let X be a normal n -dimensional projective variety with at worst Cohen-Macaulay singularities. Assume that the irrational locus of X is finite. Assume that L is ample and D is an effective (possibly empty) reduced Cartier divisor. Assume that either:*

1. $n \leq 2$; or
2. D is normal with finite irrational locus.

Then $h^i(-L - D) = 0$ for $i < n$.

Proof. If D is empty, this is part of the Kodaira vanishing theorem of [So85]. For $n = 1$, the result follows immediately from Serre duality.

Consider the sequence:

$$0 \rightarrow \omega_X \otimes L \rightarrow \omega_X \otimes L \otimes \mathcal{O}_X(D) \rightarrow \omega_D \otimes L_D \rightarrow 0.$$

If $n = 2$ then by Serre duality we have $h^1(\omega_D \otimes L_D) = 0$ and thus using [So85], $h^i(\omega_X \otimes L \otimes \mathcal{O}_X(D)) = 0$ for $i \geq 1$. This shows the lemma when $n = 2$.

If $n > 2$ then we obtain the result by using the results of [So85] on D and X .
Q.E.D.

It is conjectured that \mathcal{E} is always ample if Y is rational homogeneous with $b_2 = 1$ [KM98] or if Y is simple abelian and f does not factor through an étale map [De96].

First we make a technically very useful generalization of Lazarsfeld's theorem to the Cohen-Macaulay case.

Theorem 1.3 *Let $f : X \rightarrow \mathbb{P}^n$ be a finite surjective morphism from a pure dimensional reduced projective variety X with at worst Cohen-Macaulay singularities. Assume that either $n = 1$ or that X is normal with finite irrational locus. Then \mathcal{E} is spanned. If further X is either irreducible or a connected curve, then $\mathcal{E}(-1)$ is spanned, and, in particular, \mathcal{E} is ample.*

Proof. First assume that $n = 1$. Then the spannedness of \mathcal{E} will follow if $h^1(\mathcal{E}(-1)) = 0$. We have

$$h^1(\mathcal{E}(-1)) = h^0(\mathcal{E}^*(-1)) = h^0((f_*\mathcal{O}_X)(-1)) = h^0(f^*\mathcal{O}_{\mathbb{P}^1}(-1)) = 0.$$

The spannedness of $\mathcal{E}(-1)$ will follow from $h^1(\mathcal{E}(-2)) = 0$. We have

$$h^1(\mathcal{E}(-2)) = h^0(\mathcal{E}^*) = h^0(f_*\mathcal{O}_X) - 1 = h^0(\mathcal{O}_X) - 1.$$

Now note that if X is connected, then $h^0(\mathcal{O}_X) = 1$.

To show that \mathcal{E} is spanned for a given $n > 1$, we assume by induction that the result is true for dimensions $\leq n - 1$. Choose an arbitrary point $y \in \mathbb{P}^n$. Choose a general $D \in |\mathcal{O}_{\mathbb{P}^n}(1)|$ passing through $y \in \mathbb{P}^n$. $\overline{D} := f^{-1}(D)$ is reduced, with at worst

Cohen-Macaulay singularities. By induction the sheaf $\mathcal{E}_{f_{\overline{D}}}$ is spanned. Moreover the irrational locus is finite. Then by [So85] (see Lemma (1.2)) it follows that

$$H^1(\mathcal{E}(-1)) = H^{n-1}(X, f^* \mathcal{O}_{\mathbb{P}^n}(-n)) = 0,$$

and thus the map $H^0(\mathcal{E}) \rightarrow H^0(\mathcal{E}_D) \rightarrow 0$. Finally note that $\mathcal{E}_{f_{\overline{D}}} \cong \mathcal{E}_D$ (see (1.5)).

To show that $\mathcal{E}(-1)$ is spanned for a given $n > 1$, we assume by induction that the result is true for dimensions $\leq n - 1$. Choose an arbitrary point $y \in \mathbb{P}^n$. Choose a general $D \in |\mathcal{O}_{\mathbb{P}^n}(1)|$ passing through $y \in \mathbb{P}^n$. $\overline{D} := f^{-1}(D)$ is reduced, with at worst Cohen-Macaulay singularities. Moreover [So86], \overline{D} is irreducible if $\dim D \geq 2$ and connected if $\dim \overline{D} = 1$. Then by [So85] it follows that

$$H^1(\mathcal{E}(-2)) = H^{n-1}(X, f^* \mathcal{O}_{\mathbb{P}^n}(-n + 1)) = 0.$$

Thus the map $H^0(\mathcal{E}(-1)) \rightarrow H^0(\mathcal{E}_D(-1)) \rightarrow 0$. Now use $\mathcal{E}_{f_{\overline{D}}}(-1) \cong \mathcal{E}_D(-1)$.
Q.E.D.

By the same argument as in Theorem (1.3) we obtain

Theorem 1.4 *Let $f : X \rightarrow Q_n$ be a finite surjective morphism from a pure dimensional reduced projective variety X with at worst Cohen-Macaulay singularities onto a smooth quadric Q_n . Assume further that X is normal with finite irrational locus. Then \mathcal{E} is spanned.*

We will use (and have already used) the following easy fact several times.

Lemma 1.5 *Let $f : X \rightarrow Y$ be a finite surjective morphism from a pure dimensional reduced projective variety X , which is locally a complete intersection, onto a smooth projective manifold Y . Let $C \subset Y$ be a smooth (or a locally complete intersection) curve. Then, letting $X_C := f^{-1}(C)$, $\mathcal{E}|_C = (f|_{X_C})_*(\omega_{X_C/C})/\mathcal{O}_C$.*

Proof. Notice that X_C might not be smooth, however $X_C \subset X$ is locally a complete intersection, so that $\omega_{X/Y}$ makes good sense. Then our claim follows immediately from the adjunction formulas for $C \subset Y$ and $X_C \subset X$ and from the relation for the normal bundles $N_{X_C|X} = (f|_C)^* N_{C|Y}$.
Q.E.D.

Remark 1.6 Let \mathcal{GS} (respectively \mathcal{GS}') denote the set of all projective manifolds Y with the property that given a finite surjective morphism $f : X \rightarrow Y$ from a pure dimensional projective manifold X (respectively normal Cohen-Macaulay variety X with finite irrational locus) to Y , $H^0(Y, \mathcal{E}_f)$ spans \mathcal{E}_f over a dense Zariski open set of Y . Then it follows that given $Y \in \mathcal{GS}$ (respectively \mathcal{GS}'), $A \times Y \in \mathcal{GS}$ (respectively \mathcal{GS}') for A either projective space or the smooth quadric. This follows by the same induction used above, and the fact that given a semiample bundle L on a pure n -dimensional normal irreducible Cohen-Macaulay projective variety X with $\mathcal{I}(X)$ finite, we have the vanishing $H^j(K_X + L) = 0$ for $j > n - \kappa(L)$, where $\kappa(L)$ is the Kodaira dimension of L .

Remark 1.7 Let $f : X \rightarrow Y$ be a finite morphism from a reduced pure n -dimensional Cohen-Macaulay variety X onto a connected projective manifold Y . There are many instances when \mathcal{E}_f is spanned. In this case, what can we say about the k -ampleness of \mathcal{E}_f in the sense of [So78]? Clearly \mathcal{E} is $(n - 1)$ -ample if and only if X is connected. To see this, just note that since \mathcal{E}_f is assumed spanned, it follows that if it is not k -ample, then there exists an irreducible subvariety $Z \subset Y$ of dimension at least $k + 1$ and a trivial summand of $\mathcal{E}_{f|Z}$. Thus $h^0([f_*\mathcal{O}_X]_Z) \geq 2$. So if \mathcal{E}_f is not $(n - 1)$ -ample, we have $h^0(\mathcal{O}_X) = h^0(f_*\mathcal{O}_X) \geq 2$, which combined with X being reduced, shows that X is not connected. The same argument implies the following “criterion” for ampleness of \mathcal{E}_f when $\dim Y = 2$: “if \mathcal{E}_f is spanned, then \mathcal{E}_f is ample if the restriction of \mathcal{E}_f to the branch locus of f is ample and if inverse images under f of irreducible curves are always connected.” The key point of the argument is that if $\dim Y = 2$, then an irreducible and reduced curve C on Y is Cartier. Thus C' , the inverse image of C under f , is Cartier. Therefore, C' is Cohen-Macaulay (since effective Cartier divisors on a Cohen-Macaulay variety are Cohen-Macaulay). If C is not contained in the ramification locus of f , it follows that C' is generically reduced, and therefore (since C' is Cohen-Macaulay) it is reduced. Thus we conclude that $h^0(\mathcal{E}_{f|C}^*) = 0$ if and only if C' is connected.

Here is a general result giving “generic semipositivity” of \mathcal{E} , under the assumptions in (1.1) plus the assumption that X is smooth.

Basic Observation 1.8 *Let $f : X \rightarrow Y$ be a finite surjective morphism from a pure dimensional reduced projective variety X , which is locally a complete intersection, onto a smooth projective manifold Y . $f_*(\omega_{X/Y}) \simeq \mathcal{E} \oplus \mathcal{O}_Y$.*

Proof. Notice first that $f_*(\omega_{X/Y})$ is locally free of rank d , which follows immediately from the flatness of f via standard cohomology theorems. Then by relative duality

$$f_*(\omega_{X/Y})^* \simeq f_*(\mathcal{O}_X)$$

which gives the claim. Q.E.D.

Our purpose is to generally investigate how “positive” the vector bundle \mathcal{E} could be. To state a general result, we recall the following definition due to Viehweg.

Definition 1.9 Let Y be a projective manifold and \mathcal{F} a locally free sheaf on Y . Then \mathcal{F} is *weakly positive* if and only if for all ample line bundles H and all positive integers a the bundle $S^{ab}\mathcal{F} \otimes H^b$ is generically spanned for $b \gg 0$.

To make this definition a little more suggestive we weaken this notion in the following form

Definition 1.10 We say that \mathcal{F} is *generically nef*, if there exists a countable union $A = \cup_i A_i$ of proper analytic sets in X such that $\mathcal{F}|C$ is nef for all curves $C \not\subset A$.

For a discussion of generically nef bundles we refer to [DPS99]. We clearly have the following result.

Proposition 1.11 *If \mathcal{F} is weakly positive, then \mathcal{F} is generically nef.*

Theorem 1.12 *Let $f : X \rightarrow Y$ be a finite surjective morphism between projective manifolds. Then the bundle \mathcal{E} is weakly positive, in particular it is generically nef.*

Proof. Having in mind (1.5), this is just a special case of a result in [Vi82]. Q.E.D.

We shall see below that “in general” \mathcal{E} will be neither ample nor spanned nor even nef, even if Y is a surface, e.g., due to the existence of (-1) -curves. Nevertheless there is some tendency towards \mathcal{E} having sections. In (1.22) below we see by a very simple argument that $\mathcal{E}|_C$ is nef for the general smooth complete intersection curve C .

Lemma 1.13 *Let $f : X \rightarrow Y$ be a finite surjective morphism between projective manifolds. Then, there is an exact sequence*

$$0 \rightarrow \mathcal{E}^* \rightarrow \mathcal{E} \rightarrow \mathcal{S} \rightarrow 0,$$

with the reduced support of \mathcal{S} equal to the image under f of the ramification divisor $B \subset X$. Thus $h^0(\mathcal{E} \otimes \mathcal{E}) \geq 1$ and there is a section of $2 \det \mathcal{E}$ with zero set equal set theoretically to $f(B)$.

Proof. Since the higher direct images under f of coherent sheaves are 0, the direct image of the exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow \omega_{X/Y} \rightarrow \mathcal{O}_X(B) \rightarrow 0,$$

gives the desired exact sequence.

Since the map from \mathcal{E}^* to \mathcal{E} is an isomorphism over a Zariski dense open set, we conclude that we have a sheaf map from $-\det \mathcal{E}$ to $\det \mathcal{E}$, with cokernel having support equal to the image under f of the ramification divisor $B \subset X$. Thus $2 \det \mathcal{E}$ has a section which vanishes on the set theoretic image $f(B)$ under f of the ramification divisor $B \subset X$. Q.E.D.

There is some hope for ampleness if $\text{Pic}(Y) = \mathbb{Z}$.

Corollary 1.14 *Let $f : X \rightarrow Y$ be a finite surjective morphism between n -dimensional projective manifolds. Assume that f is not an unramified cover. If the Picard number $\rho(Y) = 1$, then $\det \mathcal{E}$ is ample.*

Proof. By Lemma (1.13), $2 \det \mathcal{E}$ has a section which vanishes on the set theoretic image $f(B)$ under f of the ramification divisor $B \subset X$. Since f is not an unramified cover, $f(B)$ is nonempty, and $\det \mathcal{E}$ is ample. Q.E.D.

Remark 1.15 Using the relative Riemann-Roch theorem it follows that given a finite degree d surjective morphism between n -dimensional projective manifolds,

$$c_1(\mathcal{E}) = \frac{f_*(\omega_{X/Y})}{2} = \frac{f_*(B)}{2},$$

where B is the branch locus of f . Similarly, the higher Chern classes of \mathcal{E} can be computed. For example, when $n = 2$, $c_2(\mathcal{E}) = d\chi(\mathcal{O}_Y) - \chi(\mathcal{O}_X) + \frac{(\omega_Y + \det \mathcal{E}) \cdot \det \mathcal{E}}{2}$.

Next we collect some trivial but basic results on the cohomology of \mathcal{E} which will be used again and again.

Proposition 1.16 *Let $f : X \rightarrow Y$ be a finite morphism from a pure dimensional projective variety X with at worst Cohen-Macaulay singularities onto a projective manifold Y . Then*

1. $h^q(X, \omega_{X/Y}) = h^q(Y, \mathcal{E}) \oplus H^q(Y, \mathcal{O}_Y)$.
2. $h^0(Y, \mathcal{E}) = h^0(X, \omega_{X/Y}) - 1$
3. $H^q(X, \mathcal{O}_X) = H^q(Y, \mathcal{E}^*) \oplus H^q(Y, \mathcal{O}_Y)$.
4. *If Y is Fano and X satisfies the additional conditions of being normal with finite irrational locus, then $H^q(Y, \mathcal{E}) = 0$ for $q \geq 1$.*
5. *If Y is Fano, and X satisfies the additional conditions of being normal with finite irrational locus, and $-\omega_X = rH$ with $r \geq 2$ (so X has index at least 2), then $H^q(Y, \mathcal{E}(-H)) = 0$ for $q \geq 1$.*

Proof. (1), (2) and (3) being clear, (4) and (5) follow by Kodaira's vanishing theorem. Q.E.D.

Notice that (5) implies that if $h^q(\mathcal{O}_X) > h^q(\mathcal{O}_Y)$ for some $1 \leq q \leq n - 1$, then \mathcal{E} cannot be Nakano positive. We will investigate \mathcal{E} mostly on Fano manifolds of large index and in low dimensions. Here is one result of more general nature.

Theorem 1.17 *Let $f : X \rightarrow Y$ be a finite surjective morphism from the projective manifold X of dimension n to the abelian variety Y . Then \mathcal{E}_f is nef.*

Proof. This is an easy adoption of the proof of [DPS94, 3.21]. Adopting the arguments of (ibid.(ii)) word by word, we need to prove the following.

Let G be a very ample line bundle on Y . Then $f_*(\omega_X) \otimes G^{n+1}$ is spanned.

This follows immediately from noting that $f_*(\omega_X) \otimes G^{n+1}$ is Castelnuovo-Mumford 0-regular. Indeed, let y be a point of Y . Choose n sections s_1, \dots, s_n

of G , whose scheme-theoretic zero set includes y as a nonsingular component. Let K^\bullet denote the tensor the Koszul complex associated to the section $s_1 \oplus \cdots \oplus s_n$ of $G^{\oplus n}$ with $f_*(\omega_X) \otimes G$. Applying Kodaira vanishing to the hypercohomology of K^\bullet shows that $f_*(\omega_X) \otimes G^{n+1}$ is spanned.

Then introduce $\lambda : Y \rightarrow Y, z \rightarrow 2z$, make the base change λ^p and argue as in the second part of [DPS94]. Q.E.D.

Corollary 1.18 *If Y is the product of an abelian variety and projective spaces, then \mathcal{E} is nef.*

Robert Lazarsfeld suggested the following problem based on the original version of Theorem (1.17). Later he found the positive solution to in, which is presented in an appendix to this paper.

Problem 1.19 *Let $f : X \rightarrow Y$ be a finite surjective morphism between n -dimensional projective manifolds. Is it true that \mathcal{E}_f is always nef modulo the discriminant locus D , i.e., $\mathcal{E}|_C$ is nef for all curves $C \not\subset D$?*

The following summarizes the general things that we know in the unramified case.

Theorem 1.20 *Let $f : X \rightarrow Y$ be a finite unramified morphism from a connected n -dimensional projective manifold X onto a connected n -dimensional projective manifold Y . Then*

1. $\mathcal{E} \cong \mathcal{E}^*$ and $h^0(\mathcal{E}) = 0$;
2. \mathcal{E} is nef;
3. the i -th Chern class $c_i(\mathcal{E})$ vanishes for $i \geq 1$; and
4. \mathcal{E} has a filtration by hermitian flat vector bundles.

Proof. Since $\omega_X \cong f^*\omega_Y$, we have that $\omega_{X/Y} \cong \mathcal{O}_X$. Taking direct images we conclude that

$$\mathcal{E}^* \oplus \mathcal{O}_Y \cong f_*(\mathcal{O}_X) \cong f_*(\omega_{X/Y}) \cong \mathcal{E} \oplus \mathcal{O}_Y,$$

and therefore that $\mathcal{E}^* \cong \mathcal{E}$.

To see that $h^0(\mathcal{E}) = 0$, simply note that

$$h^0(\mathcal{E}) = h^0(\mathcal{E}^*) = h^0(\mathcal{O}_X) - h^0(\mathcal{O}_Y) = 0.$$

To see nefness of \mathcal{E} , it is equivalent to show nefness of $(f_*\mathcal{O}_X)^*$. By definition this comes down to showing that there is no finite morphism $g : C \rightarrow Y$ of an irreducible and reduced projective curve C to Y with the property that there is a surjection $g^*[V^*] \rightarrow \mathcal{L}^* \rightarrow 0$ of the pullback under g of the vector bundle V^* associated $(f_*\mathcal{O}_X)^*$

onto \mathcal{L}^* a line bundle on C with $\deg \mathcal{L} > 0$. We argue by contradiction. Assume otherwise that there is such a surjection, or equivalently that there is an injection

$$0 \rightarrow \mathcal{L} \rightarrow g^*(f_*\mathcal{O}_X).$$

We have an inner product defined on $f_*\mathcal{O}_X$ by using the trace. Given two elements s, t of the fiber of V at $y \in Y$, define $(s, t) = \text{tr}(s\bar{t})$, where, letting $\mathcal{I}_{f^{-1}(y)}$ denote the ideal sheaf of the fiber of f over y , we identify s, t with the corresponding elements of $\mathcal{O}_X/\mathcal{I}_{f^{-1}(y)}$. Define the function $h : V \rightarrow \mathbb{R}$ by sending $s \in V$ to (s, s) . This function is a plurisubharmonic exhaustion of V , since f is unramified. In particular the restriction of h to \mathcal{L} is a nonconstant plurisubharmonic function on \mathcal{L} . But, since \mathcal{L} has positive degree, any plurisubharmonic function on \mathcal{L} is constant. This contradiction shows the nefness of \mathcal{E} .

Since the metric (s, t) is flat, i.e., its curvature form vanishes, we conclude that the Chern classes $c_i(\mathcal{E})$ of \mathcal{E} with $i > 0$ vanish. For the last statement we refer to [DPS94]. Q.E.D.

Corollary 1.21 *Let $f : X \rightarrow Y$ be a finite morphism from a connected n -dimensional projective manifold X onto a connected n -dimensional projective manifold Y . Then f is unramified if and only if $c_1(\mathcal{E}) = 0$.*

Proof. By Theorem (1.20), we can assume without loss of generality that f is ramified. If $n = 1$, then the result follows from the following Lemma.

Lemma 1.22 *Let $f : X \rightarrow Y$ be a finite morphism between smooth connected curves. Let τ denote the degree of the ramification divisor of f . Then $f_*(\omega_{X/Y})$ and \mathcal{E} are nef, and $\deg f_*(\omega_{X/Y}) = \frac{\tau}{2}$; in particular $\det(\mathcal{E})$ is ample unless f is an unramified cover.*

Proof. The nefness follows from (1.12). Let $c := \deg f_*(\omega_{X/Y}) = \deg \mathcal{E}$. By Riemann-Roch we have

$$\chi(f_*(\omega_{X/Y})) = d(1 - g(Y)) + c,$$

where g denotes the genus. On the other hand Riemann-Roch on X gives

$$\begin{aligned} \chi(f_*\omega_{X/Y}) &= \chi(\omega_{X/Y}) = \chi(\omega_X \otimes f^*(\omega_Y^{-1})) \\ &= -\chi(f^*(\omega_Y^{-1})) = g(X) - 1 - d(2g(Y) - 2). \end{aligned}$$

Putting things together and using the Riemann-Hurwitz formula, the claim follows. Q.E.D.

Thus we can assume that f is ramified and also that $n \geq 2$. In this case let C be a smooth curve on Y obtained as the intersection of $n-1$ general elements of $|L|$, where L is a very ample line bundle on Y . By Bertini's theorem, C and $X_C := f^{-1}(C)$ are smooth. By Kodaira's vanishing theorem, X_C is connected. Since X_C is the intersection of ample divisors, it meets the ramification divisor nontrivially. This gives that $\deg(f_*(\omega_{X_C/C})/\mathcal{O}_C) > 0$. Since $f_*(\omega_{X_C/C})/\mathcal{O}_C \cong \mathcal{E}_C$, we conclude the contradiction that $c_1(\mathcal{E}) \neq 0$. Q.E.D.

Proposition 1.23 *Suppose that $f : X \rightarrow Y$ factors through an unramified cover: $f = b \circ a$ with $b : Z \rightarrow Y$ unramified of degree at least 2. Then \mathcal{E} cannot be ample and moreover \mathcal{E} is not spanned at any point.*

Proof. In fact, $\mathcal{E}_b = (b_*(\mathcal{O}_Z)/\mathcal{O}_Y)^*$ is numerically flat with no sections by Theorem (1.20). Since \mathcal{E}_b is a direct summand of \mathcal{E} , the claims follow. Q.E.D.

Corollary 1.24 *Let $f : X \rightarrow Y$ be a finite surjective morphism between smooth connected n -dimensional projective manifolds. Let r denote the rank of the quotient of \mathcal{E} by the subsheaf of \mathcal{E} generated by the images of elements of $H^0(\mathcal{E})$. If f factors through an unramified cover $b : Z \rightarrow Y$ from a connected projective manifold Z onto Y , then the sheet number of b is bounded by $r + 1$. In particular, if global sections of \mathcal{E} span \mathcal{E} at some point, then*

1. f does not factor through any nontrivial unramified cover of Y ; and
2. given a smooth C obtained as the intersection of general elements $A_i \in |L_i|$, for $n - 1$ ample and spanned line bundles L_1, \dots, L_{n-1} , we have \mathcal{E}_C is ample.

Probably the generic spannedness of \mathcal{E} is unneeded in the next result.

Corollary 1.25 *Let $f : X \rightarrow Y$ be a finite surjective morphism between smooth connected n -dimensional projective manifolds. Assume that global sections of \mathcal{E} span \mathcal{E} at some point, and let C be the smooth curve obtained as the intersection of general elements $A_i \in |L_i|$, for $n - 1$ ample and spanned line bundles L_1, \dots, L_{n-1} . Then the fundamental group of $f^{-1}(C)$ surjects onto the fundamental group of C .*

Proof. First note that $C' := f^{-1}(C)$ is smooth by Bertini's theorem, and connected by the Kodaira vanishing theorem. Since $\mathcal{E}_{C'} \cong f_{C'*}(\omega_{C'/C})/\mathcal{O}_{C'}$, by Lemma (1.5), $f_{C'}$ does not factor through an unramified cover of C . This is equivalent to having the fundamental group of C' surject onto the fundamental group of C . Q.E.D.

Remark 1.26 We would like to give some examples that show that \mathcal{E} , even when ample, may have very few or no sections. Let L be a line bundle on a projective manifold Y . Assume that there exists a section s of L with a smooth zero set D . Assume that A is a line bundle on Y with $rA = L$. Let $f : X \rightarrow Y$ be the r -sheeted cover obtained by taking the r -th root of s . In this case $\mathcal{E} \cong \bigoplus_{j=1}^{r-1} jA$. Thus if we choose A and L such that A is not spanned, we have an example with \mathcal{E} not spanned. Such examples are plentiful. For example,

1. We could take L as the line bundle associated to two distinct points on a curve Y . If $h^1(\mathcal{O}_Y) > 0$, then any line bundle A with $2A = L$, is of degree one, and thus has at most one section. A dimension count shows that if $h^1(\mathcal{O}_Y) \geq 2$, we can choose the two points so that in fact $h^0(A) = 0$;

2. If Y is a Del Pezzo surface, i.e., if $-\omega_Y$ is ample, then it follows by a straightforward argument using Reider's theorem, that if $kA = L$ with $h^0(L) \geq 1$, A not spanned, and $k \geq 2$, then $L = -k\omega_Y$, $A = -\omega_Y$, and $\omega_Y \cdot \omega_Y = 1$.
3. There is a similar example for Del Pezzo threefolds. Let Y denote a Del Pezzo threefold with $-\omega_Y = 2H$ with $H^3 = 1$. Then $-\omega_Y$ is spanned, but for the double cover associated to $2H = -\omega_Y$, $\mathcal{E} = H$ is spanned except at one point.

Similar examples show that \mathcal{E} does not have to be nef. Let Y be a Hirzebruch surface F_r with $r \geq 2$, i.e., a \mathbb{P}^1 -bundle over \mathbb{P}^1 with a section E satisfying $E \cdot E = -r \leq -2$. Let f denote a fiber of the tautological surjection of F_r to \mathbb{P}^1 . Take L as the line bundle $rE + r(r-1)f$. Note that since $(r-1)[E + rf]$ is spanned we can choose a smooth divisor $D' \in |(r-1)[E + rf]|$. Since $E \cdot D' = 0$, $D := E + D'$ is a smooth divisor $\in |rE + r(r-1)f|$. Taking the r -sheeted cyclic cover associated to the r -th root of D , we have

$$\mathcal{E} \cong \bigoplus_{j=1}^{r-1} jA,$$

where A the line bundle associated to $E + (r-1)f$. Note that this bundle is negative restricted to E , and so \mathcal{E} is not even nef in this case.

It is worth emphasizing that it is hard for \mathcal{E} to be ample for all branched covers over a fixed base.

Theorem 1.27 *Let Y be an n -dimensional connected projective manifold. Assume that for all finite surjective morphisms $f : X \rightarrow Y$ from connected projective manifolds X , the associated bundle \mathcal{E} is ample. Then:*

1. *the profinite completion of the fundamental group of Y is 0, and thus in particular the first betti number of Y is 0; and*
2. *there exist no surjective morphisms $g : Y \rightarrow Z$ with $\dim Z \geq 1$, Z projective, and g having at least one positive dimensional fiber.*

Proof. The first assertion is clear. To see the second assertion, let L be a very ample line bundle on Z . Choosing the double cover associated to a smooth $D \in |2g^*L|$, we have $\mathcal{E} \cong g^*L$, which, by the hypothesis on g , cannot be ample. Q.E.D.

Corollary 1.28 *Let Y be an n -dimensional connected projective manifold. Assume that for all finite surjective morphisms $f : X \rightarrow Y$ from connected projective manifolds X , the associated bundle \mathcal{E} is ample. Then either $-\omega_Y$ is ample with $\text{Pic}(Y) = \mathbb{Z}$ or ω_Y is nef. Moreover ω_Y is ample if ω_Y is big.*

Remark 1.29 Using the prediction of the Abundance Conjecture that mK_Y is spanned for suitable large m whenever K_Y is nef, it follows in (1.28), that either $-\omega_Y$ is ample or ω_Y is ample or $\omega_Y \equiv 0$. In the last case Y is an irreducible Calabi-Yau or symplectic manifold. In particular, if $\dim Y = 2$, then Y is the projective plane or ω_Y is ample or Y is K3 without (-2) -curves. If $\dim Y = 3$, then Y is Fano with $b_2 = 1$ or ω_Y is ample or Y is Calabi-Yau without any contraction.

2 Coverings of curves

In this section we prove a result generalizing the theorems of Lazarsfeld [La80] for \mathbb{P}^1 and Debarre [De96] for elliptic curves.

Theorem 2.1 *Let $f : X \rightarrow Y$ be a finite morphism from a smooth connected projective curve X onto a smooth connected projective curve Y . The bundle $\mathcal{E} = [(f_*\mathcal{O}_X)/\mathcal{O}_Y]^*$ is ample if and only if f does not factor through a nontrivial unramified covering of Y .*

Remark 2.2 As is pointed out in (1.26), even when \mathcal{E} is ample,

1. if $h^1(\mathcal{O}_Y) > 0$, then \mathcal{E} does not have to be spanned; and
2. if $h^1(\mathcal{O}_Y) \geq 2$, then $h^0(\mathcal{E})$ can equal 0.

When $h^1(\mathcal{O}_Y) = 1$, then $h^1(\mathcal{E}) = h^0(\mathcal{E}^*) = 0$ and hence by the Riemann-Roch theorem, and Lemma (1.22),

$$h^0(\mathcal{E}) = \deg(\mathcal{E}) = h^1(\mathcal{O}_X) - 1.$$

The rest of the section is devoted to the proof of Theorem (2.1).

By (1.23) we may assume that f does not factor through an unramified covering. We assume that \mathcal{E} is not ample, and argue by contradiction. We have the following simple corollary of Hartshorne's characterization of ampleness of vector bundles on curves [Ha71].

Lemma 2.3 *Let V be a nef vector bundle on a smooth connected curve C . There exists a unique maximal ample vector subbundle A . The quotient V/A has degree zero.*

Proof. If V is ample the Lemma is trivially true. Thus we can assume that V is not ample.

By Hartshorne's characterization of ampleness, if V is not ample, then there exists a vector subbundle A of V with $\deg V/A \leq 0$. Since V is nef, V/A is nef also, and we conclude that $\deg V/A = 0$. If A is the 0 dimensional subbundle, we have $\deg V = 0$, and the Lemma is proven. Indeed, if there was a nontrivial ample subbundle A' in this case, we would have $\deg V/A' = \deg V - \deg A' < 0$, which contradicts the nefness of V . Thus we can assume that the degree of V is positive.

Thus let A be a vector subbundle of V of minimal rank with the property that V/A has degree zero. We will be done if we show that

1. A is ample; and
2. A contains any other ample subbundle of V .

To see the first assertion, note that if A is not ample, then by Hartshorne's theorem, there is a vector subbundle A' of A with $\deg A/A' \leq 0$. Thus $\deg V/A' = \deg V/A + \deg A/A' \leq 0$ contradicting the choice of A with minimal rank.

To see the second assertion, assume that there was an ample subbundle B of V with $B \not\subset A$. Then the saturation of the image of the sheaf of germs of sections of B in the sheaf of germs of sections of V/A is an ample subbundle of V/A . The inverse image of this bundle in V is an ample vector subbundle of V containing A , but the quotient of V by this bundle has negative degree. This contradicts the nefness of V . Q.E.D.

Let \mathcal{F} be the ample subbundle of \mathcal{E} given by Lemma (2.3). Clearly this bundle is also the maximal ample subbundle of $\mathcal{O}_X \oplus \mathcal{E}$.

Let $T := \mathcal{E}/\mathcal{F}$. We claim that $\mathcal{O}_Y \oplus T^*$ is a subring of $f_*\mathcal{O}_X = \mathcal{O}_Y \oplus \mathcal{E}^*$. Indeed, we have that T^* is nef since T is nef and of degree zero. Therefore, we have that $(\mathcal{O}_Y \oplus T^*) \otimes (\mathcal{O}_Y \oplus T^*)$ is nef, and that B , the saturation of its image in $\mathcal{O}_Y \oplus \mathcal{E}^*$, is nef. Since $\mathcal{O}_Y \oplus T^* \subset B$, the quotient $B/(\mathcal{O}_Y \oplus T^*)$ is a nef subbundle of the bundle \mathcal{F}^* . But since \mathcal{F} is ample, we conclude that $B = T^*$ and $\mathcal{O}_Y \oplus T^* \subset f_*(\mathcal{O}_X)$ is a subring.

Now consider the analytic spectrum

$$Z := \text{Specan}(\mathcal{O}_Y \oplus T^*).$$

It comes along with a finite map $b : Z \rightarrow Y$, and the inclusion $\mathcal{O}_Y \oplus T^* \rightarrow \mathcal{O}_Y \oplus \mathcal{E}^*$ gives a map

$$a : \text{Specan}(\mathcal{O}_Y \oplus \mathcal{E}^*) = X \rightarrow Z$$

such that $f = b \circ a$. Since T^* equals its saturation in \mathcal{E}^* , we conclude that Z is normal and hence smooth. Since $\deg T = 0$, we conclude that b is unramified. Q.E.D.

3 Coverings over surfaces

Lemma 3.1 *Let $f : X \rightarrow Y$ be a finite morphism from an irreducible normal Cohen-Macaulay projective variety X , with a finite irrational locus, onto a projective manifold Y of dimension $n \geq 2$. Let H be a line bundle on Y such that $-\omega_Y - H$ is nef and big. Then $h^1(\mathcal{E} \otimes H^*) = 0$.*

Theorem 3.2 *Let $f : X \rightarrow Y$ be a finite morphism from an irreducible normal Gorenstein projective surface X onto a smooth projective surface Y . Let Y be a Del Pezzo surface with $\omega_Y^2 = 9 - r \geq 5$ (i.e., Y is a smooth quadric, or the plane blown up in $r \leq 4$ points in sufficiently general position). Then \mathcal{E} is spanned.*

Proof. Since we already know this for the quadric, we can assume that we have a blowing up map $p : Y \rightarrow \mathbb{P}^2$. We have

$$\omega_Y = -3p^*H + \sum_{i=1}^r E_i$$

the E_i are smooth \mathbb{P}^1 s on Y with self-intersection -1 . We will identify H with $p^*(H)$.

First let $r = 1$. Let $A = 2H - E_1$. Note that $-\omega_Y - A = H$ is nef and big, and thus that $h^1(\mathcal{E}(-A)) = 0$ (3.1). We thus have $H^0(\mathcal{E}) \rightarrow H^0(\mathcal{E}_C) \rightarrow 0$. Since A is very ample, a general $C \in |A - y|$ for any $y \in Y$ is smooth. For a general $C \in |A - y|$, we conclude that $f^{-1}(C)$ is reduced. Since smooth $C \in |A|$ have genus 0, we conclude by Theorem (1.3) that \mathcal{E}_C is spanned, hence \mathcal{E} is spanned.

Now let $r \geq 2$. Let $H_i = H - E_i$. Note that $-\omega_Y - H_i = 2H - \sum_{j \neq i} E_j$ is nef and big. Since H_i is spanned with smooth $C \in |H_i|$ having genus 0 we conclude that \mathcal{E} is spanned by global sections over a generic smooth $C \in |H_i|$. Thus \mathcal{E} is spanned at all points $y \in Y$ where there is a smooth $C \in |H_i - y|$ for some i . The set of $y \in Y$ for which this is not true for a fixed i is $B_i := \cup_{j \neq i} (E_j \cup E_{ij})$ where E_{ij} is the proper transform under p of the line through $\{p(E_i), p(E_j)\}$. Note that $\cap_i B_i = \emptyset$ unless $r = 2$.

In the case $r = 2$ we have that $B_1 \cap B_2 = E_{12}$. Note that $-\omega_Y - H = 2H - E_1 - E_2$ is nef and big, and therefore we can use H in place of H_i . This shows spannedness except at the points $y_1 := E_1 \cap E_{12}$ and $y_2 := E_2 \cap E_{12}$. Thus if we show that

$$H^0(\mathcal{E}) \rightarrow H^0(\mathcal{E}_{E_{12}}) \rightarrow 0,$$

then we will be done. To see this note that $-\omega_Y - (E_i + E_{12})$ is nef and big. Therefore using the Koszul complex and (3.1)

$$0 \rightarrow \mathcal{E} \otimes (-E_1 - E_{12} - E_2) \rightarrow (\mathcal{E} \otimes (-E_1 - E_{12})) \oplus (\mathcal{E} \otimes (-E_2 - E_{12})) \rightarrow \mathcal{E} \otimes (-E_{12}) \rightarrow 0,$$

we are reduced to showing that $h^2(\mathcal{E} \otimes (-E_1 - E_{12} - E_2)) = 0$. But we have

$$h^2(\mathcal{E} \otimes (-E_1 - E_{12} - E_2)) = h^0(\mathcal{E}^* \otimes (-2H + E_1 + E_2)) = h^0(f^*(-2H + E_1 + E_2)) = 0.$$

Q.E.D.

To get spannedness over Hirzebruch surfaces, it is enough to assume that the branch locus does not contain any components of the inverse image of E .

Theorem 3.3 *Let $f : X \rightarrow Y$ be finite surjective morphism from a pure dimensional, normal variety X with at worst Cohen-Macaulay singularities onto a smooth Hirzebruch surface $Y := \mathbb{F}_r$ (with surjection $p : Y \rightarrow \mathbb{P}^1$). Let E denote a section of p with $E \cdot E = -r$. If $r \leq 1$; or $r > 1$ and E does not belong to the discriminant variety of f , then \mathcal{E} is spanned.*

Proof. By Theorem (3.2), we may assume without loss of generality that $r \geq 2$.

Let $\overline{E} := f^{-1}(E)$. Since E does not belong to the discriminant variety of f , it follows that \overline{E} is reduced. Let $d := \deg(f)$ be the degree of f .

Note that global sections of \mathcal{E} span \mathcal{E}_E . To see this, observe that by Theorem (1.3), we need only show that $h^1(\mathcal{E}(-E)) = 0$. But this is equal to $h^1(\mathcal{E}^* \otimes \omega_Y \otimes \mathcal{O}_Y(E))$. Since $-\omega_Y - E$ is ample, we conclude that $h^1(\omega_Y \otimes \mathcal{O}_Y(E)) = 0$ and $h^1(f^*(\omega_Y + E)) = 0$. Therefore

$$h^1(\mathcal{E}(-E)) = h^1(f^*(\omega_Y + E)) = 0.$$

Given an arbitrary fiber F of p we see that \mathcal{E}_F is spanned since global sections span it at $F \cap E$. Thus we will be done if we show that $h^1(\mathcal{E}(-F)) = 0$. But this is equal to $h^1(\mathcal{E}^* \otimes \omega_Y \otimes \mathcal{O}_Y(F))$. Since $h^1(\omega_Y + F) = 0$ we are reduced to showing that $h^1(f^*(\omega_Y + F)) = 0$. Since $f^*(\omega_Y + F) = -f^*(E + (r+1)F) - f^*E$, $f^*(E + (r+1)F)$ is ample, and $f^{-1}(E)$ is reduced, we have vanishing by Lemma (1.2). Q.E.D.

Remark 3.4 Theorem (3.5) is sharp for $r \geq 2$. Indeed consider the cyclic branched cover of \mathbb{F}_r with discriminant locus given by $E + (r-1)(E+rF)$.

We have not pursued the question of what is the “minimal” twist, which makes \mathcal{E} spanned, but such questions can be answered for special classes of varieties by the same techniques. Here is a simple example.

Corollary 3.5 *Let $f : X \rightarrow Y$ be a finite morphism from an irreducible normal projective surface X onto smooth Hirzebruch surface $Y := \mathbb{F}_r$. If $r \geq 1$, then $\mathcal{E} \otimes \mathcal{O}_Y((r-1)F)$ is spanned.*

Proof. Let $p : Y \rightarrow \mathbb{P}^1$ be the \mathbb{P}^1 -bundle projection of Y onto \mathbb{P}^1 . Let F be a fiber of p and let E be the section of Y with $E \cdot E = -r$. To see that $\mathcal{E} \otimes \mathcal{O}_Y((r-1)F)$ is spanned, let $y \in Y$ be a point. A general $C \in |E + (r+1)F|$ passing through y is smooth and does not belong to the branch locus. Therefore, it suffices by Theorem 1.3 to show that $H^1(\mathcal{E} \otimes \mathcal{O}_Y((r-1)F) \otimes \mathcal{O}_Y(-C)) = 0$. But the latter group is isomorphic to $H^1(f^*\mathcal{O}_Y(-E - rF))$, and is therefore zero since $E + rF$ is nef and big. Q.E.D.

For higher dimensional projective bundles, it is not so easy to prove spannedness theorems under natural conditions. There is one special case for which the simple technique of splitting the anticanonical divisor works. Since we do not need this in the rest of the paper, we leave the proof to the reader.

Theorem 3.6 *Let $f : X \rightarrow Y$ be finite surjective morphism from a pure dimensional, normal variety X with finite irrational locus and at worst Cohen-Macaulay singularities onto Y , a \mathbb{P}^{n-1} -bundle over \mathbb{P}^1 .*

1. *If $-\omega_Y$ is nef, then \mathcal{E} is spanned except if $Y = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1}^{\oplus(n-1)} \oplus \mathcal{O}_{\mathbb{P}^1}(2))$, in which case \mathcal{E} is spanned outside the exceptional divisor $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}^{\oplus(n-1)})$.*
2. *If $Y = \mathbb{P}(V)$ with $V := \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^1}(a_{n-1})$ and with $\deg V \leq 1 + a_{n-1}$, then \mathcal{E} is generically spanned.*

The bundle \mathcal{E} will fail to be ample if any curve on Y has disconnected inverse image under f . The examples in (1.26) show that the following is optimal.

Theorem 3.7 *Let $f : X \rightarrow Y$ be a finite surjective morphism from a pure dimensional, normal variety X with at worst Cohen-Macaulay singularities onto a smooth Hirzebruch surface $Y := \mathbb{F}_r$ (with surjection $p : Y \rightarrow \mathbb{P}^1$). Let E denote a section of*

p with $E \cdot E = -r$. and let F denote a general fiber of p . If $f^{-1}(F)$ is connected and $f^{-1}(E)$ is connected and reduced, then $H^0(\mathcal{E}(-E))$ spans $\mathcal{E}(-E)$ on all of Y and $H^0(\mathcal{E}(-F - E))$ spans $\mathcal{E}(-F - E)$ on the complement of E . From this it follows that \mathcal{E} is ample.

Proof. We first show that $\mathcal{E}(-E)$ is spanned by global sections. Let C be a general element of $|E + (r + 1)F|$ that contains y . Since $E + (r + 1)F$ is very ample, C is smooth and does not belong to the discriminant variety of f . Thus $\overline{C} := f^{-1}(C)$ is reduced. It is also connected since it is an ample divisor on X . Thus $\mathcal{E}_C(-1)$ is spanned by Theorem (1.3). The assertion that $\mathcal{E}(-E)$ is spanned by global sections, will therefore follow if we show that $h^1(\mathcal{E}(-E - C)) = 0$. But, the usual computation shows that this equals $h^1(f^*(-F))$. Since $f^{-1}(F)$ is connected, we conclude that the fibers of $p \circ f : X \rightarrow \mathbb{P}^1$ are all connected and thus $h^1(f^*(-F)) = 0$.

Now we will show that $\mathcal{E}(-F - E)$ is spanned by global sections on the complement of E . Let C be a general element of $|E + rF|$ that contains y . Since global sections of $E + rF$ embed $Y - E$, the curve C is smooth and does not belong to the discriminant variety. Thus $\overline{C} := f^{-1}(C)$ is reduced. It is also connected since it is a big divisor on X . Thus $\mathcal{E}_C(-1)$ is spanned by Theorem (1.3). The assertion that $\mathcal{E}(-F - E)$ is spanned by global sections, will follow if we show that $h^1(\mathcal{E}(-F - E - C)) = 0$. But, the usual computation shows that this equals $h^1(f^*(-F))$. Since $f^{-1}(F)$ is connected, we conclude that the fibers of $p \circ f : X \rightarrow \mathbb{P}^1$ are all connected and thus $h^1(f^*(-F)) = 0$.

Since \mathcal{E} is spanned by Theorem (3.3), for \mathcal{E} to fail to be ample, there must exist an irreducible and reduced curve $D \subset Y$ such that \mathcal{E}_D has a trivial line bundle as a direct summand. Since $\mathcal{E}(-E)$ is spanned, we conclude that $D \cdot E = 0$ so that $D \in |a(E + rF)|$ for some integer $a > 0$. Since $\mathcal{E}(-E - F)$ is spanned off E and since $D \subset Y \setminus E$, we conclude that $D \cdot F = 0$. Since $D \sim a(E + rF)$, we conclude that $a = 0$ and thus that D is the empty curve. Q.E.D.

Remark 3.8 Hirzebruch [Hi83] (see also, [BHH87]), showed how to associate smooth projective surfaces to configurations of lines on \mathbb{P}^2 . Let Λ denote the configuration of six lines through four points in \mathbb{P}^2 , no two of which are collinear. Associated to this configuration there is a smooth projective surfaces $\mathcal{H}(\Lambda, k)$ for each integer $k \geq 2$ with the following properties:

1. there are finite morphisms $f_k : \mathcal{H}(\Lambda, k) \rightarrow Y$, where Y is \mathbb{P}^2 blown up at the four points; and
2. the covering f_k is Galois with Galois group equal to the direct sum of five copies of \mathbb{Z}_k .

Since Y is Del Pezzo with $\omega_Y^2 = 5$, Theorem (3.2) shows that the bundle \mathcal{E} associated to f_k is spanned. This is particularly interesting because of the beautiful result of Hirzebruch that $X := \mathcal{H}(\Lambda, 5)$ is a quotient of the unit ball in \mathbb{C}^2 by a freely acting discrete group.

It would be interesting to know in general what are the properties of the bundle \mathcal{E} associated to the natural finite morphisms from line configuration surfaces to blowups of \mathbb{P}^2 . In particular:

Question 3.9 *For which line configurations Λ and which integers k are the bundles \mathcal{E} spanned?*

If there are any points where more than two lines of the configuration meet, then the surface Y fibers over \mathbb{P}^1 with the inverse image of the general fiber of this fibration disconnected in the line configuration surface, see, e.g., [So84]. This shows that \mathcal{E} will never be ample, except possibly for the trivial configuration where at most two lines meet in a point.

4 Coverings over Del Pezzo Manifolds

We always let $\rho(Y)$ denote the Picard number of Y . The guideline of this section is the following

Conjecture 4.1 *Let $f : X \rightarrow Y$ be a finite surjective morphism of degree $d \geq 2$ between projective manifolds X, Y of dimension n . If Y is Fano with $\rho(Y) = 1$, then \mathcal{E} is ample (and spanned in most cases).*

First we consider the case that Y is a Del Pezzo manifold of dimension $n = \dim Y \geq 3$, i.e. $-\omega_Y = (n - 1)H$ with some (ample) line bundle H . Let $b := H^n$. Then $1 \leq b \leq 8$, moreover all Del Pezzo manifolds are classified (see [Fu90]). In our standard situation (1.1) we obtain

$$h^0(\mathcal{E}(-H)) = h^0(X, \omega_{X/Y} \otimes f^*((n - 2)H)) \quad (1)$$

$$h^0(\mathcal{E}) = h^0(X, \omega_{X/Y} \otimes f^*((n - 1)H)) - 1. \quad (2)$$

As (1.26) shows, \mathcal{E} is not always spanned if $b = 1$. Though we expect spannedness for the cases $b \geq 2$, we can only prove it for $b \geq 5$.

Theorem 4.2 *Let $f : X \rightarrow Y$ be a finite surjective morphism from an n -dimensional normal irreducible Gorenstein projective variety X onto an n -dimensional projective manifold Y . Assume that the irrational locus of X is finite. Let Y be a Del Pezzo manifold with $b \geq 5$. Then \mathcal{E} is spanned.*

Proof. We proceed by induction on n , noticing that H is very ample. We denote a general member of $|H|$ again by H . If $n = 2$, then we conclude by (3.2). So suppose $n \geq 3$. By (1.17)(4) we obtain the exact sequence

$$0 \rightarrow H^0(\mathcal{E}(-H)) \rightarrow H^0(\mathcal{E}) \rightarrow H^0(\mathcal{E}_H) \rightarrow 0.$$

By induction hypothesis, $\mathcal{E}|_H$ is spanned, so it follows that \mathcal{E} is spanned outside a finite set Z . But, H being very ample, through a given point we can still find

a smooth member of $|H|$, hence $Z = \emptyset$. Notice that here we need to work in the Gorenstein category since the general $f^{-1}(H)$ need not be smooth! Q.E.D.

In the cases $1 \leq b \leq 4$, we do not know the spannedness of \mathcal{E} on the corresponding Del Pezzo surfaces, we are only able to prove much weaker results. We start with a general result showing that almost always \mathcal{E} has at least one section.

Theorem 4.3 *Let $f : X \rightarrow Y$ be a finite surjective morphism of degree $d \geq 2$ from an n -dimensional normal irreducible Gorenstein projective variety X onto an n -dimensional projective manifold Y . Assume that the irrational locus of X is finite. Assume that $n \geq 2$ and that Y is Del Pezzo. If $bd \geq 5$, i.e., if either $b \geq 3$; or $b = 2$ and $d \geq 3$; or $b = 1$ and $d \geq 5$; then $h^0(\mathcal{E}) \neq 0$.*

Proof. Write $-\omega_Y = (n-1)H$ with H ample. Note that if $n \geq 3$,

$$h^1(\mathcal{E}(-H)) = h^{n-1}(-f^*(n-2)H) = 0.$$

Since H is either spanned or has at most one base point, we can reduce the proof of the result to the case when $n = 2$.

So without loss of generality we assume that $n = 2$. Let $L := f^*H$. We have

$$h^0(\mathcal{E}) = h^2(\mathcal{E}^* \otimes \omega_Y) = h^2(-L) - 1 = h^0(\omega_X + L) - 1.$$

Thus we have reduced to showing that $h^0(\omega_X + L) \geq 2$. Note that since $h^0(\omega_X + L) = h^0(\mathcal{E}) + 1$, we know that $h^0(\omega_X + L) \geq 1$. From this it follows that $\omega_X + L$ is nef, see [So85]. Again using the main result of [So85], note in this case that $(\omega_X + L) \cdot L = 0$ implies that $-\omega_X = L$, which implies the absurdity that f is unramified. Thus we have that:

1. L is spanned outside of a finite set;
2. $\omega_X + L$ is nef and $(\omega_X + L) \cdot L \geq 2$.

Noting that $L \cdot L = b \deg f$ implies that $L \cdot L \geq 5$, the following Lemma will complete the proof of the theorem.

Lemma 4.4 *Let L be an ample line bundle on an irreducible normal Gorenstein projective surface X . Assume that L is spanned outside of a finite set; $h^0(\omega_X + L) \geq 1$; and $(\omega_X + L) \cdot L \geq 2$. Then either $h^0(\omega_X + L) \geq 2$ or $L \cdot L \leq 4$.*

Proof of Lemma. We assume that $h^0(\omega_X + L) = 1$ and then argue to a contradiction. Since $h^0(L) \geq 2$, we would have $h^0(\omega_X + L) \geq 2$ if $h^0(\omega_X) \geq 1$. Therefore without loss of generality we can assume that

$$h^0(\omega_X) = 0. \tag{3}$$

Since $h^1(\omega_X + L) = 0$, we conclude from

$$0 \rightarrow \omega_X \rightarrow \omega_X \otimes L \rightarrow \omega_C \rightarrow 0$$

that

$$g = q + 1, \quad (4)$$

where $q = h^1(\mathcal{O}_X)$ and $2g - 2 = (\omega_X + L) \cdot L$ defines the arithmetic genus g of a reduced curve C from $|L|$. Since $g \geq 2$ by hypothesis, we conclude further that

$$q \geq 1. \quad (5)$$

Assume first that the irrational locus $\mathcal{I}(X)$ of X is nonempty. Recall the Grauert-Riemenschneider canonical sheaf \mathcal{K}_X is defined as $\pi_*(\omega_{\overline{X}})$ where $\pi : \overline{X} \rightarrow X$ is a desingularization. Following the argument of [So85], we consider the sequence:

$$0 \rightarrow \mathcal{K}_X \otimes L \rightarrow \omega_X \otimes L \rightarrow \mathcal{S} \rightarrow 0,$$

where \mathcal{S} is a skyscraper sheaf whose support exactly equals the set of nonrational singularities $\mathcal{S}(X)$. Since $h^1(\mathcal{K}_X + L) = 0$, $h^0(\omega_X + L) = 1$, and the locus $\mathcal{I}(X)$ is nonempty, we conclude that $h^0(\mathcal{S}) = 1$. Using this, equation (3), and the sequence:

$$0 \rightarrow \mathcal{K}_X \rightarrow \omega_X \rightarrow \mathcal{S} \rightarrow 0,$$

we conclude that:

$$h^0(\omega_{\overline{X}}) = 0; \quad \overline{q} := h^1(\mathcal{O}_{\overline{X}}) = q + 1. \quad (6)$$

Therefore X has only one irrational singularity, an elliptic singularity y_0 . Consider the Albanese mapping $\alpha : \overline{X} \rightarrow Z$. By equation (6) we conclude first that $\overline{q} \geq 1$ and therefore that $\dim Z \geq 1$. Since $h^0(\omega_{\overline{X}}) = 0$, we conclude that Z is a curve, and thus smooth with α having connected fibers. Since $q \geq 1$ by equation (5), we conclude that $\overline{q} \geq 2$. Let B denote the singular set of X , and note that $\alpha(\pi^{-1}(B))$ is finite. This is clear since y_0 is elliptic and all other singularities are rational. In particular, the strict transform \overline{C} of C cannot be a fiber of α . Thus a general fiber F of α is mapped isomorphically by π onto a curve F' contained in the smooth points of X . Since L is spanned off of a finite set, we conclude that global sections of L give rise to at least two linearly independent sections of $L_{F'}$. Hence we deduce easily that $C \cdot F' \geq 2$ and \overline{C} has degree at least 2 over Z . Thus using that $\overline{q} \geq 2$, we have that

$$g \geq 2\overline{q} \geq 2q + 2 = 2g,$$

which is absurd since $g \geq 2$.

Therefore we can assume without loss of generality that the singularities of X are rational. Thus, e.g., [BS95], the Albanese map $\alpha : X \rightarrow Z$ is well defined. Since $h^0(\omega_X) = 0$ by equation (3), we conclude that by the same argument as above, but without the need of a desingularization, that if $q \geq 2$ then

$$g \geq 2q = 2g - 2,$$

which is absurd given that $g = q + 1 \geq 3$. Thus we conclude that $q = 1$ and $g = 2$. In this case we have that

$$(\omega_X + L)^2 L^2 \leq [L \cdot (\omega_X + L)]^2 = 1.$$

Since $L \cdot L \geq 5$ we conclude using the nefness of $\omega_X + L$ that $(\omega_X + L)^2 = 0$. This gives $\omega_X^2 \geq 1$. Since $q = 1$, $h^0(\omega_X) = 0$, and we have at worst rational Gorenstein singularities, we conclude from the Riemann-Roch formula for the Euler characteristic of the structure sheaf of the desingularization, that $\omega_X^2 \leq 0$.

This proves the Lemma and the Theorem. Q.E.D.

Remark 4.5 Note if Y is an elliptic curve, which can be considered as the most appropriate definition of a Del Pezzo manifold of dimension 1, then Theorem (4.2) is still true, unless f is an unramified covering.

We now show how adjunction theory can be used to give general conditions for an existence theorem, which would not follow even from knowing that \mathcal{E} is spanned. Indeed, as shown in (4.10), some of the possible exceptions to this result exist and in fact have \mathcal{E} spanned. We do not need the precise classification of Del Pezzo manifolds with $1 \leq b \leq 4$ after Fujita [Fu90], but just the following

Proposition 4.6 *Let Y be a Del Pezzo manifold of dimension at least 3. Then $\rho(Y) = 1$, unless X is $(\mathbb{P}^1)^3$, $\mathbb{P}^2 \times \mathbb{P}^2$, $\mathbb{P}(T_{\mathbb{P}^2})$ or \mathbb{P}^3 blown up in one point. Moreover H is very ample for $b \geq 3$, and spanned unless $b = 1$, in which case we have one simple base point. In all cases $|H|$ contains a smooth element.*

Theorem 4.7 *Let $f : X \rightarrow Y$ be a finite surjective morphism of degree $d \geq 2$ from an n -dimensional projective manifold X onto an n -dimensional projective manifold Y . Assume that $n \geq 3$ and that Y is Del Pezzo, i.e., $-\omega_Y = (n-1)H$ with H ample. Let Y be of degree $b := H^n \geq 2$. Then $H^0(\mathcal{E}(-H)) \neq 0$, except for the following possible cases: in the cases 3) and 4), (X_0, L_0) denotes the first reduction of (X, L) [BS95].*

1. (X, L) is a quadric fibration over a curve with $Y = (\mathbb{P}^1)^3$; or
2. (X, L) is a scroll over a surface with either $n = 3$; or
3. (X_0, L_0) is a 3-dimensional quadric, $L_0 = \mathcal{O}(2)$, $Y = \mathbb{P}^3(x_0)$, $d = 2$, and f is induced by a covering $f_0 : X_0 \rightarrow Y_0 = \mathbb{P}^3$; or
4. $n = 3$ and X_0 is a \mathbb{P}^2 -bundle over a curve with $L_F = \mathcal{O}_{\mathbb{P}^2}(2)$ on a fiber F .

Examples of dimension 3, 4, for the cases of the Theorem are given in (4.10).

Proof. Let $L = f^*(H)$. Suppose $H^0(\mathcal{E}(-H)) = 0$. Then by (1)

$$H^0(\omega_X \otimes f^*((n-2)H)) = 0.$$

Then, since $b \geq 2$ and H is spanned, [So89] applies. Using the equation (2) that $h^0(\omega_X + (n-1)L) = h^0(\mathcal{E}) + 1 \geq 1$, we obtain that either (X, L) is

- (a) a quadric fibration over a curve; or

(b) a scroll over a surface; or,

(X_0, L_0) , the first reduction of (X, L) [BS95], exists and is either:

(c) $(X_0, L_0) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(2))$ with $n = 3, 4$; or

(d) (X_0, L_0) is a 3-dimensional quadric and $L_0 = \mathcal{O}(2)$; or

(e) $n = 3$ and X_0 is a \mathbb{P}^2 -bundle over a curve with $L = \mathcal{O}(2)$ on the fibers.

We will treat these cases separately.

(a) If we have a quadric fibration $p : X \rightarrow C$, let F be a general fiber and $F' = f(F)$. Since $L^{n-1} \cdot F = 2$, the map f_F has degree 2 or 1. H being ample and spanned (recall $b \geq 2$), it follows that in the first case F' must be projective space, and the adjunction formula implies $N_{F'|Y} = \mathcal{O}(-1)$, which is absurd since F' moves in Y . In the second case, we obtain $H^{n-1} \cdot F' = 2$ and we conclude similarly that F' is a quadric. By adjunction, $N_{F'|Y} = \mathcal{O}$, hence $\rho(Y) \geq 2$. By the classification (4.6) we obtain $Y = (\mathbb{P}^1)^3$. This is in one of the exceptions.

(b) Now suppose that $p : X \rightarrow S$ is a scroll over the surface S . Then, using the same notations as before, we conclude $F' = \mathbb{P}^{n-2}$ and $H|F' = \mathcal{O}(1)$. By adjunction, $\det N_{F'} = \mathcal{O}$. Notice that $N_{F'}$ is generically generated for *general* F' , hence $N_{F'}$ must be trivial. If $b \geq 3$ and $n \geq 5$, then H is very ample, so that by Ein [Ei85], Y has a \mathbb{P}^{n-2} -bundle structure, contradicting (4.6) unless $Y = \mathbb{P}^2 \times \mathbb{P}^2$ giving rise to one of the exceptions (4.10). Therefore it remains to treat the case $b = 2$ or $n \leq 4$.

(b.1) In case $n = 3$, then F' is a line and we are in the exceptions (4.10), the cases $Y = \mathbb{P}(T_{\mathbb{P}^2}), \mathbb{P}_3(x), \mathbb{P}_1^3$ being obvious.

(b.2) Concerning $b = 2$, the linear system $|H|$ realizes Y as a $2 : 1$ -covering $h : Y \rightarrow \mathbb{P}_n$, moreover $L = f^*(H)$. Since $H|F' = \mathcal{O}(1)$, it follows that $f|F$ is $1 : 1$ for all F . This is clearly absurd, since $f^{-1}(F')$ would always consist of 1 or 2 disjoint fibers of p , which implies $\rho(Y) \geq 2$, a contradiction to the classification.

(b.3) It remains to treat the case $b \geq 3$ and $n = 4$.

(α) Suppose $b = 3$. Then $Y \subset \mathbb{P}_5$ has degree and we obtain (for general $F' = \mathbb{P}_2$) an exact sequence

$$0 \rightarrow N_{\mathbb{P}_2|Y} \rightarrow N_{\mathbb{P}_2|\mathbb{P}_5} \rightarrow N_{Y|\mathbb{P}_5}|_{\mathbb{P}_2} \rightarrow 0,$$

which reads

$$0 \rightarrow \mathcal{O} \oplus \mathcal{O} \rightarrow \mathcal{O}(1)^{\oplus 3} \rightarrow \mathcal{O}(3) \rightarrow 0.$$

This sequence is of course absurd.

(β) Suppose $b = 4$. Then Y is the intersection of 2 quadrics in \mathbb{P}_6 . Take one of the quadrics, say Q , to obtain a sequence

$$0 \rightarrow N_{\mathbb{P}_2|Y} \rightarrow N_{\mathbb{P}_2|Q} \rightarrow N_{Y|Q}|_{\mathbb{P}_2} \rightarrow 0.$$

This sequence immediately implies

$$N_{\mathbb{P}_2|Q} = \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(2),$$

which clearly contradicts the 1-ampleness of the tangent bundle T_Q .

(γ) Finally we treat $b = 5$. First notice that every line $l \subset Y$ has a normal bundle either of the form $\mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(1)$ or $\mathcal{O}(1) \oplus \mathcal{O}(1) \oplus \mathcal{O}(-1)$. This follows by choosing a smooth hyperplane section through l (which is easily seen to exist, cp. [Is77,p.505]). We are going to show that *every* plane $F' \subset Y$ has normal bundle $\mathcal{O} \oplus \mathcal{O}$. Once we know this, we can argue as in [Ei85,1.7] to obtain a \mathbb{P}_2 -bundle structure on Y . Since the planes F' cover Y , the normal bundle N is generically generated by 2 sections s_1, s_2 . Since by adjunction $\det N = \mathcal{O}$, it follows that these sections generate N everywhere so that N is trivial.

(c) Suppose $(X_0, L_0) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(2))$ with $n = 3$ or 4 . We claim that always $X = X_0$. In fact, if $X \neq X_0$, then X contains a divisor $E = \mathbb{P}^{n-1}$ with $N_E = \mathcal{O}(-1)$. Consider $E' = f(E)$. Since $L|E = \mathcal{O}(1)$, we have $H^{n-1} \cdot E' = 1$, so that since H is therefore spanned, we have that $E' = \mathbb{P}^{n-1}$. Adjunction gives $N_{E'} = \mathcal{O}(-1)$, hence by classification $Y = \mathbb{P}^3(x_0)$. Therefore $X_0 = \mathbb{P}^3$. Now, introducing the ramification divisor R of f , we have

$$\omega_X + f^*(H) = f^*(\omega_Y) + R + f^*(H) = f^*(-H) + R,$$

hence

$$H^0(X, \mathcal{O}_X(R) \otimes f^*(-H)) = 0$$

by our assumption. If $\sigma : X \rightarrow X_0$ denotes the contraction of the exceptional divisors E , we conclude that

$$R' = \sigma_*(R) \in |\mathcal{O}_{X_0}(1)|.$$

Hence the induced covering $h : X_0 \rightarrow Y_0$ is ramified exactly over a plane, which is absurd since $X_0 = Y_0 = \mathbb{P}^3$. So we shall assume $X = X_0$. But then we have the contradiction that $Y = \mathbb{P}^n$.

(d) Suppose X_0 is a three dimensional quadric with $L = \mathcal{O}_{X_0}(2)$. If $X = X_0$, then Y is \mathbb{P}^3 or a quadric. Since Y is Del Pezzo, we have $Y = \mathbb{P}^3$ and $H = \mathcal{O}_{\mathbb{P}^3}(2)$. Since $K_X + L$ has no sections, we conclude that $L = \mathcal{O}_X(2)$, and therefore that f has degree 2. From this we conclude that $\mathcal{E} = \mathcal{O}_{\mathbb{P}^3}(1)$. This gives an example. If $X \neq X_0$, then we argue as in (c) to conclude that $Y = \mathbb{P}^3(x_0)$. Hence $Y_0 = \mathbb{P}^3$, see (4.10)(4). Q.E.D.

Remark 4.8 In the preceding and following results we need $b \geq 2$ to get spannedness so that the main result of [So89] holds. Actually, we only need that $2L$ is spanned and $|L|$ has a smooth element. This would be satisfied when $b = 1$ under the extra condition that the point where H is not spanned is not an element of the discriminant divisor of the covering. Using this the statement of the result is the same, except there is one case that cannot be ruled out: the first reduction (X_0, L_0) of (X, L) exists and is equal $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(2))$ with $n = 3, 4$.

Corollary 4.9 *If $b \geq 2$, then $h^0(\mathcal{E}) \geq h^0(\mathcal{O}_Y(H))$ unless X is of the one of the exceptions in Theorem (4.7). In particular these inequalities hold if $n \geq 5$.*

We now give examples of the exceptional cases.

The Exceptional Cases 4.10 (1) $Y = (\mathbb{P}^1)^3$ and we fix a projection $q : (\mathbb{P}^1)^3 \rightarrow \mathbb{P}^1$. Let $h : C \rightarrow \mathbb{P}^1$ be a suitable ramified cover and put $X = Y \times_{\mathbb{P}^1} C$.

(2) Let Y be a Del Pezzo 3-fold and T a component of the Hilbert schemes of lines. Suppose that through every point of X there are only finitely many lines. Let S be a smooth surface and $S \rightarrow T$ be a surjective map. Let X be the \mathbb{P}^1 -bundle over S given by the graph of the family of line parameterized by T .

(3) Let $Y = \mathbb{P}^2 \times \mathbb{P}^2$. Let S be a surface and $S \rightarrow \mathbb{P}^2$ be a finite map. Put $X = S \times \mathbb{P}^2 \rightarrow Y$. Analogously we choose $S \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ finite and let $X = S \times \mathbb{P}^1$ resp. $h : S \rightarrow \mathbb{P}^2$ finite and $X = \mathbb{P}(h^*(T_{\mathbb{P}^2}))$.

(4) Suppose X_0 is a three dimensional quadric. We have a degree 2 map $f : X_0 \rightarrow \mathbb{P}^3$. In this case $\mathcal{E} = \mathcal{O}_{\mathbb{P}^3}(1)$. Since \mathbb{P}^3 is Del Pezzo with $H = \mathcal{O}_{\mathbb{P}^3}(2)$, we have $\mathcal{E}(-H)$ has no sections. Note $L_0 = \mathcal{O}_{X_0}(2)$. If $\pi : Y \rightarrow \mathbb{P}^3$ is the blowup of \mathbb{P}^3 at a point y . Then $\overline{H} := \pi^*\mathcal{O}_{\mathbb{P}^3}(2) - E$ is very ample where E is the exceptional fiber of π . Thus $-K_Y = \pi^*\mathcal{O}_{\mathbb{P}^3}(4) - 2E = 2\overline{H}$, and we obtain a Del Pezzo manifold Y . Letting X denote the blowup of X_0 at the inverse image under f of y , we have a degree 2 finite morphism $\overline{f} : X \rightarrow Y$. We have $h^0(K_X + L) = h^0(K_{X_0} + L_0) = 0$.

(5) Let $p : X_0 \rightarrow C$ be a \mathbb{P}^2 -bundle over a smooth curve C . Let $h : X_0 \rightarrow Y_0 = \mathbb{P}^3$ be a finite cover such that

$$[h^*\mathcal{O}_{\mathbb{P}^3}(1)]_F = \mathcal{O}_{\mathbb{P}^2}(1),$$

where F denotes a fiber of p . Choose y_0 not in the image of the ramification divisor of h . Then let $\tau : Y \rightarrow Y_0$ be the blow-up of Y_0 at y_0 and $\sigma : X \rightarrow X_0$ the blow-up of X_0 at the finite set $h^{-1}(y_0)$. Then h lifts to a finite cover $f : X \rightarrow Y$ and it is easy to check that $H^0(X, \omega_X \otimes f^*(H)) = 0$, where $-\omega_Y = 2H$.

In order to obtain a specific example, consider the family \mathcal{C} of hyperplanes in \mathbb{P}^3 . The canonical map

$$q : \mathcal{C} \rightarrow \mathbb{P}^3 = Y_0$$

is a \mathbb{P}^2 - bundle and so does the other projection

$$p : \mathcal{C} \rightarrow \mathbb{P}^{3*}.$$

Actually $\mathcal{C} = \mathbb{P}(T_{\mathbb{P}^3})$. Now choose $C \subset \mathbb{P}^{3*}$ to be a general smooth curve. Then $X_0 = p^{-1}(C)$ is smooth and moreover $h := q|_{X_0}$ is finite. To see this last property, consider $T(x) \subset \mathbb{P}^{3*}$, the set of hyperplanes through a given point $x \in \mathbb{P}^3$. So it is sufficient to choose C such that $C \not\subset T(x)$ for all x which comes down to choose C non-degenerate, since all $T(x)$ are linear in $(\mathbb{P}^3)^*$.

References

- [BHH87] Barthel,G.; Hirzebruch,F.; Höfer,Th.: Geradenkonfigurationen und Algebraische Flächen. Aspects of Mathematics, D4. Friedr. Vieweg & Sohn 1987
- [BS95] Beltrametti,M.; Sommese,A.J.: The adjunction theory of complex projective varieties. de Gruyter 1995
- [CP97] Campana,F.;Peternell,Th.: Towards a Mori theory on compact Kähler threefolds, I. Math. Nachr. 187, 29–59 (1997)
- [CS94] Cho,K.;Sato,E.: Smooth projective varieties dominated by smooth hyperquadrics in any characteristic. Math. Z. 217, 553–565 (1994)
- [De95] Debarre,O.: Théoremes de connexité et variétés abéliennes. Amer.J.Math. 117, 787–805 (1995)
- [De96] Debarre,O.: Sur un théoreme de connexité de Mumford pour les espace homogenes. Manusc. Math. 89, 407-425 (1996)
- [DPS94] Demailly,J.P.;Peternell,Th.;Schneider,M.: Compact complex manifolds with numerically effective tangent bundles. J. Alg. Geom. 3, 295–345 (1994)
- [DPS99] Demailly,J.P.;Peternell,Th.;Schneider,M.: Pseudo-effective line bundles on projective varieties and Kähler manifolds. Preprint in prep. (1999)
- [Ei85] Ein,L.: Varieties with small dual varieties,II. Duke Math. J. 52, 895–907 (1985)
- [Fu90] Fujita,T.: Classification theories of polarized varieties. London Math. Soc. Lect. Notes Ser. 155 (1990)
- [Ha71] Hartshorne,R.: Ample vector bundles on curves. Nagoya Math. J. 43, 73–89 (1971)
- [Hi83] Hirzebruch,F.: Arrangements of lines and algebraic surfaces. Arithmetic and geometry, Vol. II, 113–140, Progr. Math., 36, Birkhäuser, Boston, Mass., 1983
- [Is77] Iskovskih,V.A.: Fano 3-folds I. Math. USSR Izv. 11, 485-527 (1977)
- [Ki96a] Kim,M.: Barth-Lefschetz type theorem for branched coverings of Grassmannians, J. Reine Angew. Math. 470, 109–122 (1996)
- [Ki96b] Kim,M.: On branched coverings of quadrics. Arch. Math. 67, 76–79 (1996)
- [KM98] Kim,M.;Manivel,L.: On branched coverings of some homogeneous spaces. Topology 38, 1141–1160 (1999)

- [Ma97] Manivel,L.: Vanishing theorems for ample vector bundles. *Invent. Math.* 127, 401–416 (1997)
- [La80] Lazarsfeld,R.: A Barth type theorem for branched coverings of projective spaces. *Math. Ann.* 249, 153–162 (1980)
- [OSS80] Okonek,C.;Schneider,M.;Spindler,H.: *Vector bundles on complex projective spaces.* Birkhäuser 1980
- [PS89] Paranjapé,K.H.;Srinivas,V.: Selfmaps of homogeneous spaces. *Inv. Math.* 98, 425–444 (1989)
- [So78] Sommese,A.J.: Submanifolds of Abelian varieties. *Math. Ann.* 233, 229–256 (1978)
- [So84] Sommese,A.J.: On the density of ratios of Chern numbers of algebraic surfaces. *Math. Ann.* 268, 207–221 (1984)
- [So85] Sommese,A.J.: Ample divisors on normal Gorenstein surfaces. *Abh. Math. Sem. Univ. Hamburg* 55, 151–170 (1985)
- [So86] Sommese,A.J.: On the adjunction theoretic structure of projective varieties. In: *Proceedings of the Complex Analysis and Algebraic Geometry Conference, Göttingen, 1985*, ed. by H. Grauert, *Lecture Notes in Math.* 1194, 175–213. Springer 1986
- [So89] Sommese,A.J.: On the nonemptiness of the adjoint linear system of a hyperplane section of a threefold. *J. reine u. angew. Math.* 402, 211–220 (1989); Erratum, 411, 122–123 (1990)
- [Vi82] Viehweg,E.: Die Additivität der Kodairadimension für projektive Faserräume über Varietäten des allgemeinen Typs. *J.reine u. angew. Math.* 330, 132–142 (1982)
- [Vi95] Viehweg,E.: *Quasi-projective moduli for polarized manifolds.* Springer 1995

Thomas Peternell
 Mathematisches Institut
 Universität Bayreuth
 D-95440 Bayreuth, Germany
 fax: Germany + 921-552999
 thomas.peternell@uni-bayreuth.de

Andrew J. Sommese
 Department of Mathematics
 University of Notre Dame
 Notre Dame, Indiana 46556, U.S.A,
 fax: U.S.A. + 219-631-6579
 sommese@nd.edu
 URL: www.nd.edu/~sommese

Appendix: Nefness Modulo the Branch Locus of the Bundle Associated to a Branched Covering

Robert Lazarsfeld

Dedicated to Robin Hartshorne on his 60th birthday

We give the solution to a problem posed in preceding paper of Peternell and Sommese, whose notation and bibliography we follow. The paper [La80] which we are revisiting here was very much influenced by the work of Robin Hartshorne. It is a pleasure to use this occasion to express our debt to Hartshorne and to convey to him our best wishes. We also wish to acknowledge the support of the NSF (Grant DMS 97-13149) and the J. S. Guggenheim Foundation.

We start by restating (in slightly greater generality) the result in question:

Proposition A *Let $f : X \rightarrow Y$ be a finite and flat covering of complex projective varieties, with Y smooth and irreducible, and X reduced (but possibly reducible). Let $C \subset Y$ be any irreducible curve not contained in the branch locus of f . Then $\mathcal{E}_f|_C$ is nef, i.e. $(f_*\mathcal{O}_X)|_C$ is the dual of a nef vector bundle on C .*

The hypothesis on C is that f is generically étale over C . Note that then $D = f^{-1}(C) \subset X$, being generically reduced and flat over the reduced and irreducible variety C , is itself (everywhere) reduced. Observe also that since f is flat and finite, the direct image functor f_* commutes with base-change. Moreover, by normalizing we can assume without loss of generality that C is smooth. Therefore:

Reduction B *It suffices to prove Proposition A under the additional assumption that X and Y are curves. \square*

The essential remark is the following

Lemma C *Let $f : X \rightarrow Y$ be a branched covering of curves satisfying the hypotheses of Proposition A. Then there exists a finite covering $p : Y' \rightarrow Y$ of smooth irreducible curves, plus a Cartesian square:*

$$\begin{array}{ccc} X' & \xrightarrow{p'} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{p} & Y \end{array} ,$$

where $X' = X \times_Y Y'$ is a reduced curve each of whose irreducible components maps isomorphically via f' to Y' .

Grant this for the moment. Since the nefness of \mathcal{E}_f can be tested after pulling back by p , and since f_* commutes with base-change, it is enough to prove that $f'_*\mathcal{O}_{X'}$ is the dual of a nef bundle on Y' . But since X' is a union of copies of Y' , $f'_*\mathcal{O}_{X'}$ sits naturally as a

subsheaf with torsion cokernel inside a direct sum of copies of $\mathcal{O}_{Y'}$. And in this case the nefness of $(f'_* \mathcal{O}_{X'})^*$ is clear.

It remains only to give the proof of Lemma C. Denote by X_1, \dots, X_t the irreducible components of X , and set

$$d^* = d^*(f) = \sum_i \left(\deg(X_i \rightarrow Y) - 1 \right).$$

We argue by decreasing induction on d^* , the goal being to arrange that $d^* = 0$. Assuming that $d^* > 0$, suppose (after reindexing) that $\deg(X_1 \rightarrow Y) \geq 2$. Let $Y^\sharp \rightarrow X_1$ be the normalization of X_1 , and consider the fibre product $X^\sharp = X \times_Y Y^\sharp \rightarrow Y^\sharp$:

$$\begin{array}{ccc} X^\sharp & \xrightarrow{p^\sharp} & X \\ f^\sharp \downarrow & & \downarrow f \\ Y^\sharp & \xrightarrow{p} & Y \end{array} .$$

Then Y^\sharp embeds in X^\sharp as an irreducible component of $(p^\sharp)^{-1}(X_1)$, and consequently $(p^\sharp)^{-1}(X_1)$ splits into at least two components. Therefore

$$d^*(f^\sharp) < d^*(f),$$

and we are done.

Robert Lazarsfeld
 Department of Mathematics
 University of Michigan
 Ann Arbor, MI 48109
 rlaz@math.lsa.umich.edu