

v_* -TORSION SPACES AND THICK CLASSES

W. CHACHÓLSKI, W. G. DWYER, AND M. INTERMONT

1. INTRODUCTION

Suppose that p is some chosen prime number. A topological space is said to be v_* -torsion if its homotopy groups are concentrated at p and its v_n -periodic homotopy groups vanish for $n \geq 1$ (1.1). A similar definition can be made for spectra, and in this stable case it is easy to see that a spectrum is v_* -torsion if and only if it can be built (in the sense of [9]) from *any* finite spectrum with nontrivial mod p homology. The unstable analog of this statement is not true, but in this paper we provide a close substitute. For any space W , let \mathcal{B}_W denote the class of spaces which can be built from W , and $\mathcal{T}(\mathcal{B}_W)$ the smallest class of spaces which contains \mathcal{B}_W and is thick with respect to fibrations (1.6). We prove that a space X is v_* -torsion if and only if it belongs to $\mathcal{T}(\mathcal{B}_W)$ for *any* finite complex W with nontrivial reduced mod p homology. We go on to show that, although the category of v_* -torsion spaces is not closed under arbitrary homotopy colimits, it is closed in many cases with respect to realizations of simplicial objects. Finally, we consider the Stover-Blanc simplicial resolution $R_W(X)$ of a space X , which comes up in a natural attempt to construct the W -cellular approximation $\text{Cell}_W(X)$. We show that for appropriate W the fibres of the map from the geometric realization of $R_W(X)$ to $\text{Cell}_W(X)$ are v_* -torsion. Since v_* -torsion spaces are in some respects negligible objects, this can be interpreted as saying that the realization of the Stover-Blanc construction is closer to $\text{Cell}_W(X)$ than one might suspect.

1.1. v_* -torsion spaces. Let p be a prime number, which is fixed for the rest of the paper. For each integer $n \geq 1$, let V_n be a pointed finite p -torsion complex *of type n* in the sense of chromatic homotopy theory [13], so that the reduced Morava K -theory module $\tilde{K}(i)_*V_n$ is trivial for $i < n$ and nontrivial for $i = n$. By a theorem of Hopkins-Smith [11], after replacing V_n by a large enough suspension we can find an integer $d > 0$ and a map $\omega : \Sigma^d V_n \rightarrow V_n$ which is a v_n -self-map, in the sense that it induces an isomorphism on $\tilde{K}(n)_*$ and induces the

Date: July 28, 2004.

zero map on $\tilde{K}(i)_*$ for $i \neq n$. For a pointed space X , the v_n -periodic homotopy groups $v_n^{-1}\pi_*(X; V_n)$ are obtained by inverting the action of ω on $\pi_*(X; V_n)$ [2, 11.3].

1.2. Definition. Let $n \geq 1$ be an integer. An unpointed space X is v_n -torsion if for each choice of basepoint in X , the v_n -periodic homotopy groups $v_n^{-1}\pi_*(X; V_n)$ vanish.

1.3. Remark. By [11], the answer to the question of whether $v_n^{-1}\pi_*(X; V_n)$ is trivial does not depend upon the choice of the complex V_n or on the choice of the map ω .

Say that a group G is *concentrated at p* if it has a finite subnormal series

$$\{1\} = G_0 \triangleleft G_1 \triangleleft \cdots \triangleleft G_m = G$$

such that each quotient G_i/G_{i-1} is a p -primary torsion abelian group. We will say that an unpointed space X is *concentrated at p* if for each basepoint in X and each $k \geq 1$, the homotopy group $\pi_k(X)$ is concentrated at p .

1.4. Definition. An unpointed space X is said to be v_* -torsion if it is concentrated at p , and it is v_n -torsion for each $n \geq 1$.

1.5. Remark. The condition of being concentrated at p can be interpreted as being “ v_0 -torsion”.

1.6. Thick classes. Let \mathcal{C} be some class of unpointed spaces; all the classes we consider will be closed under retracts, and also *closed under equivalences*, in the sense that whenever $X \rightarrow Y$ is a weak equivalence, then $X \in \mathcal{C}$ if and only if $Y \in \mathcal{C}$. If $q : E \rightarrow B$ is a map of spaces, then F is said to be *a fibre of q* if for some choice of basepoint $b_0 \in B$, F is equivalent to the homotopy fibre of q over b_0 . Consider the following three conditions on $q : E \rightarrow B$, where F runs through the fibres of q :

- (1) $B \in \mathcal{C}$ and $E \in \mathcal{C}$ implies all $F \in \mathcal{C}$;
- (2) $B \in \mathcal{C}$ and all $F \in \mathcal{C}$ implies $E \in \mathcal{C}$;
- (3) $E \in \mathcal{C}$, all F connected, and all $F \in \mathcal{C}$ implies $B \in \mathcal{C}$.

1.7. Definition. Let \mathcal{C} be a class of unpointed spaces, closed under retracts and equivalences. Then \mathcal{C} is said to be *thick* (with respect to fibrations) if for any map $q : E \rightarrow B$ all three of the above conditions hold.

1.8. Remark. If the first condition holds, then \mathcal{C} is said to be *closed under fibres*; if the second condition holds, then \mathcal{C} is said to be *closed under extension by fibrations*.

1.9. *Examples.* The class of spaces concentrated at p is thick, as is the class of v_n -torsion spaces for fixed n , or the class of v_* -torsion spaces.

If \mathcal{C} is a class of spaces, we write $\mathcal{T}(\mathcal{C})$ for the smallest thick class of unpointed spaces containing \mathcal{C} . If W is a pointed CW-complex, let \mathcal{B}_W be the class of spaces *built from* W in the sense of [9]; \mathcal{B}_W is the smallest class of spaces containing W and closed under equivalences and pointed homotopy colimits. Our first main theorem is the following one.

1.10. **Theorem.** *The class of v_* -torsion spaces is the intersection of the classes $\mathcal{T}(\mathcal{B}_W)$, where W ranges over all finite complexes such that $\tilde{H}_*(W; \mathbb{Z}/p)$ is nontrivial.*

1.11. *Remark.* The class \mathcal{B}_W is naturally a class of pointed spaces, but in forming $\mathcal{T}(\mathcal{B}_W)$ we forget the basepoints and work in the unpointed category. The class \mathcal{B}_W does not depend on the choice of basepoint in W .

1.12. *Remark.* The proof of 1.10 shows that $\mathcal{T}(\mathcal{B}_W)$ can also be described as the smallest class of spaces containing \mathcal{B}_W and closed under fibres and extension by fibrations.

1.13. **Closure and Resolutions.** The class of v_* -torsion spaces is not closed under homotopy colimits; for instance, it clearly contains all discrete spaces, and any space is weakly equivalent to a homotopy colimit of discrete spaces. However, it is closed under one special kind of homotopy colimit.

1.14. **Theorem.** *Suppose that X is a simplicial space with the property that each space X_i , $i \geq 0$ is connected, is v_* -torsion, and has an abelian fundamental group. Then the geometric realization $|X|$ of X is also v_* -torsion.*

Note that by the geometric realization $|X|$ of a simplicial space X we mean its homotopy colimit; this is equivalent to the ordinary geometric realization if X satisfies suitable cofibrancy conditions.

Now for the resolution construction. Suppose that W is a pointed CW-complex. For any pointed space X , let $\text{Cell}_W(X)$ denote the *W-cellular approximation to* X [9, 2.A]; this is the closest approximation to X by a space which is built from W . Stover [14] and Blanc [1] have constructed a natural simplicial resolution $R_W(X)$ of X together with a map $|R_W(X)| \rightarrow \text{Cell}_W(X)$; this map is an equivalence if W is a sphere [14] and the question is whether or not it is an equivalence in general. Similar questions are considered in [6]. What we prove is the following.

1.15. Theorem. *Suppose that X is a pointed connected space, and that W is a pointed finite CW-complex which is 1-connected, concentrated at p , and homotopy equivalent to a suspension. Then the homotopy fibres of the map $|\mathbf{R}_W(X)| \rightarrow \mathbf{Cell}_W(X)$ are v_* -torsion.*

This paper is heavily dependent on ideas of Bousfield [2] and Dror-Farjoun [9].

1.16. Organization of the paper. Section 2 recalls some properties of nullification functors, which are a key tool in what we do; §3 uses these functors to give a proof of 1.10. Section 4 gives a recognition principle for v_* -torsion spaces. Section 5 has a brief discussion of the simplest kind of v_* -torsion spaces, the *polygems*, and §6 points out that certain simplicial constructions give polygems. The final section uses the results of the preceding three to prove 1.14 and 1.15.

1.17. Notation and terminology. If W is a space, $c(W)$ denotes the smallest integer k such that $\tilde{H}_*(W; \mathbb{Z}/p)$ is nonzero. If there is no such k , $c(W) = \infty$. Similarly, $\kappa(W)$ denotes the smallest integer n such that $\tilde{K}(n)_*(W)$ is nonzero. We say that a space is *connected* if it is pathwise connected, and that a map of spaces is an *equivalence* if it is a weak homotopy equivalence. The notation $\mathbf{Map}(X, Y)$ denotes the space of maps from X to Y , taken in the derived sense, i.e., before forming the mapping space, X is to be replaced by an equivalent cofibrant object and Y by an equivalent fibrant one. In the category of topological spaces, this involves replacing X by a CW-approximation; in the category of simplicial sets, replacing Y by a Kan complex. If X and Y are pointed, similar considerations apply to the space $\mathbf{Map}_*(X, Y)$ of pointed maps, or to its null component $\mathbf{Map}_*^0(X, Y)$.

2. NULLIFICATION

One of the main tools we use in this paper is the *nullification functor* P_W associated to a CW-complex W [2] [9]. Recall that a space X is said to be *W -null* if the map $W \rightarrow *$ induces an equivalence $X \rightarrow \mathbf{Map}(W, X)$. The nullification functor P_W assigns to any X a map $\epsilon_X : X \rightarrow P_W(X)$ from X to a W -null space $P_W(X)$; this is a *universal* map, in the sense that if Y is another W -null space, $\mathbf{Map}(\epsilon_X, Y)$ is an equivalence. The functor P_W is idempotent and $P_W(\epsilon_X)$ is an equivalence.

The space $P_W(X)$ is called the *nullification of X (with respect to W)*. A map f is said to be a *P_W -equivalence* if $P_W(f)$ is an equivalence. It is easy to see that X is W -null if and only if $P_W(X) \sim X$. If, on the

other hand, $P_W(X) \sim *$, then X is said to be *killed by W* ; we denote by \mathcal{K}_W the class of all such spaces.

Our interest in P_W and its “kernel” \mathcal{K}_W is explained by the following theorem

2.1. Theorem. [7] *For any pointed CW-complex W , the class \mathcal{K}_W is the smallest class of spaces containing \mathcal{B}_W and closed under equivalences and extension by fibrations. In particular $\mathcal{T}(\mathcal{K}_W) = \mathcal{T}(\mathcal{B}_W)$*

In this section we recall some properties of the functor P_W ; we will use these properties in §3 to characterize \mathcal{K}_W for suitable W and to prove 1.10.

2.2. Remark. If S^{k+1} is the $k + 1$ -sphere, then $P_{S^{k+1}}(X)$ is the k 'th Postnikov stage of X . At what we hope is only a minor risk of confusion we will denote the nullification functor $P_{S^{k+1}}$ by the usual Postnikov notation P_k . For each choice of basepoint in X , the natural map $\pi_i X \rightarrow \pi_i P_k X$ is an isomorphism for $i \leq k$, while $\pi_i P_k X$ is trivial for $i > k$.

The following proposition is easy to check directly from the above definitions. The space ΣW is the reduced suspension of W with respect to some chosen basepoint.

2.3. Proposition. *For any W , $\mathcal{K}_{\Sigma W} \subset \mathcal{K}_W$.*

2.4. Remark. This proposition implies that $\mathcal{T}(\mathcal{K}_{\Sigma W}) \subset \mathcal{T}(\mathcal{K}_W)$, and so in order to study the intersection which appears in 1.10, it is enough to consider \mathcal{K}_W for W a suspension. The proposition also implies that $P_W P_{\Sigma W}(X) \sim P_W(X)$, so that at least up to canonical homotopy there is a natural map $P_{\Sigma W}(X) \rightarrow P_W(X)$. Similarly, $P_{\Sigma W} P_W(X) \sim P_W(X)$.

2.5. Proposition. [2, 4.4] *If W is connected, the functor P_W respects coproducts of spaces. If W is k -connected, then for any choice of basepoint in X the map $X \rightarrow P_W(X)$ induces isomorphisms $\pi_i X \cong \pi_i P_W(X)$ for $i \leq k$.*

2.6. Proposition. [2, 4.3, 4.8] *If $f : E \rightarrow B$ is a map all of whose fibres are killed by W , then $P_W(f)$ is an equivalence. All of the fibres of the map $X \rightarrow P_W(X)$ are killed by W .*

2.7. Proposition. (Partial nullification) [2, 4.1] *Suppose that B is connected and that $F \rightarrow E \rightarrow B$ is a fibre sequence. Then there is a natural map*

$$\begin{array}{ccccc} F & \longrightarrow & E & \longrightarrow & B \\ \downarrow & & \downarrow & & \downarrow \\ P_W(F) & \longrightarrow & E' & \longrightarrow & P_{\Sigma W}(B) \end{array}$$

in the homotopy category of homotopy fibre sequences. All three vertical arrows are P_W -equivalences, and E' is ΣW -null.

2.8. Proposition. [2, 7.4] *If G is an abelian group which is concentrated at p and $n \geq c(W)$, then $P_W(K(G, n)) \sim *$.*

2.9. The standard assumptions. We will say that W satisfies the standard assumptions if W is a finite CW-complex which is equivalent to a suspension, is not contractible, and is concentrated at p . Under these assumptions, W is simply connected and $c(W)$ is obtained by adding one to the connectivity of W .

The next proposition is clear from the way in which $P_W(X)$ is constructed by attaching copies of W and its suspensions to X .

2.10. Proposition. [2, 2.8] *If W satisfies the standard assumptions, then X is concentrated at p if and only if $P_W(X)$ is concentrated at p .*

2.11. Definition. A space X is said to be of finite Postnikov type if there is some integer k such that $X \sim P_k X$, i.e., if regardless of the choice of basepoint, $\pi_i X = 0$ for $i > k$.

2.12. Proposition. (cf. [2, 10.5]) *If W satisfies the standard assumptions and X is a space which is concentrated at p and of finite Postnikov type, then $P_W(X) \sim P_\nu(X)$ for $\nu = c(W) - 1$.*

Proof. We can assume that X is connected (2.5). It is clear that $P_\nu(X)$ is W -null, so it is only necessary to show that $X \rightarrow P_\nu(X)$ is a P_W -equivalence. Since X is of finite Postnikov type, it is enough to show that for $j > \nu$, the map $P_j(X) \rightarrow P_{j-1}(X)$ is a P_W -equivalence. But the fibre of this map is $K(\pi_j X, j)$, which is W -null by 2.8. The desired result follows from 2.6. \square

2.13. Proposition. [3, 1.8] *Suppose that W satisfies the standard assumptions, that X is a space, and that F is a homotopy fibre of $P_{\Sigma W} X \rightarrow P_W(X)$. Then $F \sim K(G, c(W))$ for some abelian group G which is concentrated at p .*

2.14. Definition. A map $X \rightarrow Y$ of pointed, connected spaces is a v_i^{-1} -equivalence if it induces an isomorphism on $v_i^{-1}\pi_*(-; V_i)$.

2.15. Proposition. [2, 11.5, 11.8] *Suppose that W satisfies the standard assumptions. Then for any X the map $X \rightarrow P_W(X)$ is a v_i^{-1} -equivalence for $1 \leq i < \kappa(W)$, whereas $P_W(X)$ is v_i -torsion for $i \geq \kappa(W)$.*

2.16. Remark. Note in connection with 2.15 that Bousfield works with a functor P_{v_n} which is nullification with respect to the suspension of a

particular complex of chromatic type n with minimal connectivity [2, 10.1]. However the arguments apply as stated to P_W .

3. THICK CLASSES

We are now ready to give a simple description of $\mathcal{T}(\mathcal{K}_W)$, at least when W satisfies the standard assumptions (2.9), and to prove 1.10.

3.1. Definition. A space X is said to be *almost killed by W* if $P_W(X)$ is concentrated at p and of finite Postnikov type. The class of all such spaces is denoted \mathcal{K}_W^+ .

3.2. Proposition. *If W satisfies the standard assumptions, then $\mathcal{T}(\mathcal{K}_W) = \mathcal{K}_W^+$.*

3.3. Proposition. *If W satisfies the standard assumptions, then \mathcal{K}_W^+ is the class of spaces which are concentrated at p , and v_i -torsion for $1 \leq i < \kappa(W)$.*

Proof of 1.10. By 2.1 and 2.3, it is enough to show that a space X is v_* -torsion if and only if X belongs to $\mathcal{T}(\mathcal{K}_W)$ for any finite complex W which is nontrivial mod p and is, say, a double suspension. Pick such a W , and let C be the homotopy cofibre of the degree p map $W \rightarrow W$. Then C satisfies the standard assumptions and $C \in \mathcal{B}_W$, so that $\mathcal{B}_C \subset \mathcal{B}_W$, $\mathcal{K}_C \subset \mathcal{K}_W$ and $\mathcal{T}(\mathcal{K}_C) \subset \mathcal{T}(\mathcal{K}_W)$. It is thus enough to show that X is v_* -torsion if and only if $X \in \mathcal{T}(\mathcal{K}_W)$ for any W which satisfies the standard assumptions. But for such a W , $\mathcal{T}(\mathcal{K}_W) = \mathcal{K}_W^+$ (3.2), and so the result follows immediately from 3.3 and the fact that there are finite complexes W with $\kappa(W)$ arbitrarily large [12] [13, 3.4]. \square

The rest of this section contains the proofs of 3.2 and 3.3.

3.4. Lemma. *If W satisfies the standard assumptions and $X \in \mathcal{K}_W^+$, then $P_W(X) \sim P_\nu(X)$, for $\nu = c(W) - 1$.*

Proof. By 2.12, $P_W(X) \sim P_W P_W(X)$ is equivalent to $P_\nu P_W(X)$. But by 2.5, $P_\nu P_W(X)$ is equivalent to $P_\nu(X)$. \square

3.5. Remark. It follows from 2.5 and 3.4 that if W satisfies the standard assumptions, a space belongs to \mathcal{K}_W^+ if and only if all of its components do.

3.6. Lemma. *If W satisfies the standard assumptions, then $\mathcal{T}(\mathcal{K}_W)$ contains all spaces which are concentrated at p and of finite Postnikov type.*

Proof. Since \mathcal{K}_W contains $*$ and $\mathcal{T}(\mathcal{K}_W)$ is closed under fibres, $\mathcal{T}(\mathcal{K}_W)$ is closed under taking loop spaces. Looping down Eilenberg-Mac Lane spaces (2.8) now shows that $\mathcal{T}(\mathcal{K}_W)$ contains discrete spaces of arbitrarily large cardinality, and taking fibres of maps between these discrete spaces shows that $\mathcal{T}(\mathcal{K}_W)$ contains all discrete spaces. For any X the homotopy fibres of $X \rightarrow \pi_0 X$ are the connected components of X ; since $\mathcal{T}(\mathcal{K}_W)$ is closed under extension by fibres, it is now clear then that X belongs to $\mathcal{T}(\mathcal{K}_W)$ if and only if all of its components do. It is thus enough to show that any connected space X of the indicated type belongs to $\mathcal{T}(\mathcal{K}_W)$. Since every Eilenberg-Mac Lane space $K(G, n)$ in which G is concentrated at p and $n \geq 1$ belongs to $\mathcal{T}(\mathcal{K}_W)$ (2.8), the desired result is easy to prove by an induction on the Postnikov tower of X (slightly refined in the case of $\pi_1 X$). \square

Proof of 3.2. Note by 2.13 that $\mathcal{K}_W^+ = \mathcal{K}_{\Sigma W}^+$. To obtain the inclusion $\mathcal{T}(\mathcal{K}_W) \subset \mathcal{K}_W^+$, it is enough to show that \mathcal{K}_W^+ is thick with respect to fibrations. By 3.5, it is sufficient to take a fibration sequence $F \rightarrow E \rightarrow B$ with B connected, and show that it satisfies the three conditions of 1.6. Consider the diagram

$$\begin{array}{ccccc} F & \longrightarrow & E & \longrightarrow & B \\ \downarrow & & \downarrow & & \downarrow \\ P_W(F) & \longrightarrow & E' & \longrightarrow & P_{\Sigma W}(B) \end{array}$$

from 2.7, in which $E \rightarrow E'$ is a P_W -equivalence and E' is ΣW -null. Say that a space Y is *good* if it is concentrated at p and has finite Postnikov type. Then $P_W(F)$ is good if and only if $F \in \mathcal{K}_W^+$; E' is good if and only if $E' \in \mathcal{K}_{\Sigma W}^+ = \mathcal{K}_W^+$, or, since $E \rightarrow E'$ is a P_W -equivalence, if and only if $E \in \mathcal{K}_W^+$; and $P_{\Sigma W}(B)$ is good if and only if $B \in \mathcal{K}_{\Sigma W}^+ = \mathcal{K}_W^+$. The desired result follows immediately from the long exact homotopy sequence of a fibration. Note that $P_W(F)$ is connected if and only if F is connected.

It remains to show that $\mathcal{K}_W^+ \subset \mathcal{T}(\mathcal{K}_W)$. Pick $X \in \mathcal{K}_W^+$, and consider the map $X \rightarrow P_W(X)$. By 2.6, the homotopy fibres of this map are in \mathcal{K}_W ; by 3.6, the target of the map is in $\mathcal{T}(\mathcal{K}_W)$. Since $\mathcal{T}(\mathcal{K}_W)$ is closed under extension by fibrations, $X \in \mathcal{T}(\mathcal{K}_W)$.

Proof of 3.3. By 3.5, we can assume that X is connected and show that X belongs to \mathcal{K}_W^+ if and only if it is concentrated at p and v_i -torsion for $1 \leq i \leq n$. This follows from results of Bousfield. Let $\nu = c(W) - 1$, and F the fibre of the Postnikov map $X \rightarrow P_\nu(X)$. By 2.10, 2.6, and 3.4, $X \in \mathcal{K}_W^+$ if and only if X is concentrated at p and $P_W(F) \sim *$.

Since the map $F \rightarrow X$ induces an isomorphism on $v_i^{-1}\pi_*(-; V_i)$ for all i , the result follows from 2.15. \square

4. RECOGNIZING v_* -TORSION SPACES

The aim of this section is to prove the following proposition, which gives a way of recognizing v_* -torsion spaces.

4.1. Proposition. *Suppose that W is a pointed space that satisfies the standard assumptions, and that X is a pointed connected space. Then the following three conditions are equivalent:*

- (1) X is v_* -torsion;
- (2) $\text{Map}_*^0(W, X)$ and $P_W(X)$ are v_* -torsion;
- (3) $\text{Cell}_W(X)$ and $P_W(X)$ are v_* -torsion.

This is a consequence of two more detailed statements, which depend on work of Bousfield and complement 2.15.

4.2. Proposition. *Suppose that W satisfies the standard assumptions, and that $f : X \rightarrow Y$ is a map of pointed connected spaces. If $i \geq \kappa(W)$, then $\text{Map}_*^0(W, f)$ is a v_i^{-1} -equivalence (2.14) if and only if f is a v_i^{-1} -equivalence. Moreover, $\text{Map}_*^0(W, X)$ is v_i -torsion for $i < \kappa(W)$.*

4.3. Proposition. *Suppose that W satisfies the standard assumptions and that X is a connected pointed space. Then $\text{Cell}_W(X) \rightarrow X$ is a v_i^{-1} -equivalence for $i \geq \kappa(W)$, whereas $\text{Cell}_W(X)$ is v_i -torsion for $i < \kappa(W)$.*

4.4. Remark. Taken together, 2.15 and 4.3 suggest that $\text{Cell}_W(X) \rightarrow X \rightarrow P_W(X)$ is a fibre sequence. This is not quite the case [7], but the sequence certainly acts like a fibre sequence as far as v_* -periodic homotopy is concerned.

Proof of 4.2. Suppose that $i \geq \kappa(W)$. Then $\kappa(V_i \wedge W) = i$; this comes from applying the Kunnetth formula for Morava K -theory, and using the fact that $\tilde{K}(i)_*W \neq 0$ if $i \geq \kappa(W)$ [13, 1.5.2]. It is also clear that any v_i -self-map of V_i induces a v_i -self-map of $V_i \wedge W$. By adjointness, then, there is an isomorphism

$$v_i^{-1}\pi_*(X; V_i \wedge W) \cong v_i^{-1}\pi_*(\text{Map}_*^0(W, X); V_i),$$

as well as a similar isomorphism for Y . Since the question of whether a map is a v_i^{-1} -equivalence does not depend upon the complex of type i used to compute the v_i -periodic homotopy groups, the result follows.

To see that the v_i -periodic homotopy of $\text{Map}_*^0(W, X)$ vanishes for $i < \kappa(W)$, consider the tower

$$V_i \wedge W \rightarrow \Sigma^d V_i \wedge W \rightarrow \Sigma^{2d} V_i \wedge W \rightarrow \dots$$

where $\Sigma^d V_i \rightarrow V_i$ is the v_i -self-map used to construct the v_i -periodic homotopy groups. By the Kunneth theorem, the maps in this tower induce the zero homomorphism on $\tilde{K}(n)_*$ for all $n \geq 1$. It follows from the nilpotence theorem [8] that the tower is pro-trivial (i.e. composites of sufficiently long chains of maps are null homotopic). Mapping the tower into X , however, and taking a colimit, gives the v_i -periodic homotopy of $\text{Map}_*^0(W, X)$. See the proof of [2, 11.4]) for another instance of this argument. \square

Proof of 4.3. The first statement is a consequence of 4.2, since the map $\text{Cell}_W(X) \rightarrow X$ induces an equivalence on $\text{Map}_*(W, -)$. The fact that $\text{Cell}_W(X)$ is v_i -torsion for $i < \kappa(W)$ follows from 3.3 together with the chain $\text{Cell}_W(X) \in \mathcal{B}_W \subset \mathcal{K}_W \subset \mathcal{K}_W^+$. \square

Proof of 4.1. Note that X is concentrated at p if and only if $P_W(X)$ is concentrated at p (2.10). The space $\text{Cell}_W(X)$ is always concentrated at p , since $P_W(\text{Cell}_W(X)) \sim *$; $\text{Map}_*^0(W, X)$ is concentrated at p by inspection. The first two conditions are now equivalent by 2.15 and 4.3; the first and third by combining 2.15 with an application of 4.2 to the map $X \rightarrow *$.

5. POLYGEMS

The purpose of this short section is to recall some properties of *polygems*, which are in some sense the simplest v_* -torsion spaces.

5.1. Definition. [9, 4.B] A *generalized Eilenberg-Mac Lane space*, or *GEM*, is a space which is equivalent to a (possibly infinite) product of Eilenberg-Mac Lane spaces whose homotopy groups are abelian.

5.2. Remark. A space X is a GEM if and only if X is equivalent to a topological abelian group.

5.3. Definition. The class of *polygems* is the smallest class of spaces which contains GEMs and is closed under fibres and under extension by fibrations.

5.4. Example. A connected space X of finite Postnikov type is a polygem if and only if $\pi_1(X)$ is solvable.

5.5. Remark. All of the components of a polygem are polygems. If X is a polygem and W is a pointed finite CW-complex, then for any choice of basepoint in X , the spaces $\text{Map}(W, X)$, $\text{Map}_*(W, X)$ and $\text{Map}_*^0(W, X)$ are also polygems.

It is easy to see that any GEM is v_i -torsion for $i \geq 1$. Since the class of spaces which are v_i -torsion for $i \geq 1$ is closed under fibres and

extension by fibrations (in fact it is thick with respect to fibrations), we immediately obtain the following proposition.

5.6. Proposition. *Any polygem X is v_i -torsion for $i \geq 1$. If in addition X is concentrated at p , then X is v_* -torsion.*

6. SIMPLICIAL SPACES

In this section we use some results from Bousfield and Friedlander [4] in order to prove the following two theorems, which identify certain spaces as polygms (§5).

6.1. Theorem. *Let X be a simplicial pointed topological space with the property that each space X_i is connected and has an abelian fundamental group. Suppose there is an integer m such that for any $i \geq 0$ and $j > m$, $\pi_j X_i$ is trivial. Then $|X|$ is a polygem.*

6.2. Theorem. *Let X be a simplicial pointed topological space with the property that each space X_i is connected and has an abelian fundamental group. Let A be a pointed finite-dimensional CW-complex which is equivalent to a suspension. Then the homotopy fibre of the natural map*

$$|\mathrm{Map}_*(A, X)| \rightarrow \mathrm{Map}_*(A, |X|)$$

is a polygem.

Since we will be using simplicial machinery from [4], we will prove these theorems in the case in which X is a simplicial pointed simplicial set and A is a finite-dimensional pointed simplicial set which is weakly equivalent to a suspension. The results as stated can be obtained by using the singular complex functor and the geometric realization functor to compare spaces to simplicial sets.

6.3. Warning. *For the remainder of this section, the word “space” will mean “simplicial set”.* In particular, a *simplicial space* is a *bisimplicial set* [4, §B] [10, IV].

Theorem 6.1 is a special case of a more general result. If X is a simplicial pointed space, then $\pi_j(X)$ ($j \geq 1$) is a simplicial group, and we let $\pi_i \pi_j(X)$ denote the i 'th homotopy group of this object.

6.4. Remark. There is a first quadrant spectral sequence with $E_{i,j}^2 = \pi_i \pi_j(X)$, which converges to $\pi_* |X|$ if each space X_i is connected [4, B.5].

6.5. Theorem. *Let X be a simplicial pointed space with the property that each space X_i is connected and has an abelian fundamental group.*

Suppose there is an integer m such that for any $i \geq 0$ and $j > m$, $\pi_i \pi_j X$ is trivial. Then $|X|$ is a polygem.

Proof. This is a corollary of the proof of [4, B.5]. According to the inductive fibration argument used in that proof, it is enough to show that if there is a single integer t such that for all $i \geq 0$, $\pi_j(X_i) = 0$ for $j \neq t$, then $|X|$ is a GEM. But, as indicated in the proof, in this case there is a natural bisimplicial map $X \rightarrow Y$, where Y is the simplicial space with Y_i given by the minimal Eilenberg-Mac Lane complex $K(\pi_t X_i, t)$; see also the discussion following [10, IV.4.12]. Since Y is a simplicial object in the category of simplicial abelian groups, its homotopy colimit, which is equivalent to its diagonal [4, B.1] [5, XII.3.4], is a simplicial abelian group and hence a GEM [10, III.2.20]. \square

The proof of 6.2 depends on the following result of Bousfield and Friedlander

6.6. Theorem. [4, B.4] [10, IV.4.9] *Suppose that $F \rightarrow E \rightarrow B$ is a sequence of simplicial spaces, with B a simplicial pointed space, and each B_i and E_i connected. Assume that for any $i \geq 0$ the sequence $F_i \rightarrow E_i \rightarrow B_i$ is a homotopy fibre sequence. Then $|F| \rightarrow |E| \rightarrow |B|$ is also a homotopy fibre sequence.*

Proof of 6.2. Let $M(-)$ stand for $\text{Map}_*^0(A, -)$, so that we have to study the homotopy fibre of the natural map $|MX| \rightarrow M|X|$. Suppose that $d = \dim(A)$. Let B_i be the Postnikov stage $P_d X_i$, and F_i the fibre of the map $X_i \rightarrow B_i$. Since each space F_i is d -connected, an induction over the skeleta of A using 6.6 shows that $|MF| \rightarrow M|F|$ is an equivalence. There is a commutative diagram

$$(6.7) \quad \begin{array}{ccccc} |MF| & \longrightarrow & |MX| & \longrightarrow & |MB| \\ \sim \downarrow & & \downarrow & & \downarrow \\ M|F| & \longrightarrow & M|X| & \longrightarrow & M|B| \end{array}$$

in which both rows are fibration sequence by 6.6. Note in this connection that because of the connectivity of F_i , $MF_i = \text{Map}_*(A, F_i)$ and $MF_i \rightarrow MX_i \rightarrow MB_i$ is a fibration sequence. Similar remarks apply to the bottom row, since $|F|$ is d -connected [10, IV.4.13]. It follows that the right hand square in 6.7 is a homotopy fibre square, and so the homotopy fibre of $|MX| \rightarrow M|X|$ is the same as the homotopy fibre of $|MB| \rightarrow M|B|$. Since A is a suspension, the fundamental groups of the spaces B_i are abelian, and so MB satisfies the assumptions of 6.1 and $|MB|$ is a polygem. Similarly, B satisfies the assumptions of 6.1, and so $|B|$ and hence $M|B|$ are polygems (5.5). It follows that the homotopy fibre of $|MB| \rightarrow M|B|$ is also a polygem.

7. CLOSURE AND RESOLUTION

In this section we prove 1.14 and 1.15. The proof of 1.14 depends on the following slightly more elaborate result.

7.1. Proposition. *Suppose that W satisfies the standard assumptions (2.9), and that X is a simplicial space with the property that for any $i \geq 0$, the space X_i is connected, has an abelian fundamental group, and belongs to \mathcal{K}_W^+ . Then the realization $|X|$ also belongs to \mathcal{K}_W^+ .*

Proof. Choosing a basepoint in X_0 and propagating it to the other spaces X_i with degeneracy maps converts X into a simplicial pointed space. Now apply the functor P_W in each simplicial degree, and let F_i be the fibre of $X_i \rightarrow P_W(X_i)$, so that $F \rightarrow X \rightarrow P_W(X)$ is a sequence of simplicial spaces which in each degree is a fibration sequence. By 2.5 and 6.6, $|F| \rightarrow |X| \rightarrow |P_W(X)|$ is a homotopy fibre sequence of spaces. By 3.4 and 5.6, $P_W(X_i)$ has trivial homotopy in dimensions $\geq c(W)$, and so (6.1) $|P_W(X)|$ is a polygem. It is clear from 6.4 that $|P_W(X)|$ is concentrated at p , and so this space belongs to \mathcal{K}_W^+ . For any i the space F_i is killed by W (2.6), and hence $|F|$ is killed by W [2, 2.5] and belongs to \mathcal{K}_W^+ . Since $\mathcal{K}_W^+ = \mathcal{T}(\mathcal{K}_W)$ (3.2) and so is closed under extension by fibrations, $|X| \in \mathcal{K}_W^+$. \square

Proof of 1.14. Note that as in the proof of 1.10, a space X is v_* -torsion if and only if it belongs to \mathcal{K}_W^+ for each space W which satisfies the standard conditions. Now apply 7.1. \square

Proof of 1.15. Let $M(-) = \text{Map}_*^0(W, -)$, $R = R_W(X)$, and $C = \text{Cell}_W(X)$. Then R is a simplicial pointed space which is augmented by a map to X and in each degree is equivalent to a wedge of suspensions of W . Essentially by construction [14] [1], the natural map $|M(R)| \rightarrow M(X)$ is a weak equivalence; in fact R is designed so that the homotopy spectral sequence (6.4) for $\pi_*|M(R)|$ collapses to $\pi_*M(X)$. Consider the natural sequence

$$|M(R)| \rightarrow M(|R|) \rightarrow M(X).$$

in which the composite is an equivalence. Let F_1 be the fibre of the left hand map and F_2 the fibre of the right hand map. Then $F_1 \sim \Omega F_2$, and F_1 is a polygem (6.2). In particular F_1 is v_i -torsion for $i \geq 1$ (5.6), so the same is true of F_2 . It follows that $M(|R|) \rightarrow M(X)$ is a v_i^{-1} -equivalence for $i \geq 1$ (2.14). The space $|R|$ is built from W , and so the map $|R| \rightarrow X$ lifts canonically (up to homotopy) to a map $|R| \rightarrow C$. Since $C \rightarrow X$ induces an equivalence $M(C) \rightarrow M(X)$, it's clear that the map $M(|R|) \rightarrow M(C)$ is a v_i^{-1} -equivalence for $i \geq 1$; hence (4.2) $|R| \rightarrow C$ is a v_i^{-1} -equivalence for $i \geq \kappa(W)$. However

$|R| \rightarrow C$ is trivially a v_i^{-1} -equivalence for $i < \kappa(W)$, since both of these spaces are built from, hence killed by, W , and consequently are v_i -torsion for $1 \leq i < \kappa(W)$ (3.3). It follows that the fibre F of $|R| \rightarrow C$ is concentrated at p and is v_i -torsion for $i \geq 1$; in other words, F is v_* -torsion. \square

REFERENCES

- [1] David Blanc, *Mapping spaces and M - CW complexes*, Forum Math. **9** (1997), no. 3, 367–382. MR **98i**:55015
- [2] A. K. Bousfield, *Localization and periodicity in unstable homotopy theory*, J. Amer. Math. Soc. **7** (1994), no. 4, 831–873. MR **95c**:55010
- [3] ———, *Unstable localization and periodicity*, Algebraic topology: new trends in localization and periodicity (Sant Feliu de Guíxols, 1994), Progr. Math., vol. 136, Birkhäuser, Basel, 1996, pp. 33–50. MR **98c**:55014
- [4] A. K. Bousfield and E. M. Friedlander, *Homotopy theory of Γ -spaces, spectra, and bisimplicial sets*, Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II, Lecture Notes in Math., vol. 658, Springer, Berlin, 1978, pp. 80–130. MR **80e**:55021
- [5] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Springer-Verlag, Berlin, 1972, Lecture Notes in Mathematics, Vol. 304. MR 51 #1825
- [6] W. Chachólski, W. G. Dwyer, and M. Interfont, *The A -complexity of a space*, J. London Math. Soc. (2) **65** (2002), no. 1, 204–222. MR **2002j**:55009
- [7] Wojciech Chachólski, *On the functors CW_A and P_A* , Duke Math. J. **84** (1996), no. 3, 599–631. MR **97i**:55023
- [8] Ethan S. Devinatz, Michael J. Hopkins, and Jeffrey H. Smith, *Nilpotence and stable homotopy theory. I*, Ann. of Math. (2) **128** (1988), no. 2, 207–241. MR **89m**:55009
- [9] Emmanuel Dror Farjoun, *Cellular spaces, null spaces and homotopy localization*, Lecture Notes in Mathematics, vol. 1622, Springer-Verlag, Berlin, 1996. MR **98f**:55010
- [10] Paul G. Goerss and John F. Jardine, *Simplicial homotopy theory*, Progress in Mathematics, vol. 174, Birkhäuser Verlag, Basel, 1999. MR **2001d**:55012
- [11] Michael J. Hopkins and Jeffrey H. Smith, *Nilpotence and stable homotopy theory. II*, Ann. of Math. (2) **148** (1998), no. 1, 1–49. MR **99h**:55009
- [12] Stephen A. Mitchell, *Finite complexes with $A(n)$ -free cohomology*, Topology **24** (1985), no. 2, 227–246. MR **86k**:55007
- [13] Douglas C. Ravenel, *Nilpotence and periodicity in stable homotopy theory*, Annals of Mathematics Studies, vol. 128, Princeton University Press, Princeton, NJ, 1992, Appendix C by Jeff Smith. MR **94b**:55015
- [14] Christopher R. Stover, *A van Kampen spectral sequence for higher homotopy groups*, Topology **29** (1990), no. 1, 9–26. MR **91h**:55011

YALE UNIVERSITY, NEW HAVEN, CT 06520 USA
E-mail address: chacholskiwojciech@math.yale.edu

UNIVERSITY OF NOTRE DAME, NOTRE DAME IN 46556 USA
E-mail address: `dwyer.1@nd.edu`

KALAMAZOO COLLEGE, KALAMAZOO MI, 49006 USA
E-mail address: `intermon@kzoo.edu`