

Section 4.1 Numerical Differentiation

Example 4.4.1 Use forward difference formula with $h = 0.1$ to approximate the derivative of $f(x) = \ln(x)$ at $x_0 = 1.8$. Determine the bound of the approximation error.

1st derivative approximation (obtained by Lagrange interpolation)

The interpolation points are given as:


$$(x_0, f(x_0))$$

$$(x_1, f(x_1))$$

$$(x_2, f(x_2))$$

...

$$(x_N, f(x_N))$$



By Lagrange Interpolation Theorem (**Thm 3.3**):

$$f(x) = \sum_{k=0}^n f(x_k) L_{N,k}(x) + \frac{(x-x_0) \cdots (x-x_N)}{(N+1)!} f^{(N+1)}(\xi(x)) \quad (1)$$

Take 1st derivative for Eq. (1):

$$f'(x) = \sum_{k=0}^n f(x_k) L'_{N,k}(x) + \frac{(x-x_0) \cdots (x-x_N)}{(N+1)!} \left(\frac{d \left(f^{(N+1)}(\xi(x)) \right)}{dx} \right) + \frac{1}{(N+1)!} \left(\frac{d \left((x-x_0) \cdots (x-x_N) \right)}{dx} \right) f^{(N+1)}(\xi(x))$$

Set $x = x_j$, with x_j being x-coordinate of one of interpolation nodes. $j = 0, \dots, N$.

$$f'(x_j) = \sum_{k=0}^n f(x_k) L'_{N,k}(x_j) + \frac{f^{(N+1)}(\xi(x_j))}{(N+1)!} \prod_{\substack{k=0; \\ k \neq j}}^N (x_j - x_k) \text{ ---- (N+1)-}$$

point formula to approximate $f'(x_j)$ **(4.2)**

The error of (N+1)-point formula is $\frac{f^{(N+1)}(\xi(x_j))}{(N+1)!} \prod_{\substack{k=0; \\ k \neq j}}^N (x_j - x_k)$.

Remark: $f'(x_j) \approx \sum_{k=0}^n f(x_k) L'_{N,k}(x_j)$


Example. Derive the three-point formula with error to approximate $f'(x_j)$.

Let interpolation nodes be $(x_0, f(x_0))$, $(x_1, f(x_1))$ and $(x_2, f(x_2))$.

$$\begin{aligned} f'(x_j) = & f(x_0) \left[\frac{2x_j - x_1 - x_2}{(x_0 - x_1)(x_0 - x_2)} \right] + f(x_1) \left[\frac{2x_j - x_0 - x_2}{(x_1 - x_0)(x_1 - x_2)} \right] \\ & + f(x_2) \left[\frac{2x_j - x_0 - x_1}{(x_2 - x_0)(x_2 - x_1)} \right] + \frac{f^{(3)}(\xi(x_j))}{6} \prod_{\substack{k=0; \\ k \neq j}}^2 (x_j - x_k) \end{aligned}$$

Mostly used three-point formula (see Figure 1)

Let x_0, x_1 , and x_2 be **equally spaced** and the grid spacing be h .



Thus $x_1 = x_0 + h$; and $x_2 = x_0 + 2h$.

$$1. \quad f'(x_0) = \frac{1}{2h} [-3f(x_0) + 4f(x_1) - f(x_2)] + \frac{h^2}{3} f^{(3)}(\xi(x_0)) \quad (\text{three-point endpoint formula}) \quad (4.4)$$

$$2. \quad f'(x_1) = \frac{1}{2h} [-f(x_0) + f(x_2)] + \frac{h^2}{6} f^{(3)}(\xi(x_1)) \quad (\text{three-point midpoint formula}) \quad (4.5)$$

$$3. \quad f'(x_2) = \frac{1}{2h} [f(x_0) - 4f(x_1) + 3f(x_2)] + \frac{h^2}{3} f^{(3)}(\xi(x_2)) \quad (\text{three-point endpoint formula}) \quad (4.4.1)$$


Remark: Eq. (4.4) in textbook is:

$$f'(x_0) = \frac{1}{2h} [-3f(x_0) + 4f(x_0 + h) - f(x_0 + 2h)] + \frac{h^2}{3} f^{(3)}(\xi(x_0))$$

h can be both positive and negative


Mostly used five-point formula

1. Five-point midpoint formula



$$f'(x_0) = \frac{1}{12h} [f(x_0 - 2h) - 8f(x_0 - h) + 8f(x_0 + h) - f(x_0 + 2h)] + \frac{h^4}{30} f^{(5)}(\xi) \quad (4.6)$$

2. Five-point endpoint formula




$$f'(x_0) = \frac{1}{12h} [-25f(x_0) + 48f(x_0 + h) - 36f(x_0 + 2h) + 16f(x_0 + 3h) - 3f(x_0 + 4h)] + \frac{h^4}{5} f^{(5)}(\xi) \quad (4.7)$$

Example 4.1.2 Values for $f(x) = xe^x$ are given in the following table. Use all applicable 3-point and 5-point formulas to approximate $f'(2.0)$.

x	1.8	1.9	2.0	2.1	2.2
$f(x)$	10.889365	12.703199	14.778112	17.148957	19.855030

2nd derivative approximation (obtained by Taylor polynomial)



Approximate $f(x_0 + h)$ by expansion about x_0 :

$$f(x_0 + h) = f(x_0) + f'(x_0)h + \frac{1}{2}f''(x_0)h^2 + \frac{1}{6}f'''(x_0)h^3 + \frac{1}{24}f^{(4)}(\xi_1)h^4 \quad (3)$$

Approximate $f(x_0 - h)$ by expansion about x_0 :

$$f(x_0 - h) = f(x_0) - f'(x_0)h + \frac{1}{2}f''(x_0)h^2 - \frac{1}{6}f'''(x_0)h^3 + \frac{1}{24}f^{(4)}(\xi_2)h^4 \quad (4)$$

Add Eqns. (3) and (4): $f(x_0 - h) + f(x_0 + h) = 2f(x_0) + f''(x_0)h^2 + \left[\frac{1}{24}f^{(4)}(\xi_1)h^4 + \frac{1}{24}f^{(4)}(\xi_2)h^4\right]$

Thus

Second derivative midpoint formula

$$f''(x_0) = \frac{1}{h^2} [f(x_0 - h) - 2f(x_0) + f(x_0 + h)] - \frac{h^2}{12} f^{(4)}(\xi)$$

Example 3. Values for $f(x) = xe^x$ are given in the following table. Use second derivative approximation formula to approximate $f''(2.0)$.

x	1.8	1.9	2.0	2.1	2.2
$f(x)$	10.889365	12.703199	14.778112	17.148957	19.855030

Solution: $f''(2.0) \approx \frac{1}{(0.1)^2} [f(1.9) - 2f(2.0) + f(2.1)] = 29.5932$

Or

$$f''(2.0) \approx \frac{1}{(0.1)^2} [f(1.8) - 2f(2.0) + f(2.2)] = 29.704275$$